

Gaussian integration formulas for logarithmic weights and application to 2-dimensional solid-state lattices.

Alphonse P. Magnus
Université catholique de Louvain,
Institut de mathématique pure et appliquée,
2 Chemin du Cyclotron,
B-1348 Louvain-La-Neuve, Belgium
alphonse.magnus@uclouvain.be

The 1st version: April 8, 2015 (incomplete and unfinished)
2nd version: Aug 20, 2016 (incomplete and unfinished)
3rd version: Dec 8, 2016 (incomplete and unfinished)
4th version: Apr 11, 2017 (incomplete and unfinished)
This version: November 18, 2017(incomplete and unfinished)

*Nothing exists per se except atoms and the void
... however solid objects seem,*

Lucretius, *On the Nature of Things*,
Translated by William Ellery Leonard

Abstract. The making of Gaussian numerical integration formulas is considered for weight functions with logarithmic singularities. Chebyshev modified moments are found most convenient here. The asymptotic behavior of the relevant recurrence coefficients is stated in two conjectures. The relation with the recursion method in solid-state physics is summarized, and more details are given for some two-dimensional lattices (square lattice and hexagonal (graphene) lattice).

Keywords: Numerical analysis, Integration formulas, Logarithmic singularities, Solid-state physics, 2-dimensional lattices, Graphene.

2010 Mathematics Subject Classification:

42A16 Fourier coefficients, Fourier series of functions with special properties, special Fourier series, 42C05 Orthogonal functions and polynomials, general theory, 42C10 Fourier series in special orthogonal functions (Legendre polynomials, etc.), 47B36 Jacobi (tridiagonal) operators (matrices) and generalizations, 65D32 Quadrature and cubature formulas, 81Q05 Closed and approximate solutions to the Schrödinger eq. etc.

MSC[2010]42A16, 42C05, 42C10, 47B36, 65D32, 81Q05

CONTENTS

1. Orthogonal polynomials and Gaussian quadrature formulas.	2
2. Power moments and recurrence coefficients.	3
2.1. Recurrence coefficients and examples.	3
2.2. Asymptotic behavior of recurrence coefficients.	5
3. Modified moments.	5
3.1. Main properties and numerical stability.	5
3.2. Legendre examples	6
3.3. Chebyshev examples	7
3.4. The algorithm.	8
4. Expansions in functions of the second kind.	8
4.1. Theorem.	9
4.2. Chebyshev functions of the second kind.	10

4.3. Corollary.	10
5. Weights with logarithmic singularities. Interior singularity.	10
5.1. Known results.	10
5.2. Conjecture.	11
5.3. About the Szegő asymptotic formula.	12
6. Weights with logarithmic singularities. Endpoint singularity.	18
6.1. Numerical tests.	18
6.2. Conjecture.	18
6.3. Trying multiple orthogonal polynomials.	20
7. About matrices in solid-state physics.	21
7.1. Matrix approximation of the Hamiltonian operator.	21
7.2. Density of states.	23
7.3. The recursion (Lanczos) method.	24
7.4. Perfect crystals.	24
8. Two famous 2-dimensional lattices.	25
8.1. The square lattice.	25
8.2. Hexagonal lattice: <i>graphene</i>	27
9. Acknowledgements.	33
References	33

1. ORTHOGONAL POLYNOMIALS AND GAUSSIAN QUADRATURE FORMULAS.

Let μ be a positive measure on a real interval $[a, b]$, and P_n the related monic orthogonal polynomial of degree n , i.e., such that

$$P_n(x) = x^n + \cdots, \quad \int_a^b P_n(t)P_m(t)d\mu(t) = 0, m \neq n, \quad n = 0, 1, \dots \quad (1)$$

An enormous amount of work has been spent since about 200 years on the theory and the applications of these functions. One of their most remarkable properties is the recurrence relation

$$P_{n+1}(x) = (x - b_n)P_n(x) - a_n^2 P_{n-1}(x), \quad n = 1, 2, \dots, \quad (2)$$

with $P_0(x) \equiv 1, P_1(x) = x - b_0$. See, among numerous other sources, books by Chihara [17], Gautschi [39, 41], Ismail [55], chap. 18 of NIST handbook [89], and other surveys [44, 69, 70].

Orthogonal polynomials are critically involved in the important class of Gaussian integration formulas. A classical integration formula $\int_a^b f(t)d\mu(t) \approx w_1 f(x_1) + \cdots + w_N f(x_N)$ (Newton-Cotes, Simpson, etc.) is the integral $\int_a^b p(t) d\mu(t)$ of the polynomial interpolant p of f at the points x_1, \dots, x_N . Interpolation errors can sometimes become quite wild, to the opposite of least squares approximations made with a polynomial q of degree N , achieving

$$\min_q \int_a^b (f(t) - q(t))^2 d\mu(t) \quad \Rightarrow \quad \int_a^b (f(t) - q(t))r(t) d\mu(t) = 0$$

for any polynomial r of degree $< N$. We want the favorable aspects of both sides! i.e., easy use of numerical integration formulas, and safety of least squares approximation. Take at least for f a polynomial of degree N , say, $f(t) = t^N$, see that $f - p$ vanishes at x_1, \dots, x_N and will be orthogonal to all polynomials of degree $< N$ if it is a constant times P_N , so if x_1, \dots, x_N are the zeros of P_N . All least squares problems are then satisfactorily solved with the discrete scalar product $(f, g)_N = \sum_1^N w_j f(x_j)g(x_j)$. See Davis & Rabinowitz [22, § 2.7], Boyd [11, chap. 4] for this discussion.

Approximate integration formulas are not only used in the area or volume calculations from time to time, they are also used massively in pseudospectral solutions of big partial derivative equations and other functional equations. As an example of numerical procedure, a polynomial approximation to the solution of a functional equation $F(u) = 0$ is determined by orthogonality conditions $\int_a^b F(u(t))r(t) d\mu(t) = 0$ for any polynomial r of degree $< N$ (Galerkin method), where the integral is replaced by its Gaussian formula $(F(u), r)_N = 0$. See for instance Boyd [11, chap. 3, 4], Fornberg [32, § 4.7], Mansell & al. [75], Shizgal [98].

2. POWER MOMENTS AND RECURRENCE COEFFICIENTS.

2.1. Recurrence coefficients and examples. Let us consider the generating function of the moments μ_n , which is called here the Stieltjes function of the measure $d\mu$

$$S(x) = \int_a^b \frac{d\mu(t)}{x-t} = \frac{\mu_0}{x} + \frac{\mu_1}{x^2} + \dots, \quad x \notin [a, b], \quad \mu_n = \int_a^b t^n d\mu(t). \quad (3)$$

Sometimes, S is called the Stieltjes transform of $d\mu$, but technically, the Stieltjes transform of a measure is the integral of $(x+t)^{-1}d\mu(t)$ on the positive real line [52, chap. 12]. For measures on the whole real line, one should use the name ‘‘Hamburger transform’’. P. Henrici [53, §14.6] speaks of ‘‘Cauchy integrals on straight line segments’’, Van Assche [103] calls S ‘‘Stieltjes transform’’ for (3) in all cases. ‘‘Markov function’’ is also used [4, 44] when it is clear that there is no danger of confusion with random processes.

The power expansion (3) is an asymptotic expansion. If $[a, b]$ is finite, the expansions converges when $|x| > \max(|a|, |b|)$.

The function S is also the first function of the second kind [5, 6]

$$Q_n(x) = \int_a^b \frac{P_n(t) d\mu(t)}{x-t}.$$

The recurrence relation (2) holds for the Q_n s too. Indeed,

$$Q_{n+1}(x) = \int_a^b \frac{[(t-b_n = t-x+x-b_n)P_n(t) - a_n^2 P_{n-1}(t)] d\mu(t)}{x-t}$$

$= -\mu_0 \delta_{n,0} + (x-b_n)Q_n(x) - a_n^2 Q_{n-1}(x)$. At $n=0$, $Q_1(x) - (x-b_0)Q_0(x) + \mu_0 = 0$. We have

$$\frac{Q_n(x)}{Q_{n-1}(x)} = \frac{a_n^2}{x-b_n - \frac{Q_{n+1}(x)}{Q_n(x)}}$$

[38, eq.(5.1-5.2)], [42, eq.(2.15)], and

$$S(x) = Q_0(x) = \frac{\mu_0}{x-b_0 - \frac{a_1^2}{x-b_1 - \dots - \frac{Q_{n+1}(x)}{Q_n(x)}}},$$

[107, eq. (59.2)]. For bounded $[a, b]$, the continued fraction converges for all $x \notin [a, b]$ [38, 53, 107].

Some examples, which will be inspiring later on, are

$$S(x) = \frac{1}{2} \int_{-1}^1 \frac{dt}{x-t} = \frac{1}{2} \log \frac{x+1}{x-1} = \frac{1}{x} + \frac{1}{3x^3} + \frac{1}{5x^5} + \dots \quad (4)$$

$$S(x) = \int_{-1}^1 \frac{|t|dt}{x-t} = \int_0^1 \frac{2txdt}{x^2-t^2} = x \log \frac{x^2}{x^2-1} = \frac{1}{x} + \frac{1}{2x^3} + \frac{1}{3x^5} + \dots \quad (5)$$

This shows how logarithmic singularities are often seen in Stieltjes functions.

Here is a case with an explicit logarithmic singularity in the weight function

$$S(x) = - \int_0^1 \frac{\log t \, dt}{x-t} = \text{Li}_2(x^{-1}) = \sum_1^{\infty} \frac{1}{n^2 x^n}, \quad (6)$$

where Li_2 is the dilogarithm function [89, §25.12].

A last example with Euler's Beta function:

$$S(x) = \int_0^1 \frac{t^{q-1}(1-t)^{p-q} \, dt}{x-t} = \frac{\mu_0}{x} + \frac{\mu_1}{x^2} + \frac{\mu_2}{x^3} + \dots, \quad (7)$$

$$\mu_n = B(n+q, p-q+1) = \frac{\Gamma(n+q)\Gamma(p-q+1)}{\Gamma(n+p+1)}$$

The recurrence relation (2) is needed in various applications, whence the importance of getting the recurrence coefficients (Lanczos constants) from the moments μ_n (Schwarz constants, see [19] for these names). Some of our examples have been solved in the past, see the results in Table 1.

	(4) Legendre	(5) mod. Jacobi Chihara [17, chap. 5, § 2 (G)]	(7) Jacobi on (0, 1) Abramowitz [1, § 22.2.2, § 22.7.2]
a_n^2	$\frac{n^2}{4n^2-1}$	$\frac{2n+1-(-1)^n}{4(2n+1+(1)^n)}$	$\frac{n(n+p-1)(n+q-1)(n+p-q)}{(2n+p-2)(2n+p-1)^2(2n+p)}$
b_n	0	0	$\frac{2n(n+p)+q(p-1)}{(2n+p+1)(2n+p-1)}$

TABLE 1. Some known recurrence coefficients formulas.

General formulas for the recurrence coefficients from the power moments follow from the set of linear equations $\sum_0^{n-1} \mu_{i+j} c_j^{(n)} = -\mu_{i+n}$, $i = 0, \dots, n-1$ for the coefficients $c_j^{(n)}$ of $P_n(x) = x^n + \sum_0^{n-1} c_j^{(n)} x^j$, yielding $b_0 + \dots + b_{n-1} = -c_{n-1}^{(n)}$ and $\mu_0 a_1^2 \dots a_n^2 = D_{n+1}/D_n$, where D_n is the determinant of the stated set of equations (Hankel determinant). Various algorithms organize the progressive construction of the recurrence coefficients from the power moments but have an enormous condition number for large degree, whence the importance of alternate numerical methods [39], which will be considered in next section.

In some serendipitous cases, as seen in Table 1, closed-form formulas have been found [17, chapters 5 and 6] [89, § 18.3-18.37].

n	0	1	2	3	4	5
a_n^2		$\frac{7}{144}$	$\frac{647}{11025}$	$\frac{71180289}{1172105200}$	$\frac{332349955856}{5405644687527}$	$\frac{39672481023099631594375}{641525900508218274561936}$
b_n	$\frac{1}{4}$	$\frac{13}{28}$	$\frac{8795}{18116}$	$\frac{124351943}{252694908}$	$\frac{43450203422161}{87773135347044}$	$\frac{23506086742557104854013941}{47335997944735259180626044}$

TABLE 2. First recurrence coefficients for the weight $-\log t$ on (0, 1).

No formula is known for the dilogarithm case (6), and nothing simple must be expected, as seen from the first ones in table 2, whose asymptotic behavior will be investigated in table 7; see also [80].

2.2. Asymptotic behavior of recurrence coefficients. Asymptotic behavior of a_n and b_n has been enormously investigated. The simplest, and most meaningful, result is that, if the derivative $w = \mu'$ of the absolutely continuous¹ part is positive a.e. on (a, b) , then

$$a_n \rightarrow a_\infty = \frac{b-a}{4}, \quad b_n \rightarrow b_\infty = \frac{a+b}{2}, \quad n \rightarrow \infty. \quad (8)$$

This seemingly simple result took decades to receive a complete proof, see the surveys by D.S. Lubinsky [69, §3.2], P. Nevai [84, §4.5], [85], and Van Assche's book [103, §2.6] for accurate statements and story.

A closer look to the Jacobi recurrence coefficients (7), Table 1 gives $a_n = \frac{1}{4} - \frac{(q-1)^2 + (p-q)^2 - 1/2}{16n^2} + o(n^{-2})$, $b_n = \frac{1}{2} + \frac{(q-1)^2 - (p-q)^2}{8n^2} + o(n^{-2})$.

For a general interval (a, b) , the Jacobi weight is $(b-x)^\alpha(x-a)^\beta$, and the relevant asymptotic behavior is

$$\begin{aligned} a_n &= \frac{b-a}{4} \left(1 - \frac{\alpha^2 + \beta^2 - 1/2}{4n^2} + o(n^{-2}) \right), \\ b_n &= \frac{a+b}{2} - \frac{(b-a)(\alpha^2 - \beta^2)}{8n^2} + o(n^{-2}). \end{aligned} \quad (9)$$

This behavior is thought to be present for all weights behaving like powers near the support's endpoints. Interior singularities create wilder oscillating perturbations, as it will be recalled later on. Lambin and Gaspard [63, Appendix] made interesting numerical tests on problems of solid-state physics by reducing the oscillating terms through sums and products, their formulas² are:

$$a_1 \cdots a_n = \text{constant} \times \left(\frac{b-a}{4} \right)^n \left(1 + \frac{\alpha^2 + \beta^2 - 1/2}{4n} + o(1/n) \right),$$

$b_0 + \cdots + b_n = n \frac{a+b}{2} + \text{constant} + \frac{(b-a)(\alpha^2 - \beta^2)}{8n} + o(1/n)$. The constants are known from Szegő's theory, see § 5.3. Complete expansions in powers of $1/n$ have been established when the weight function is $(b-x)^\alpha(x-a)^\beta$ times a positive analytic function on $[a, b]$ [62, Thm. 1.10], quite a strong condition. Perturbation of a Jacobi weight is considered by Nevai and Van Assche [86, § 5.2], but their trace-class condition $\sum |a_n - a_\infty| + |b_n - b_\infty| < \infty$ is rather strong too, as we will encounter expected $O(1/n)$ and $O(1/(n \log n))$ perturbations. See also L.Lefevre et al. [66] for more applications with Jacobi polynomials.

Many more refinements will be studied in § 5 and 6.

3. MODIFIED MOMENTS.

A very efficient technique for computing large numbers of recurrence coefficients is described here.

3.1. Main properties and numerical stability. We consider a sequence of polynomials $\{R_0, R_1, \dots\}$ with R_n of degree n . Here, R_n need not be monic. The related modified moment of degree n is defined as

$$\nu_n = \int_a^b R_n(t) d\mu(t). \quad (10)$$

We want to compute the recurrence relation coefficients (2) from the modified moments of $d\mu$. The algebraic contents of the problem is the same as before, as each modified moment is a finite linear combination of the power moments, but the numerical accuracy in finite precision can be

¹ $d\mu = d\mu_{\text{absolutely continuous}} + d\mu_{\text{singular}}$.

²The product $b_1 \cdots b_n$ of [63] is here $a_1^2 \cdots a_n^2$.

strongly enhanced: with the notation (f, g) for the scalar product $\int_a^b f(x)g(x) d\mu(x)$, we again compute the values $(P_n, R_j), n, j = 0, 1, \dots, N-1$ by

$$\begin{aligned} \mathbf{G}_N &= \begin{bmatrix} (R_0, R_0) & \cdots & (R_0, R_{N-1}) \\ \vdots & \vdots & \vdots \\ (R_{N-1}, R_0) & \cdots & (R_{N-1}, R_{N-1}) \end{bmatrix} \\ &= \begin{bmatrix} (R_0, P_0) & 0 & \cdots & 0 \\ (R_1, P_0) & (R_1, P_1) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ (R_{N-1}, P_0) & (R_{N-1}, P_1) & \cdots & (R_{N-1}, P_{N-1}) \end{bmatrix} \begin{bmatrix} 1/\|P_0\|^2 & & & \\ & 1/\|P_1\|^2 & & \\ & & \ddots & \\ & & & 1/\|P_{N-1}\|^2 \end{bmatrix} \\ &\quad \begin{bmatrix} (P_0, R_0) & (P_0, R_1) & \cdots & (P_0, R_{N-1}) \\ 0 & (P_1, R_1) & \cdots & (P_1, R_{N-1}) \\ & & \ddots & \vdots \\ 0 & 0 & \cdots & (P_{N-1}, R_{N-1}) \end{bmatrix} \end{aligned} \quad (11)$$

Where the left-hand side is the Gram matrix of the basis $\{R_0, \dots, R_{N-1}\}$, factored in (11) as a lower triangular matrix times a diagonal matrix times an upper triangular matrix which happens to be the transposed of the first factor. The equation (11) is the matrix writing of the Gaussian (!) elimination method, also known for a positive definite matrix as Cholesky's method [12, 13]. See also Bultheel & Van Barel [15, § 4.2] for this connection of the Gram-Schmidt method with modified moments.

The numerical stability of the computation of the factors of the right-hand side of (11) is measured by the condition number of the matrix \mathbf{G}_N , which is the ratio of the extreme eigenvalues of the matrix (for a general nonsymmetric matrix, singular values must be considered [45, 109]), after a convenient scaling replacing $R_n(x)$ by $R_n(x)/\rho_n$.

The extreme eigenvalues are easily seen as the inf and sup on real vectors $[c_0, \dots, c_{N-1}]$ of the ratio $\frac{\sum_j \sum_k c_j c_k (R_j/\rho_j, R_k/\rho_k) = \int_a^b p^2(x) d\mu(x)}{\sum_j c_j^2}$, where $p(x) = \sum_j c_j R_j(x)/\rho_j$ (Rayleigh quotient [109, §54]). Now, in the important special case where the R_n/ρ_n s are the orthonormal polynomials with respect to a measure $d\mu_R$ with the same support as $d\mu$, the extreme eigenvalues are the inf and sup on the real polynomials p of degree $< N$ of $\frac{\int p^2(x) d\mu(x)}{\int_a^b p^2(x) d\mu_R(x)}$ so, these eigenvalues remain bounded and bounded from below if $d\mu(x)/d\mu_R(x)$ is similarly bounded [8].

3.2. Legendre examples.

Let $R_n(x) = P_n\left(\frac{2x-a-b}{b-a}\right)$. From tables and formulas of Legendre polynomials [1, 89] etc., one has $R_0 = 1, R_1(x) = (2x-a-b)/(b-a), R_{n+1}(x) = [(2n+1)(2x-a-b)R_n(x)/(b-a) - nR_{n-1}(x)]/(n+1), R_n(a) = (-1)^n, R_n(b) = 1, \|R_n\|_R^2 = \int_a^b R_n^2(x) dx = (b-a)/(2n+1)$. The integral of R_n is of special interest, it is $\int_a^x R_n(t) dt = (b-a)(R_{n+1}(x) - R_{n-1}(x))/(2(2n+1))$ when $n > 0$ [32, p.157].

For $-\log t$ on $(0, 1)$, the modified moments are $\nu_0 = 1, \nu_n = -\int_0^1 R_n(t) \log t dt$
 $= \int_0^1 \frac{R_{n+1}(t) - R_{n-1}(t)}{2(2n+1)t} dt = -\int_0^1 \frac{R_n(t) + R_{n-1}(t)}{2(n+1)t} dt = \frac{(-1)^n}{n(n+1)}, n = 1, 2, \dots$ [89, 14.18.6
 Christoffel Darboux], also a special case of Jacobi polynomials formulas by Gautschi [40, eq. (16)].

For $-\log|t|$ on $(-1, 1)$: $\nu_{2n} = -\int_{-1}^1 \log|t|P_{2n}(t)dt = -2\int_0^1 \log(t)P_{2n}(t)dt =$
 $\frac{2}{4n+1}\int_0^1 \frac{P_{2n+1}(t) - P_{2n-1}(t)}{t}dt = -\frac{2}{2n+1}\int_0^1 \frac{P_{2n-1}(t)}{t}dt = (-1)^n 2^n \frac{(n-1)!}{(2n+1)!}, n = 1, 2, \dots$
 $(\nu_0 = 2)$, from [29, 10.10 (52)] with $\lambda = -1$.

For $-|t|\log|t|$ on $(-1, 1)$: $\nu_{2n} = -\int_{-1}^1 |t|\log|t|P_{2n}(t)dt = -2\int_0^1 t\log(t)P_{2n}(t)dt =$
 $(-1)^{n+1} \frac{d}{d\lambda} \left[\frac{(-\lambda/2)(-\lambda/2+1)\cdots(-\lambda/2+n-1)}{(1/2+\lambda/2)(3/2+\lambda/2)\cdots(1/2+n+\lambda/2)} \right]_{\lambda=1}$
 $= (-1)^{n-1} \frac{(2n-3)!!}{2^{n+1}(n+1)!} \left(\frac{2}{3} + \cdots + \frac{2}{2n-3} + 1 + \frac{1}{2} + \cdots + \frac{1}{n+1} \right), n = 2, 3, \dots (\nu_0 = 1/2, \nu_2 =$
 $-1/16)$ from [29, 10.10 (51)].

3.3. Chebyshev examples.

We now have $R_n(x) = T_n\left(\frac{2x-a-b}{b-a}\right)$. One has $R_0 = 1, R_1(x) = (2x-a-b)/(b-a)$,

$$R_{n+1}(x) = 2(2x-a-b)R_n(x)/(b-a) - R_{n-1}(x),$$

$$R_n(a) = (-1)^n, R_n(b) = 1, \|R_n\|_R^2 = \int_a^b \frac{R_n^2(x)}{\sqrt{(x-a)(b-x)}}dx = \pi \text{ if } n = 0, \pi/2 \text{ if } n > 0.$$

Chebyshev modified moments examples are easy to capture through expansions of functions: if $F(x) = c_0/2 + \sum_1^\infty c_n T_n\left(\frac{2x-a-b}{b-a}\right)$, then $c_n = \frac{2}{\pi} \int_a^b \frac{F(t)}{\sqrt{(t-a)(b-t)}} T_n\left(\frac{2t-a-b}{b-a}\right) dt$.

Consider first the generating function of Chebyshev polynomials $\frac{1}{A-x} = \frac{4/(b-a)}{z+z^{-1}-2u} =$
 $\frac{4/(b-a)}{z-z^{-1}} [1 + 2\sum_1^\infty T_n(u)/z^n]$, where $A = (a+b)/2 + (b-a)(z+z^{-1})/4$ and $x = (a+b)/2 + (b-a)u/2 \in [a, b]$ if $-1 \leq u \leq 1$ ([77, § 5.2], [90], etc.) so,

$$\int_a^b T_n\left(\frac{2t-a-b}{b-a}\right) \frac{dt}{(A-t)\sqrt{(t-a)(b-t)}} = \frac{4\pi}{(b-a)(z-z^{-1})z^n}, \quad n = 0, 1, \dots \quad (12)$$

where $A = (a+b)/2 + (b-a)(z+z^{-1})/4$ with $|z| > 1$ if $A \notin [a, b]$, or also $z = [2A - a - b + 2\sqrt{(A-a)(A-b)}]/(b-a)$.

A logarithmic example: $\log(x-x_0)$, with $x_0 \in [a, b]$. As before, let $x = (a+b)/2 + (b-a)(z+z^{-1})/4$, $x_0 = [(a+b) + (b-a)\cos\theta_0]/2$, then, $\log(x-x_0) = \log[(a-b)\exp(-i\theta_0)(1-z\exp(i\theta_0))(1-z^{-1}\exp(i\theta_0))/4] = \log((a-b)/4) - i\theta_0 - \sum_1^\infty \frac{z^n + z^{-n}}{n} \exp(in\theta_0)$
 $= \log((a-b)/4) - i\theta_0 - 2\sum_1^\infty \frac{T_n\left(\frac{2x-a-b}{b-a}\right)}{n} \exp(in\theta_0)$ [77, 90], etc. Remark that the imaginary part is a simple discontinuity (jump, sign function) at x_0 . Other examples can be found in Milovanović & Cvetković's paper [82, Thm 2.2].

The examples of § 8 of the present study will be worked through Chebyshev modified moments expansions too, see (40) and (45).

3.4. The algorithm. Stable and efficient computation of the recurrence coefficients of (2) from the modified moments (10) has been first published by Sack and Donovan in 1969 [94, 95], immediately enthusiastically commented and expanded by W. Gautschi [37] whose exposition is summarized here (see also [39, § 2.1.7], [41, viii, pp. 28,36, 58–64, 92–100, 162]).

One does not compute the matrix of the left-hand side of (11) to get the orthogonal polynomials P_n . Instead, we use polynomials R_n satisfying themselves a known recurrence formula

$$xR_k(x) = A_k R_{k+1}(x) + B_k R_k(x) + \cdots + Z_k R_{k-s}(x), \quad (13)$$

containing the ordinary moments case when $s = 0$, some other (possibly formal) orthogonal polynomials when $s = 1$, and we shall even try an example where $s = 2$!

We make vectors

$$\mathbf{v}^{(n)} = \left[\int_a^b P_n(t)R_0(t)d\mu(t), \int_a^b P_n(t)R_1(t)d\mu(t), \dots, \int_a^b P_n(t)R_{2N-n}(t)d\mu(t) \right],$$

looking like the rows of the last factor of (11), for $n = 0, 1, \dots, N - 1$, starting of course with the modified moments at $n = 0$. By orthogonality of P_n and polynomials of degree $< n$, one must have $\mathbf{v}_0^{(n)} = \mathbf{v}_1^{(n)} = \cdots = \mathbf{v}_{n-1}^{(n)} = 0$. We also define $\mathbf{v}^{(-1)}$ to be the null vector. Then, by (2) and (13),

$$\begin{aligned} \mathbf{v}_k^{(n+1)} &= \int_a^b P_{n+1}(t)R_k(t)d\mu(t) = \int_a^b (t - b_n)P_n(t)R_k(t)d\mu(t) - a_n^2 \int_a^b P_{n-1}(t)R_k(t)d\mu(t) \\ &= \int_a^b [A_k R_{k+1}(t) + (B_k - b_n)R_k(t) + \cdots + Z_k R_{k-s}(t)]P_n(t)d\mu(t) - a_n^2 \int_a^b P_{n-1}(t)R_k(t)d\mu(t) \\ &= A_k \mathbf{v}_{k+1}^{(n)} + (B_k - b_n) \mathbf{v}_k^{(n)} + \cdots + Z_k \mathbf{v}_{k-s}^{(n)} - a_n^2 \mathbf{v}_k^{(n-1)}, \end{aligned}$$

using therefore elements of $\mathbf{v}^{(n)}$ and $\mathbf{v}^{(n-1)}$.

As one must have $\mathbf{v}_{n-1}^{(n+1)} = 0$, $a_n^2 = A_{n-1} \mathbf{v}_n^{(n)} / \mathbf{v}_{n-1}^{(n-1)}$ if $n > 0$ follows, and $\mathbf{v}_n^{(n+1)} = 0 \Rightarrow b_n = B_n + [A_n \mathbf{v}_{n+1}^{(n)} - a_n^2 \mathbf{v}_n^{(n-1)}] / \mathbf{v}_n^{(n)} = B_n + A_n \mathbf{v}_{n+1}^{(n)} / \mathbf{v}_n^{(n)} - A_{n-1} \mathbf{v}_n^{(n-1)} / \mathbf{v}_{n-1}^{(n-1)}$.

See also that $\mathbf{v}_k^{(n+1)}$ needs $\mathbf{v}_{k+1}^{(n)}$, that's why we need the $\mathbf{v}_k^{(0)}$'s up to $k = 2N$, in order to have a correct $\mathbf{v}_N^{(N)}$.

There will be much ado later on about the Chebyshev polynomials on $[a, b]$: $R_0(t) \equiv 1$, $R_1(t) = T_1((2t - a - b)/(b - a)) = (2t - a - b)/(b - a)$, $R_2(t) = T_2((2t - a - b)/(b - a)) = 2((2t - a - b)/(b - a))^2 - 1, \dots$ satisfying

$$tR_n(t) = (b - a)R_{n-1}(t)/4 + (a + b)R_n(t)/2 + (b - a)R_{n+1}(t)/4.$$

Remark. If we have a software allowing fast shift vector operations $\text{shiftright}([a_1, \dots, a_N]) = [a_2, \dots, a_N, 0]$, $\text{shiftright}([a_1, \dots, a_N]) = [0, a_1, \dots, a_{N-1}]$, then $\mathbf{v}^{(n+1)} = (b - a)[\text{shiftright}(\mathbf{v}^{(n)}) + \text{shiftright}(\mathbf{v}^{(n)})]/4 + (a + b)\mathbf{v}^{(n)}/2 - a_n^2 \mathbf{v}^{(n-1)} - b_n \mathbf{v}^{(n)}$ [40].

4. EXPANSIONS IN FUNCTIONS OF THE SECOND KIND.

We proceed with modified moments and related expansions. The weight function w is not always given in such an explicit form allowing a fast way to compute the modified moments. It is often better to use the generating function $S(x)$ of the power moments, but how is $S(x)$ an expansion involving modified moments?

From now on, we choose R_n to be an orthogonal polynomial of degree n with respect to a weight function w_R on $[a, b]$, and the searched P_n orthogonal with respect to the weight function w so that $d\mu(x) = w(x)dx$. We will often need the ratio w/w_R , a writing more realistic than the Radon-Nykodim derivative $d\mu/d\mu_R$ in most cases.

We saw that the Laurent expansion of the Stieltjes function of w with the power moments is $S(x) = \int_a^b w(t)(x-t)^{-1} dt = \sum_0^\infty \mu_k x^{-k-1}$. See here the expansion involving the modified moments $\nu_n = \int_a^b R_n(t) w(t) dt$:

4.1. Theorem. *Let R_n , $n = 0, 1, \dots$ be orthogonal polynomials related to a weight w_R on $[a, b]$, with $\|R_n\|_R^2 = \int_a^b R_n^2(t) w_R(t) dt$, and $S(x) = \int_a^b (x-t)^{-1} w(t) dt$ be the Stieltjes function of the weight function w . Then,*

$$S(x) = \sum_0^\infty \frac{\nu_n}{\|R_n\|_R^2} Q_n(x), \quad (14)$$

for $x \notin [a, b]$, where ν_n is the modified moment $\int_a^b R_n(t) w(t) dt$, and where

$Q_n(x) = \int_a^b (x-t)^{-1} R_n(t) w_R(t) dt$ is the n^{th} function of the second kind related to the weight w_R .

Indeed, as R_n is a finite linear combination of powers, which may be inverted as $t^k = \sum_{n=0}^k c_{n,k} R_n(t)$, we have $S(x) = \sum_{k=0}^\infty \int_a^b t^k w(t) dt x^{-k-1} = \sum_{k=0}^\infty \int_a^b [\sum_{n=0}^k c_{n,k} R_n(t)] w(t) dt x^{-k-1} = \sum_{n=0}^\infty \nu_n [\sum_{k=n}^\infty c_{n,k} x^{-k-1}]$. Remark now the Laurent expansion $Q_n(x) = \sum_{k=n}^\infty \int_a^b t^k R_n(t) w_R(t) dt x^{-k-1} = \sum_{k=n}^\infty c_{n,k} \|R_n\|_R^2 x^{-k-1}$. \square

There is no convergence problem, at least if a and b are finite, as the Laurent expansions converges exponentially fast when $|x| > \max(|a|, |b|)$.

My first idea was to expand the ratio w/w_R in the $\{R_n\}$ basis, by $w(t)/w_R(t) = \sum_{n=0}^\infty [\int_a^b (w(u)/w_R(u)) w_R(u) R_n(u) dt = \nu_n] R_n(t) / \|R_n\|_R^2$ for t almost everywhere in $[a, b]$, but we do not need to discuss the validity of this expansion. It seems however strange that the theorem seems to be true in some eerie situations where w and w_R have different supports. The price is that the modified moments are unusually large, which makes them completely useless. This is obvious if the support of w is bigger than the support of w_R , as the R_n s are free to become large outside the support of w_R . But things are not better if the support of w is too small! Recall that the condition number of the Gram matrix \mathbf{G}_N in (11) depends also on the smallest eigenvalue, which is the infimum on polynomials p of degree $< N$ of the Rayleigh ratio $\int_a^b p^2 w dx / \int_a^b p^2 w_R dx$, and we may choose p to be very small on the part of (a, b) which is the support of w . See also Beckermann & Bourreau [8].

Expansions with functions of the second kind share properties of Laurent expansions, such as exponential speed of convergence outside $[a, b]$, and orthogonal expansions, such as the use of recurrence relations, see Barrett [7], Gautschi [38].

For Legendre functions, the connection between Laurent expansions and expansions in functions of the second kind is given by Heine's series $(x-t)^{-1} = \sum_0^\infty (2m+1) P_m(t) Q_m(x)$, $-1 < t < 1$, $x \notin [-1, 1]$ (NIST [89, § 14.28.2], etc.), so that, gathering the t^n terms,

$\frac{1}{x^{n+1}} = \sum_0^n \frac{d^n P_m(0)/dt^n}{n!} Q_m(x)$, showing how the Q_n expansion is a rearrangement of the Laurent expansion.

As a matter of fact, Heine's series is valid for any choice of orthogonal polynomials: expand $(x-t)^{-1}$ in orthogonal expansion of the R_n s:

$$\frac{1}{x-t} = \sum_{m=0}^\infty \frac{\int_a^b \frac{R_m(u) w_R(u) du}{x-u}}{\|R_m\|_R^2} R_m(t). \text{ See Area } et \text{ al. [5, 6] for more identities.}$$

The subject matter will now be strongly simplified by turning to the Chebyshev case:

4.2. Chebyshev functions of the second kind. *The functions of second kind related to the Chebyshev polynomials*

$R_n(x) = T_n((2x - a - b)/(b - a))$ are

$$Q_n(x) = \int_a^b \frac{T_n((2t - a - b)/(b - a)) dt}{(x - t)\sqrt{(t - a)(b - t)}} = \frac{\pi}{a_\infty z^n(z - 1/z)}, \quad (15)$$

where $z = [2x - a - b + 2\sqrt{(x - a)(x - b)}]/(b - a) \sim 4x/(b - a) = x/a_\infty$ for large $|x|$.

This has been established in (12) of § 3.3, where the A of (12) is the x of (15).

This formula (15) is seen as an exercise in many textbooks, as Davis & Rabinowitz [22, § 1.13], sometimes from the finite part (Hilbert transform) of (15) when $x \in (a, b)$, known to be $-(\pi/(2a_\infty))U_{n-1}(\cos \theta)$ [1, 22.13.3], [77, eq. 9.22a], also used by Weisse & al. [108, eq. (14)]. Then, when $x = b_\infty + 2a_\infty \cos \theta \pm i\varepsilon$ is close to $[a, b]$, we add $\pm \pi i T_n(\cos \theta)/\sqrt{(x - a)(b - x)}$ (Sokhotskyi-Plemelj [53, §14.1]), and (15) is restored in a neighborhood of $[a, b]$.

It will also be recalled in § 5.3 that the asymptotic formula (21b) is exact in the Bernstein-Szegő case (when $\sqrt{(t - a)(b - t)}/w(t)$ is a polynomial, and when $n >$ half the degree of this polynomial). Henrici gives (15) in [53, § 14.6, Problem 2] with the symbol " U_n " for our Q_n .

4.3. Corollary. *Chebyshev modified moments are the coefficients of the expansion of the Stieltjes function in negative powers of z*

$$\frac{(b - a)(z - z^{-1})}{2} S \left(x = \frac{a + b}{2} + \frac{(b - a)(z + z^{-1})}{4} \right) = 2\nu_0 + \sum_1^\infty \frac{4\nu_n}{z^n}. \quad (16)$$

Indeed, put (15) in (14)

$$S(x) = \sum_0^\infty \frac{\nu_n}{\|R_n\|_R^2} Q_n(x) = \frac{\nu_0}{\|R_0\|_R^2} \frac{\pi/a_\infty}{z - z^{-1}} + \sum_1^\infty \frac{\nu_n}{\|R_n\|_R^2} \frac{\pi/a_\infty}{z^n(z - z^{-1})}. \quad \square$$

Consider also (see § 3.3) the generating function of Chebyshev polynomials.

5. WEIGHTS WITH LOGARITHMIC SINGULARITIES. INTERIOR SINGULARITY.

5.1. Known results. The influence of an algebraic singularity at $c \in (a, b)$ on the recurrence coefficients has been discussed in [33, 71, 73], it has been observed, and sometimes proved, that

$$\begin{aligned} a_n - a_\infty &= f_n \cos(2n\theta_c + \varphi_c) + o(f_n), \\ b_n - b_\infty &= 2f_n \cos((2n + 1)\theta_c + \varphi_c) + o(f_n), \end{aligned} \quad (17)$$

where $c = \frac{a + b}{2} + \frac{b - a}{2} \cos \theta_c$, with $0 < \theta_c < \pi$, $a_\infty = \frac{b - a}{4}$, $b_\infty = \frac{a + b}{2}$ from (8), and where f_n and φ_c depend on the kind of singularity.

We first consider here **weak singularities**, i.e. when the weight function remains continuous, bounded and bounded from below by a positive number in a neighborhood of the singular point. For an algebraic weak singularity,

$$\begin{aligned} w(x) &\approx w(c) + \text{constant } |x - c|^\alpha, \alpha > 0, 0 < w(c) < \infty \\ &\Rightarrow f_n = \text{constant } n^{-\alpha-1}, \end{aligned} \quad (18)$$

has been shown in [71, p.156], see also the first part of [73].

The proof in [71] makes heavy use of Toeplitz determinants based on Szegő's theory. The subsections 5.3-5.3.5 that will follow here try to suggest simpler approaches. No new proof will be presented, but the puzzling fact is that the form (17) holds for all kinds of interior singularities encountered up to now, whether as proved results, or as conjectures.

In the much more difficult situation of a *strong singularity* where $w(c) = 0$ or ∞ , the power of n in f_n does not depend on α , it happens that

$$w(x) \sim \text{constant } |x - c|^\alpha, \alpha > -1 \Rightarrow f_n = -(b - a)|\alpha|/(8n), \quad (19)$$

as conjectured in [73], and brilliantly proved in [33], through new Riemann-Hilbert techniques. Initially a way to reconstruct systems of differential equations from the properties of the solutions about their branchpoints (monodromy matrix), "the Riemann-Hilbert method reduces a particular problem to the reconstruction of an analytic function from jump conditions", in the words of A. Its [56, p. 1389]. The method has been applied with striking results to orthogonal polynomials asymptotics [56, p. 1391] [4, 23, 31, 61, 62], up to, as already noted, the proof of conjectures on interior singularities [33, 34]. For our orthogonal polynomials problems, we must limit ourselves to weight functions that can be analytically continued in a domain containing the support, up to a finite number of singular points, and cuts that join them. When such a singular point is a *branchpoint*, the continuation on a surrounding contour (from "before" to "after", when the contour's turn is achieved) leads to new values, best described by

$$\begin{bmatrix} P_n(x) & Q_n(x)/w(x) \\ P_{n-1}(x) & Q_{n-1}(x)/w(x) \end{bmatrix}_{\text{after}} = \begin{bmatrix} P_n(x) & Q_n(x)/w(x) \\ P_{n-1}(x) & Q_{n-1}(x)/w(x) \end{bmatrix}_{\text{before}} \begin{bmatrix} 1 & C_1 \\ 0 & C_2 \end{bmatrix}$$

with suitable C_1, C_2 making the relevant *monodromy matrix* around the singular point [74, §3.4], with a very subtle scale dilation in some regions, this is the new feature found in present day developments of the Riemann-Hilbert method [4, 23, 33, 34, 31, 61, 62], [55, chap. 22].

In the year 2000, Lubinsky wrote illuminating remarks on the virtues of various techniques, see here a (slightly shortened) part of the conclusion of his survey: " At present it seems that the Deift-Fokas-Its-Kitaev-Kriecherbauer- McLaughlin & al. method [= the new RH] will lead to very precise asymptotics for restricted classes of weights, while I believe that Bernstein-Szegő-Rakhmanov's one will lead to asymptotics for more general weights, but with weaker error estimates. " [70, p. 251].

$w(t)$	1	2	3	4	5	6	7
$-\log t $	$\frac{1}{9}$	$\frac{56}{225}$	$\frac{1863}{8575}$	$\frac{162776}{639009}$	$\frac{1070243125}{4586681781}$	$\frac{25913014364304}{101783375460481}$	$\frac{1934965659283089461}{8086464983903415525}$
$- t \log t $	$\frac{1}{4}$	$\frac{7}{36}$	$\frac{17}{63}$	$\frac{647}{2975}$	$\frac{294777}{1099900}$	$\frac{134451657}{593353348}$	$\frac{5943701732}{22386249711}$

TABLE 3. First recurrence coefficients a_n^2 for the weights $-\log |t|$ and $-|t| \log |t|$ on $(-1, 1)$.

5.2. Conjecture. *If the weight function has one or several logarithmic singularities of the form $w(x) \sim \text{constant} \times |x - c|^\alpha \log |x - c|$ near one or several values of $c \in (a, b)$, the first terms of the asymptotic behavior of the related amplitude in (17) are*

$$f_n = \frac{A}{n} + \frac{B}{n \log n} + \frac{C}{n(\log n)^2} + o(1/(n(\log n)^2)), \quad (20)$$

with $A = -(b - a)|\alpha|/8$. When $\alpha = 0$, $B = (b - a) \sin(\theta_c)/8$ is conjectured, where $c = (a + b + (b - a) \cos \theta_c)/2$.

One has also $(b - a) \sin \theta_c = 2\sqrt{(c - a)(b - c)}$. The formula for A in the conjecture is the same as in (19). As a matter of fact, the validity of the formulas for A and especially B in f_n is the weakest part of the conjecture, and may be wrong.

The examples of tables 3-5 have been computed from the Legendre modified moments of § 3.2.

n	a_n	$n(1/2 - a_n)$	B	A	C
1	0.5000000000	0			
2	0.4409585518	0.1180828963			
3	0.5194628810				
4	0.4663464287	0.1346142850	0.02292049	-0.15114792	-0.12482
5	0.5176930736				
8	0.4812117607	0.1503059146	0.06526631	-0.18169237	-0.088775
16	0.4896880521	0.1649911660	0.12215957	-0.20905091	-0.039663
32	0.4944429849	0.1778244819	0.17792267	-0.22916211	0.000490
64	0.4970563023	0.1883966532	0.21986008	-0.24126182	0.025789
128	0.4984622524	0.1968316976	0.24558204	-0.24744598	0.038691
256	0.4992050878	0.2034975345	0.25876286	-0.25016202	0.043609
512	0.4995922083	0.2087893758	0.26411801	-0.25112734	0.044200
1024	0.4997919518	0.2130413275	0.26527060	-0.25131178	0.042822
2048	0.4998942823	0.2165098115	0.26447849	-0.25119724	0.040791
4096	0.4999464395	0.2193838028	0.26297659	-0.25100005	0.038747
8192	0.4999729247	0.2218007622	0.26136741	-0.25080641	0.036952
16384	0.4999863366	0.2238608480	0.25990460	-0.25064392	0.035477
32768	0.4999931141	0.2256377520	0.25864776	-0.25051440	0.034304

TABLE 4. Analysis of the recurrence coefficients for the weight $-|t| \log |t|$ on $(-1, 1)$.

Table 4 shows a check of conjecture 5.2 for the even weight function $-|t| \log |t|$ on $(-1, 1)$. So, $\theta_c = \pi/2$ and, as $b_n \equiv 0$, $\varphi_c = 0$ or π . Having chosen $A < 0$, as in (19), $\varphi = 0$. Then, A and B are estimated from linear interpolation of $n(1/2 - a_n)$ in the variable $1/\log n$. The limit value $-1/4$ for A is quite credible, that $B = 1/4$ is suggested, and C is estimated from $(\log n)^2[n(a_n - 1/2) + 1/4 - 1/(4 \log n)]$, thus assuming $A = -1/4$ and $B = 1/4$.

The same experiment with $-\log |t|$, so $\alpha = 0$, yields indeed $A = 0$ and, with $\varphi = 0$, very likely, $B = 1/4$ in table 5, using an estimate of C as the slope of $n(\log n)(a_n - 1/2)$ as a function of $1/\log n$. Assuming $B = 1/4$, a new estimate of C is found from $[n(\log n)(a_n - 1/2) - 1/4] \log n$.

No tentative formula for C can be launched from the small number of available cases explored here.

The conjecture 5.2 is motivated by interesting solid-state problems in §8, where weight functions with logarithmic singularities are encountered, and numerical results are seen in Tables 8 and 9.

5.3. About the Szegő asymptotic formula.

Warning: up to the end of the present section 5, I develop some ideas on the origin of the formulas (17), trying to show in several ways how they are related to the Szegő's theory.

No new proof follows, alas, but perhaps valuable work will be inspired by what follows.

The Szegő's theory is centered on the asymptotic behavior of the orthogonal polynomials of large degree. This description has been achieved by G. Szegő long ago, and is available of course in his book [101, chap. 12], also in the surveys by Lubinsky [69, 70] and Nevai [84], and in Van Assche's book [103, §1.3.1] the formula for the orthonormal polynomial is

$$p_n(x) \approx p_n^{(0)}(x) = (2\pi)^{-1/2} [z^n \exp(\lambda(z^{-1})) + z^{-n} \exp(\lambda(z))], \quad (21a)$$

where $x = b_\infty + 2a_\infty \cos \theta$, $z = e^{i\theta}$, and

$$\lambda(z) = \lambda_0/2 + \lambda_1 z + \lambda_2 z^2 + \dots$$

n	a_n	$n(\log n)(a_n - 1/2)$	C	B	C
1	0.3333333333				
2	0.4988876516	-0.0015420423			
3	0.4661110117				
4	0.5047097547	0.0261164255	-0.038348	0.053778	-0.310368
5	0.4830498129				
8	0.5039795051	0.0662011853	-0.166724	0.146378	-0.382198
16	0.5023625151	0.1048045226	-0.321122	0.220625	-0.402567
32	0.5012250826	0.1358660071	-0.430640	0.260122	-0.395558
64	0.5005968024	0.1588500221	-0.477978	0.273779	-0.379082
128	0.5002823026	0.1753268095	-0.479713	0.274195	-0.362316
256	0.5001318186	0.1871251805	-0.458004	0.269720	-0.348652
512	0.5000612761	0.1957173003	-0.428836	0.264459	-0.338633
1024	0.5000284796	0.2021431261	-0.400894	0.259979	-0.331718
2048	0.5000132625	0.2070968803	-0.377733	0.256638	-0.327119
4096	0.5000061941	0.2110317678	-0.360050	0.254318	-0.324128
8192	0.5000029023	0.2142428737	-0.347245	0.252778	-0.322204
16384	0.5000013644	0.2169243207	-0.338296	0.251785	-0.320968
32768	0.5000006434	0.2192063024	-0.332167	0.251154	-0.320168

TABLE 5. Some recurrence coefficients of large index for the weight $-\log|t|$ on $(-1, 1)$.

is the half constant term and the part with positive powers of the Laurent-Fourier expansion

$$-\log[w(x)\sqrt{(x-a)(b-x)}] = -\log[2a_\infty w(b_\infty + 2a_\infty \cos \theta) |\sin \theta|] = \sum_{-\infty}^{\infty} \lambda_k z^k = \lambda(z) + \lambda(z^{-1})$$

on $|z = e^{i\theta}| = 1$. The condition of asymptotic validity is the minimal condition $\log[w(x)\sqrt{(x-a)(b-x)}] \in L_1$ (Szegő class). The function $D(z) = \exp(-\lambda(z))$ is the *Szegő function* associated to the weight w , it is analytic without zero in the unit disk, and satisfies $|D(z)|^2 \rightarrow w(x)\sqrt{(x-a)(b-x)}$ when $|z| \rightarrow 1$. Remark that the $\lambda_{-n} = \lambda_n$ s are real. When $\exp(\lambda(z))$ is a polynomial of degree, say d , the formula (21a) is exact for $n > d/2$ (Bernstein-Szegő class [70, § 2.1]). For a general weight in the Szegő class, $p_n^{(0)}(x)$ in (21a) is not a polynomial but has an expansion $\kappa_n^{(0)} x^n + \kappa_n^{\prime(0)} x^{n-1} + \dots$ for large x . Also, the two sides of (21a) tend to be equal in some L_2 norm [70, eq. (12)] etc.

In the simplest case $w(x) = 1/\sqrt{(x-a)(b-x)}$, $p_n(x) = \sqrt{2/\pi} \cos n\theta = \sqrt{2/\pi} T_n((x-b_\infty)/(2a_\infty))$, $\lambda(z) \equiv 0$. For Chebyshev polynomials of the second kind, $w(x) = \sqrt{(x-a)(b-x)}$, $p_n(x) = \frac{\sin(n+1)\theta}{a_\infty \sqrt{2\pi} \sin \theta} = (1/\sqrt{2\pi}) U_n((x-b_\infty)/(2a_\infty))$, $w(x)\sqrt{(x-a)(b-x)} = (x-a)(b-x) = 4a_\infty^2 \sin^2 \theta = -a_\infty^2 (z - z^{-1})^2 = a_\infty^2 (1 - z^2)(1 - z^{-2})$, $e^{\lambda(z)} = 1/[a_\infty(1 - z^2)]$.

For the function of the second kind $q_n(x) = \int_a^b \frac{p_n(t)w(t) dt}{x-t}$,

$$q_n(x) \approx q_n^{(0)}(x) = (2\pi)^{1/2} \frac{4}{b-a} \frac{\exp(-\lambda(z^{-1}))}{z^n(z - z^{-1})}, \quad (21b)$$

see Barrett[7], also Van Assche[103, §5.4].

We also have $z = \cos \theta + i \sin \theta = (b-a)^{-1} [2x - a - b + 2\sqrt{(x-a)(x-b)}]$, with the square root such that $|z| > 1$ if $x \notin [a, b]$, in which case only the term containing z^n has to be considered in (21a). Remark that $x = b_\infty + a_\infty(z + 1/z) \Rightarrow z = (x - b_\infty)/a_\infty + O(1/x)$ when x is large, allowing to estimate the coefficients of x^n and x^{n-1} : with $p_n^{(0)}(x)$ and $q_n^{(0)}(x)$ of (21a)-(21b), when

x is large, $p_n(x) \approx p_n^{(0)}(x) = \kappa_n^{(0)} x^n + \kappa_n^{\prime(0)} x^{n-1} + \dots$, and

$$\kappa_n^{(0)} = \frac{\exp(\lambda_0/2)}{\sqrt{2\pi} (a_\infty)^n}, \quad \frac{\kappa_n^{\prime(0)}}{\kappa_n^{(0)}} = -nb_\infty + a_\infty \lambda_1. \quad (22)$$

Indeed, $p_n^{(0)}(x) = (2\pi)^{-1/2} z^n \exp(\lambda_0/2 + \lambda_1 z^{-1} + \lambda_2 z^{-2} + \dots)$ and $z = x/a_\infty - b_\infty/a_\infty - a_\infty/x + O(x^{-2})$, so, the coefficient of x^n , $\kappa_n^{(0)} = (2\pi)^{-1/2} \exp(\lambda_0/2)/a_\infty^n$, and $z = x/a_\infty - b_\infty/a_\infty - a_\infty/x + O(x^{-2}) \Rightarrow p_n^{(0)}(x)/\kappa_n^{(0)} = (a_\infty z)^n \exp(\lambda_1 z^{-1} + \lambda_2 z^{-2} + \dots) = (a_\infty z)^n + a_\infty \lambda_1 (a_\infty z)^{n-1} + a_\infty^2 (\lambda_1^2/2 + \lambda_2) (a_\infty z)^{n-2} + \dots = x^n - (nb_\infty - a_\infty \lambda_1) x^{n-1} + \dots$.

For $p_n(x) = \kappa_n x^n + \kappa_n' x^{n-1} + \dots$ itself, from the recurrence relation (2) $P_n(x) = p_n(x)/\kappa_n = (x - b_{n-1})P_{n-1}(x) - a_{n-1}^2 P_{n-2}(x)$, and $\|P_n\|^2 = \mu_0 a_1^2 \dots a_n^2$:

$$\kappa_n = \frac{1}{\sqrt{\mu_0} a_1 \dots a_n}, \quad \frac{\kappa_n'}{\kappa_n} = -b_0 - \dots - b_{n-1}. \quad (23)$$

Each term of (23) behaves like the corresponding term of (22) when $n \rightarrow \infty$.

In terms of z such that $x = a_\infty z + b_\infty + a_\infty/z$:

$$p_n(x)/\kappa_n = a_\infty^n z^n + a_\infty^{n-1} (nb_\infty - b_0 - \dots - b_{n-1}) z^{n-1} + \dots \quad (24)$$

We have gathered useful asymptotic material, and try to see how it allows to explain the formulas (17).

First, a quick check of (21a): $\int_a^b p_n(x) p_m(x) w(x) dx \approx \int_a^b p_n^{(0)}(x) p_m^{(0)}(x) w(x) dx$

$$= \frac{1}{2\pi} \int_a^b [z^n \exp(\lambda(z^{-1})) + z^{-n} \exp(\lambda(z))] [z^m \exp(\lambda(z^{-1})) + z^{-m} \exp(\lambda(z))] \frac{\exp(-\lambda(z) - \lambda(z^{-1})) dx}{a_\infty |z - z^{-1}|}.$$

With $x = b_\infty + a_\infty(z + z^{-1})$, $dx = a_\infty(z - z^{-1}) \frac{dz}{z}$, we have the integral on the unit circle

$\frac{1}{4\pi i} \oint [z^{n+m} \exp(\lambda(z^{-1}) - \lambda(z)) + z^{n-m} + z^{m-n} + z^{-n-m} \exp(\lambda(z) - \lambda(z^{-1}))] \frac{dz}{z}$. The central terms leave no residue if $m \neq n$. When $m = n$, the result is unity, together with perturbations involving high index Fourier coefficients of $\exp(\lambda(z) - \lambda(z^{-1}))$.

The same technique is now used in order to get these high index coefficients, and how they enter the following estimate of recurrence coefficients finer asymptotics:

$$a_n - a_\infty \approx \frac{a_\infty}{2} (\psi_{-2n+2} - \psi_{-2n}), \quad b_n - b_\infty \approx a_\infty (\psi_{-2n+1} - \psi_{-2n-1}), \quad (25)$$

where the ψ s are the Fourier coefficients of

$$\psi(e^{i\theta}) = \exp(\lambda(e^{-i\theta}) - \lambda(e^{i\theta})) = \sum_{-\infty}^{\infty} \psi_k e^{ik\theta}. \quad (26)$$

The formula (25) has been established by a long and painful proof through Toeplitz determinants in [71, p. 153, 158, 167] for weak singularities (the weight function w being continuous and bounded from below by a positive number at the singular point). It is wondered here if we can reach (25) more easily, and if these formulas still hold for stronger singularities, but to which strength?

Remark that $\sum 2(a_n - a_\infty)z^{-2n} + (b_n - b_\infty)z^{-2n-1} \sim a_\infty \sum [\psi_{-k+2} z^{-k} - \psi_{-k} z^{-k}] =$ sum of negative exponents of z in $(z^{-2} - 1)\psi(z)$.

Remark also that $\exp(\lambda(z^{-1}) - \lambda(z)) = D(z)/D(z^{-1})$ in Szegő's notation, is an *inner function*³, i.e., of modulus unity when $|z| = 1$, so $\sum \psi_n \psi_{n+k} = \delta_{k,0}$.

³ D itself is an *outer function* [44, § 8], [70, p.211].

An even stronger estimate follows from a refinement of the asymptotic matching of (22) and (23):

$$\sqrt{\frac{\mu_0 e^{\lambda_0}}{2\pi} \frac{a_1 \cdots a_n}{a_\infty^n}} - 1 \sim -\frac{\psi_{-2n}}{2}, \quad b_0 + \cdots + b_{n-1} - nb_\infty + a_\infty \lambda_1 \sim -a_\infty \psi_{-2n+1}. \quad (27)$$

Can we find a quick and dirty argument for (25) and (27)?

5.3.1. *From orthogonal polynomials coefficients.* Consider the square of the norm of the monic orthogonal polynomial $\mu_0 a_1^2 \cdots a_n^2 = \|P_n\|^2 \approx \|P_n^{(0)}\|^2 = \int_a^b (p_n^{(0)}(t))^2 w(t) dt / (\kappa_n^{(0)})^2$ with $p_n^{(0)}$ being the right-hand side of (21a). We take a better look at the integral $\int_a^b (p_n^{(0)}(t))^2 w(t) dt = \frac{1}{4\pi i} \oint [z^{2n} \exp(\lambda(z^{-1}) - \lambda(z)) + 2 + z^{-2n} \exp(\lambda(z) - \lambda(z^{-1}))] \frac{dz}{z} = 1 + \psi_{-2n}/2 +$ the half of the coefficient of $\exp 2ni\theta$ in the complex conjugate $\overline{\psi(e^{i\theta})}$ which is $\psi_{-2n}/2$ again. We now need $\kappa_n^{(0)}$, already estimated in (22), but we need again a refined estimation. The coefficient of x^n in $p_n^{(0)}$ is estimated through the projection on the n^{th} degree element of an orthonormal basis of polynomials, so, by $p_n(x)$ times the scalar product of $p_n^{(0)}$ and the unknown p_n , which we replace by $\dots p_n^{(0)}$ (this part of the argument is very weak), and we get refined $\kappa_n^{(0)}$ = the $\kappa_n^{(0)}$ of (22) times the square of the norm of $p_n^{(0)}$, which is $1 + \psi_{-2n}$ as seen above, and $\mu_0 a_1^2 \cdots a_n^2 \approx \frac{2\pi \exp(-\lambda_0) (a_\infty)^{2n}}{1 + \psi_{-2n}}$ follows, leading to the first part of (27). For the second part, see that $-b_0 \cdots -b_{n-1}$ is the coefficient of x^{n-1} of $p_n(x)/\kappa_n \approx p_n^{(0)}(x)/\kappa_n^{(0)}$ estimated by its projections on p_n and p_{n-1} again replaced (same caution) by $p_n^{(0)}$ and $p_{n-1}^{(0)}$. Result is $-b_0 \cdots -b_{n-1} \approx \kappa_n^{(0)'} / \kappa_n^{(0)} + (\kappa_{n-1}^{(0)} / \kappa_n^{(0)})$ times the scalar product of $p_n^{(0)}$ and $p_{n-1}^{(0)} = \frac{1}{4\pi i} \oint [z^{2n-1} \exp(\lambda(z^{-1}) - \lambda(z)) + z + 1/z + z^{-2n+1} \exp(\lambda(z) - \lambda(z^{-1}))] \frac{dz}{z} = \psi_{-2n+1}$ as seen before in similar situations, and the second part of (27) follows.

This way to get (27) is far from being a valuable proof! The argument repeatedly confuses p_n and $p_n^{(0)}$, ignoring that $p_n^{(0)}$ is normally NOT a polynomial, so that various ways of estimating coefficients yield various results, of which the most convenient ones are kept. I even turned to some numerical tests to be sure of the numerical credibility of (25), actually of the first part of (27): with $w(x) = (1 - x^2)^{-1/2} \exp(|x|)$ on $(-1, 1)$, $\lambda(e^{i\theta}) + \lambda(e^{-i\theta}) = -|\cos \theta|$, $\lambda_{2n} = 2(-1)^n / ((4n^2 - 1)\pi)$, $\lambda(z) = (2/\pi)(-1/2 - z^2/3 + z^4/15 - \dots) = -(1 + z^2) \frac{1}{2\pi i z} \log \frac{1 + iz}{1 - iz} = i\pi^{-1} \cos \theta \log[i \cot(\pi/4 + \theta/2)]$ on $|z = e^{i\theta}| = 1$, actually $\cos \theta [-1/2 + i\pi^{-1} \log \cot(\pi/4 + \theta/2)]$ when $-\pi/2 \leq \theta \leq \pi/2$, $\cos \theta [1/2 + i\pi^{-1} \log \cot(-\pi/4 + \theta/2)]$ when $\pi/2 \leq \theta \leq 3\pi/2$: $\lambda(e^{-i\theta}) - \lambda(e^{i\theta}) = 2i\pi^{-1} \cos \theta \log \cot(\pm\pi/4 + \theta/2)$. We check that $\lambda(e^{-i\theta}) + \lambda(e^{i\theta}) = -|\cos \theta|$ indeed.

From (27), the product $M_n = M_0 a_1 \cdots a_n / a_\infty^n \rightarrow 1$ when $M_0 = \sqrt{\mu_0 \exp(\lambda_0) / (2\pi)}$. Here, $\mu_0 = \int_{-1}^1 w(t) dt = 6.2088$, $\lambda_0 = -0.63662$, $M_0 = 0.72306$, some a_n, λ_n, ψ_n are shown, and $2M_n - 2$ shows how M_n is close to $1 - \psi_{-2n}/2$ according to (27).

n	0	1	2	3	4	5	10
a_n		0.77414	0.43434	0.52081	0.49034	0.50548	0.49893
λ_{2n}	-0.63662	-0.21221	0.042441	-0.01819	0.01011	-0.00643	0.00160
ψ_{2n}	0.95317	0.21745	-0.022831	0.01252	-0.00738	0.00489	-0.00134
ψ_{-2n}	0.95317	-0.19755	0.058329	-0.02507	0.01353	-0.00838	0.00191
$2M_n - 2$		0.23899	-0.055036	0.02590	-0.01325	0.00851	-0.00189

TABLE 6. Results for $e^{|t|} / \sqrt{1 - t^2}$.

We see that the relevant information is indeed contained in the ψ_{-2n} coefficients of (26).

5.3.2. *From scattering theory.* An alternate source of knowledge is therefore most welcome: Van Assche gave in [104] a survey on how Case, Geronimo, and Nevai (and himself too, see [86]) investigated the relation between recurrence coefficients and weight function modification, by introducing a function $\phi(x) = \lim_{n \rightarrow \infty} \frac{(z - z^{-1})P_n(x)}{a_\infty^n z^{n+1}}$ outside $[a, b]$ for x , i.e., when $|z| > 1$, and where $P_n(x)$ is the monic polynomial $p_n(x)/\kappa_n \sim \sqrt{2\pi} a_\infty^n e^{-\lambda_0/2} p_n(x)$, so that $\phi(x) = (1 - z^{-2}) \exp(\lambda(z^{-1}) - \lambda_0/2) = 1 + \lambda_1/z + (\lambda_1^2/2 + \lambda_2 - 1)/z^2 + \dots$, and it is shown in [104] that

$$\phi(x) = 1 - \sum_0^\infty \left[\frac{b_n - b_\infty}{a_\infty z^{n+1}} + \frac{a_{n+1}^2 - a_\infty^2}{a_\infty^2 z^{n+2}} \right] \frac{P_n(x)}{a_\infty^n} \quad (28)$$

valid for x up to the sides of the cut $[a, b]$ in the trace-class case ($\sum_1^\infty |a_n - a_\infty| + |b_n - b_\infty| < \infty$). Check Chebyshev polynomials of the first kind: $\lambda(z^{-1}) \equiv 0$, $\phi(x) = \lim(z - z^{-1})[2T_n(x) = z^n + z^{-n}]/z^{n+1} = 1 - z^{-2}$, OK, as only $a_1^2 = 2a_\infty^2$ is different from a_∞^2 ; Chebyshev polynomials of second kind: $\lambda(z) + \lambda(z^{-1}) = -\log(1 - (z + z^{-1})^2/4)$, $\lambda(z) = \log 2 - \log(1 - z^2)$, $\phi(x) = \lim(z - z^{-1})[U_n(x) = (z^{n+1} - z^{-n-1})/(z - z^{-1})]/z^{n+1} = 1$.

Can we extract from (28) information on $F(z) = \sum_0^\infty \left[\frac{b_n - b_\infty}{a_\infty z^{2n+1}} + \frac{a_{n+1}^2 - a_\infty^2}{a_\infty^2 z^{2n+2}} \right]$?

From (21a) and (22), $P_n(x)/(a_\infty z)^n = p_n(x)/(\kappa_n a_\infty^n z^n)$ contains a part with strongly negative powers of z which tend to be close to the corresponding part of $\exp(\lambda(z) - \lambda_0/2)z^{-2n}$, and the corresponding part of $\sum_0^\infty \left[\frac{b_n - b_\infty}{a_\infty z} + \frac{a_{n+1}^2 - a_\infty^2}{a_\infty^2 z^2} \right] \frac{P_n(x)}{a_\infty^n z^n}$, so, $(1 - z^{-2}) \exp(\lambda(z^{-1}) - \lambda_0/2) \approx 1 - e^{-\lambda_0/2} \sum_0^\infty \left[\frac{b_n - b_\infty}{a_\infty} z^{-n-1} + \frac{a_{n+1}^2 - a_\infty^2}{a_\infty^2} z^{-n-2} \right] [z^n \exp(\lambda(z^{-1})) + z^{-n} \exp(\lambda(z))]$, or, after division by $e^{\lambda(z) - \lambda_0/2}$, $(1 - z^{-2})\psi(z) \approx e^{\lambda_0/2 - \lambda(z)} - \sum_0^\infty \left[\frac{b_n - b_\infty}{a_\infty} z^{-1} + \frac{a_{n+1}^2 - a_\infty^2}{a_\infty^2} z^{-2} \right] \psi(z) - \sum_0^\infty \left[\frac{b_n - b_\infty}{a_\infty} z^{-2n-1} + \frac{a_{n+1}^2 - a_\infty^2}{a_\infty^2} z^{-2n-2} \right]$. This confirms that the latter series is related to the negative powers part of $\psi(z)$ precisely as stated in (25).

5.3.3. *From recurrence relations as matrix products.* An interesting exercise is also to recover (21a-21b) from (25) by working a linearization of a product of 2×2 matrices containing the recurrence coefficients:

$$\begin{bmatrix} p_{N-1}(x) & q_{N-1}(x) \\ p_N(x) & q_N(x) \end{bmatrix} = \prod_{n=0}^{N-1} \begin{bmatrix} 0 & 1 \\ -a_{N-1-n}/a_{N-n} & (x - b_{N-1-n})/a_{N-n} \end{bmatrix} \times \begin{bmatrix} 0 & \sqrt{\mu_0}/a_0 \\ 1/\sqrt{\mu_0} & S(x)/\sqrt{\mu_0} \end{bmatrix},$$

see ([79]), and we write each matrix factor above as $A + E_{N-n}$, with $A = \begin{bmatrix} 0 & 1 \\ -1 & (x - b_\infty)/a_\infty \end{bmatrix}$.

Then, $(A + E_{N-1}) \cdots (A + E_0)$

$\approx A^N + \sum_{n=0}^{N-1} A^{N-1-n} E_n A^n$, seeing that $A = \begin{bmatrix} 1 & 1 \\ z & z^{-1} \end{bmatrix} \begin{bmatrix} z & 0 \\ 0 & z^{-1} \end{bmatrix} \begin{bmatrix} 1 & 1 \\ z & z^{-1} \end{bmatrix}^{-1} = U \text{diag}(z, z^{-1}) U^{-1}$,

where $z + z^{-1} = (x - b_\infty)/a_\infty$, so $A^{N-1-n} E_n A^n = U \text{diag}(z^{N-1-n}, z^{-N+1+n}) U^{-1} E_n U \text{diag}(z^n, z^{-n}) U^{-1}$

and we find $U^{-1} E_n U \approx (z^{-1} - z)^{-1} \begin{bmatrix} e_n(z) & e_n(z^{-1}) \\ -e_n(z) & -e_n(z^{-1}) \end{bmatrix}$, where

$e_n(z) = [(a_{n+1} - a_\infty)z^2 + (b_n - b_\infty)z + a_n - a_\infty]/a_\infty$, so that off-diagonal elements of the sum are $z^{\pm(N-1)} \sum z^{\mp 2n} e_n(z^{\mp 1})$ containing again the sum $\sum 2(a_n - a_\infty)z^{\mp 2n} + (b_n - b_\infty)z^{\mp(2n+1)}$.

It seems here that much energy has been spent on incomplete proofs, and that somebody should achieve a decent one!

5.3.4. *Relation with Fourier coefficients asymptotics.* So, (25)-(27) relates recurrence coefficients asymptotics to Fourier coefficients of large index, a well worked subject.

The main influence of a singularity at $\theta = \theta_c$ on the Fourier coefficient $\int_{-\pi}^{\pi} f(\theta) \exp(in\theta) d\theta$ of a function f is $\exp(in\theta_c) \hat{f}(n/(2\pi))$, see Lighthill [67, p. 8, p.43, p.72], where \hat{f} is the Fourier transform⁴ of f . An algebraic singularity of the form $|\theta - \theta_c|^\alpha$ is shown to correspond to an $n^{-\alpha-1}$ behavior. This case is also given with much detail by A. Erdélyi [30, §2.8], and Zygmund [114, chap. 5, §2.24]. The nature of a weak singularity $w(c) + \text{const. } |x - c|^\alpha$ with $0 < w(c) < \infty$, is left unchanged by taking logarithms or exponentials, also in conjugate functions [114, chap.5, §2.6 and 2.24], so, the $1/n^{\alpha+1}$ is kept unchanged up to the ψ_n s and (18) is confirmed.

Stretching the argument for weak singularity to a strong singularity such as $w(t) \sim \text{constant } |t - c|^\alpha$ near c , the logarithm of w behaves like $\alpha \log |\cos \theta - \cos \theta_c|$
 $= \text{constant} + \alpha \text{Re} \log(1 - e^{i\theta}/z_c)(1 - e^{-i\theta}/z_c)$ whence $\lambda_n \sim -\alpha \text{Re } z_c^{-n}/n = -\alpha \cos(n\theta_c)/n$, $\lambda(z) \sim (\alpha/2) \log((1 - ze^{i\theta_c})(1 - ze^{-i\theta_c})) = (\alpha/2) \log(2e^{i\theta}(\cos \theta - \cos \theta_c))$ on the circle. Keeping logarithms of positive numbers to be real, $\lambda(e^{i\theta}) \sim (\alpha/2)[\log 2 + i\theta + \log(\cos \theta - \cos \theta_c)]$ when $-\theta_c < \theta < \theta_c$, $(\alpha/2)[\log 2 + i\theta - i\pi + \log(\cos \theta_c - \cos \theta)]$ otherwise. Then, $\lambda(e^{-i\theta}) - \lambda(e^{i\theta}) \sim -i\alpha\theta$ on the first arc, $i\alpha(\pi - \theta)$ on the second arc, and its exponential has $\psi_n = (2\pi)^{-1} [\int_{-\theta_c}^{\theta_c} \exp(-i(n + \alpha)\theta) d\theta + \exp(i\alpha\pi) \int_{\theta_c}^{2\pi - \theta_c} \exp(-i(n + \alpha)\theta) d\theta] = \frac{2 \sin(\alpha\pi/2) \cos(n\theta_c + \alpha(\theta_c - \pi/2))}{\pi(\alpha + n)}$ showing an $1/n$ asymptotic behavior, but the amplitude is not right, it should have been $-a_\infty|\alpha|/2$ from (19).

So, (25)-(27) hold for weak singularities $0 < w(c) < \infty$ as they should, but fail for at least one important class of strong singularities $w(c) = 0$ or ∞ .

Remind that the correct result has been established through the new Riemann-Hilbert method [33, 34].

And what about a logarithmic singularity of the conjectures of § 5.2-6.2, as encountered with 2-dimensional crystals?

Let $w(x) - A \log|x - c|$ be continuous in a neighborhood of $c \in (a, b)$.

Now, in the case of conjecture 5.2, $\log w(t)$ has a $\log \log$ singularity! There is probably not much literature on Fourier coefficients of a $\log(\log|t - c|)$ singularity, but Zygmund [114, chap. 5, §2.31] and Wong & Lin [111] show how to arrive at a $n^{-m-1}(\log n)^{\beta-1}$ from a $|t - c|^m(\log|t - c|)^\beta$ singularity, when m is an integer. Take $m = 0$ and $\beta \rightarrow 0$, as $\log(\log|t - c|)$ is the limit when $\beta \rightarrow 0$ of $\beta^{-1}[(\log|t - c|)^\beta - 1]$, we may expect the $1/(n \log n)$ of the conjecture. Two meaningful examples will be considered in § 8.

5.3.5. *Relation between jumps and logarithmic singularities.* The Fourier series conjugate to the real part of $\sum c_k e^{ik\theta}$ is the imaginary part of the same expansion [114, § 1]. Jumps and logarithmic singularities are conjugate phenomena. A simple demonstration is given by the real part of $\log(1 - z/e^{i\theta_c}) = -\sum_{k=1}^{\infty} e^{ik(\theta - \theta_c)}/k$ when $z = e^{i\theta}$. When $|z| < 1$ and z close to $e^{i\theta}$, $1 - z/e^{i\theta_c}$ is almost pure imaginary, and the complex logarithm is about $i\pi/2 + \log|\theta - \theta_c|$ when $\theta < \theta_c$, and $-i\pi/2 + \log|\theta - \theta_c|$ otherwise, so, a logarithm in the real part corresponds to a jump in the imaginary part, and these two kind of singularities create similar asymptotic behaviors in

⁴The Fourier transform in Lighthill's book is written with kernel $\exp(-2\pi ixy)$.

the Fourier coefficients, maybe the work done for a jump [34] can be the basis for a proof of the conjecture 5.2.

Unfortunately, the loose considerations of the preceding subsection suggest to look at the logarithm of the weight function. If the logarithm of a jump (between two positive values) is still a jump, $\log(\log)$ is something new.

6. WEIGHTS WITH LOGARITHMIC SINGULARITIES. ENDPOINT SINGULARITY.

6.1. Numerical tests. B. Danloy [21] considered the generation of orthogonal polynomials of degrees up to N related to $d\mu(x) = -\log x$ on $(0, 1)$ through the exact and stable computation of integrals $J(F) = -\int_0^1 F(x) \log x \, dx$ of some polynomials F of degree $\leq 2N - 1$ by $J(F) = \int_0^1 x^{-1} G(x) dx$, where G is the integral of F vanishing at 0. If G is numerically available everywhere on $[0, 1]$, an N -point Legendre integration formula will do. As $G(x) = \int_0^x F(t) dt = x \int_0^1 F(xu) du$, another Legendre formula, x being now a known value, may be used for $G(x)$ itself.

This technique is probably close to using Legendre modified moments, with $R_n(x)$ = the Legendre polynomial of argument $2x - 1$, as in § 3.2. Powers of logarithms are considered by Sidi [99]. It is then possible to compute safely thousands of recurrence coefficients, as done in table 7.

As the weight function $-\log x$ vanishes at the upper endpoint, we certainly have $\alpha = 1$ in a comparison with the Jacobi weight $(1 - x)^\alpha x^\beta$. With $\beta = 0$, one should have limit values $(\alpha^2 + \beta^2 - 1/2)/4 = 1/8$ and $(\alpha^2 - \beta^2)/2 = 1/2$, so $a_n = 1/4 - \frac{1/32}{n^2} + o(n^{-2})$, $b_n = \frac{1}{2} - \frac{1}{8n^2} + o(n^{-2})$, from (9) when $a = 0$, $b = 1$, and $\alpha = 1$, $\beta = 0$.

The next terms in the asymptotic description of a_n and b_n are suspected to behave like $1/(n^2 \log n)$:

6.2. Conjecture. *If the weight function has no interior singularity, and if $w(x)/[-(x-a)^\beta \log(x-a)]$ and $w(x)/(b-x)^\alpha$ have positive bounded limits at $x = a$ and $x = b$, then there are constants A, A', B , etc. such that*

$$\begin{aligned} a_n &= \frac{b-a}{4} + \frac{A}{n^2} + \frac{B}{n^2 \log n} + \frac{C}{(n \log n)^2} + o((n \log n)^{-2}), \\ b_n &= \frac{a+b}{2} + \frac{A'}{n^2} - \frac{2B}{n^2 \log n} - \frac{2C}{(n \log n)^2} + o((n \log n)^{-2}), \end{aligned} \tag{29}$$

with $A = -(b-a) \frac{\alpha^2 + \beta^2 - 1/2}{16}$, $A' = -\frac{(b-a)(\alpha^2 - \beta^2)}{8}$, as in (9).

Why emphasize these three terms of what is likely an expansion in powers of $1/\log n$? because the simplest test $w(x) = -\log x$ on $(0, 1)$ leads to a $B = 0$ case! With $\alpha = 1$ and $\beta = 0$, $A = -1/32$, $A' = -1/8$ again from (9).

We therefore try to guess the behavior of the corrections ε_n and η_n in $a_n = \frac{1}{4} - \frac{1}{32n^2} + \varepsilon_n$ and $b_n = \frac{1}{2} - \frac{1}{8n^2} + \eta_n$. As logarithmic terms vary so slowly, we only look at powers $n = 2^k$, $k = 2, 3, \dots$. A little amount of luck is always welcome: we first look at $\frac{1}{n^2(a_n - 1/4)}$ to be sure of the $1/32$. That being settled, let $\rho_n = \frac{1}{\frac{1}{n^2(a_n - 1/4)} + 32} \sim -\frac{1}{1024n^2\varepsilon_n}$. Values of ρ_n make the third column of table 7, and it is pleasing to see for $n = 512, 1024, \dots$, the unmistakable quadratic

n	a_n	ρ_n	$-C$	b_n	σ_n	$2C$
0				0.250000000000000000		
1	0.22047927592204921588			0.464285714285714286		
2	0.24224947318051246417			0.485482446456171340		
3	0.24643170234147606161			0.492103081871360859		
4	0.24795568192105814439	0.700546	0.006028	0.495028498758353905	-0.21874	-0.13728
8	0.24947732897332924671	0.474951	0.015806	0.498497801978252749	-0.41642	-0.16225
16	0.24986904694972974076	0.460701	0.025461	0.499581244730037590	-0.75288	-0.15954
32	0.24996748208328437450	0.508006	0.033249	0.499888698235926998	-1.29198	-0.14526
64	0.24999194570788315582	0.592370	0.038811	0.499971199714645076	-2.09634	-0.12892
128	0.24999800446165694617	0.707118	0.042466	0.499992656969610461	-3.20529	-0.11476
256	0.24999950495806105131	0.849780	0.044723	0.499998142922055425	-4.61770	-0.10405
512	0.24999987701903611419	1.018952	0.046047	0.499999532447336313	-6.29481	-0.09660
1024	0.24999996940989087467	1.213451	0.046786	0.499999882584593121	-8.18165	-0.09176
2048	0.24999999238321795744	1.432144	0.047177	0.499999970557423757	-10.2303	-0.08879
4096	0.24999999810192130108	1.673977	0.047368	0.499999992623709105	-12.4114	-0.08710
8192	0.24999999952670709122	1.938042	0.047451	0.499999998153046342	-14.7130	-0.08623
16384	0.24999999988192528940	2.223611	0.047476	0.499999999537711061	-17.1353	-0.08587
32768	0.24999999997053220987	2.530139	0.047473	0.499999999884319298	-19.6838	-0.08581
65536	0.24999999999264358437	2.857242	0.047457	0.499999999971057923	-22.3658	-0.08593

TABLE 7. Data for $-\log x$ on $(0, 1)$: a_n, ρ_n (see text), $(\log 2n)^2/(1024\rho_n)$; $b_n, \sigma_n, (\log n)^2/(64\sigma_n)$.

behavior $\rho_n = 1.02, 1.21, 1.43, 1.67, \dots$, almost one hundredth of the squares of the binary logarithm of n (of $2n$, actually), and $-C$ is estimated from $\frac{(\log 2n)^2}{1024\rho_n}$.

For $b_n, \sigma_n = \frac{1}{\frac{1}{n^2(b_n - 1/2)} + 8} \sim -\frac{1}{64n^2\eta_n}$, and $-2C$ must be the limit of $\frac{(\log n)^2}{64\sigma_n}$ (last

column).

We find here C suspiciously close to the theoretical value $-3/64 = -0.046875$ (see below), but the values seen in the fourth column of table 7 suggest an enormous wave with a crest at 16384 before more oscillations. A similar wave must hold in the last column, we are here not so close to the theoretical value $2C = -3/32 = -0.9375$. One would like to investigate further the dependence of A, B and C on α, β , more powers of logarithms. . .

Stop press: Percy Deift and Oliver Conway [24] sent this to me on March 31, 2017: "For the log weight $-\log(x)$ on $[0, 1]$ we prove that as $n \rightarrow \infty$, the recurrence coefficients a_n, b_n have the following asymptotic behavior: $a_n = 1/4 - 1/(32n^2) + C/(n \log n)^2 + O(1/(n^2(\log n)^3))$, $b_n = 1/2 - 1/(8n^2) - 2C/(n \log n)^2 + O(1/(n^2(\log n)^3))$, where $C = -3/64 \approx -0.047$, which is close to your conjectured value.

To prove this result we use Riemann-Hilbert/steepest-descent methods, but not in the standard way." End of quote.

Note that this also means that $B = 0$ has been proved as well for this weight.

We have at least an interesting relation to the interior singularity conjecture of §5.2 (for which no proof has yet been announced): let $x = a + (b - a)t^2$, then $P_n(x) = (b - a)^n \tilde{P}_{2n}(t)$, where \tilde{P}_{2n} is the even monic orthogonal polynomial of degree $2n$ with respect to the even weight function $|t|w(a + (b - a)t^2)$ on $[-1, 1]$, as well known ([17, chap. 1, §8], etc.), as is the recurrence relation:

$$\tilde{P}_{2n+2}(t) = t \left[\tilde{P}_{2n+1}(t) = t\tilde{P}_{2n}(t) - \tilde{a}_{2n}^2 \frac{\tilde{P}_{2n}(t) + \tilde{a}_{2n-1}^2 \tilde{P}_{2n-2}(t)}{t} \right] - \tilde{a}_{2n+1}^2 \tilde{P}_{2n}(t),$$

so, $a_n = (b-a)\tilde{a}_{2n}\tilde{a}_{2n-1}$, $b_n = a + (b-a)(\tilde{a}_{2n}^2 + \tilde{a}_{2n+1}^2)$. In our example $-\log|t|$ on $(0, 1)$, the \tilde{a} s make the last row of table 3, $\tilde{a}_1^2 = 1/4$, $\tilde{a}_2^2 = 7/36$, $\tilde{a}_3^2 = 17/63 \dots$ and check that sums and products reconstruct the recurrence coefficients of $-\log t$ of table 2. As $\tilde{a}_n = 1/2 + (-1)^n s[\tilde{A}/n + \tilde{B}/(n \log n) + \dots]$ is expected from the conjecture of § 5.2 (here, $c = 0$ and $\theta_c = \pi/2$, $s = \cos \varphi_c = -1$ in (17)), then $O(n^{-2})$ perturbations follow. We must therefore go up to $O(n^{-2})$ terms in the $\tilde{a}_n s$, including the Jacobi polynomials endpoint effect (9)

$$\tilde{a}_n = \frac{1}{2} + s(-1)^n \left[\frac{\tilde{A}}{n} + \frac{\tilde{B}}{n \log n} + \dots \right] - 2 \frac{2\alpha^2 - 1/2}{16n^2} + \dots, \text{ as } |t|w(t^2) \text{ behaves like } |t \mp 1|^\alpha \text{ near}$$

$t = \pm 1$, and where the next term in the brackets is most likely a $1/(n \log^2 n)$ term.

Then, in $a_n = \tilde{a}_{2n}\tilde{a}_{2n-1} = [(\tilde{a}_{2n} + \tilde{a}_{2n-1})^2 - (\tilde{a}_{2n} - \tilde{a}_{2n-1})^2]/4$, and $b_n = \tilde{a}_{2n}^2 + \tilde{a}_{2n+1}^2 = [(\tilde{a}_{2n} + \tilde{a}_{2n+1})^2 + (\tilde{a}_{2n} - \tilde{a}_{2n+1})^2]/2$, $(\tilde{a}_{2n} - \tilde{a}_{2n\pm 1})^2$ is easily found to be $O(n^{-2})$

with the same expansion as in $4 \left[\frac{\tilde{A}}{2n} + \frac{\tilde{B}}{2n \log 2n} + \dots \right]^2 = \frac{\tilde{A}^2}{n^2} + \frac{2\tilde{A}\tilde{B}}{n^2 \log 2n} + \dots$. For $\tilde{a}_{2n} + \tilde{a}_{2n\pm 1}$,

one must take differences into account: $\frac{\tilde{A}}{2n} - \frac{\tilde{A}}{2n \pm 1} \sim \frac{\pm \tilde{A}}{4n^2}$ etc.

$$\tilde{a}_{2n} + \tilde{a}_{2n\pm 1} = 1 - \frac{2\alpha^2 - 1/2}{16n^2} \pm \frac{s\tilde{A}}{4n^2} \pm \frac{s\tilde{B}}{4n^2 \log 2n} + \dots,$$

$$(\tilde{a}_{2n} + \tilde{a}_{2n\pm 1})^2 = 1 + \frac{1/2 - 2\alpha^2 \pm 4s\tilde{A}}{8n^2} \pm \frac{s\tilde{B}}{2n^2 \log 2n} + \dots,$$

$$2a_n \text{ and } b_n \sim \frac{1}{2} + \frac{1/2 - 2\alpha^2 \mp 4s\tilde{A}}{16n^2} \mp \frac{s\tilde{B}}{4n^2 \log 2n} \mp \left[\frac{\tilde{A}^2}{2n^2} + \frac{\tilde{A}\tilde{B}}{n^2 \log 2n} + \frac{2\tilde{A}\tilde{C} + \tilde{B}^2}{2n^2 \log^2 2n} + \dots \right].$$

If $w(t) \approx (x-a)^\beta \log(x-a)$ near a , $\tilde{w}(t) = |t|w(a + (b-a)t^2) \approx |t|^{1+2\beta} \log|t|$ near $t = 0$. Then, with the power $1 + 2\beta$ in conjecture 5.2, $\tilde{A} = |1 + 2\beta|/4$, and

$$2a_n \text{ and } b_n \sim \frac{1}{2} + \frac{1/2 - 2\alpha^2 \mp 2[(2\tilde{A} + s/2 = |\beta|)^2 - 1/4]}{16n^2} \mp \frac{\tilde{B}(4\tilde{A} + s)}{4n^2 \log n} + \dots. \text{ Finally:}$$

$$a_n \sim \frac{1}{4} + \frac{1/2 - \alpha^2 - \beta^2}{16n^2} - \frac{\tilde{B}(|1 + 2\beta| + s)}{8n^2 \log n}, b_n \sim \frac{1}{2} + \frac{\beta^2 - \alpha^2}{8n^2} + \frac{\tilde{B}(|1 + 2\beta| + s)}{4n^2 \log n}.$$

As $\beta = 0$ and $s = -1$ hold for the weight $-\log t$ on $(0, 1)$, the $1/(n^2 \log n)$ term is indeed vanishing. The $1/(n^2 \log^2 n)$ term has not been worked further here...

6.3. Trying multiple orthogonal polynomials. Warning: this subsection deals with a trend which proved not to be fruitful in the present case, but could be inspiring in other situations. Fact is that I do not know any other explicit family of orthogonal (of some sort) polynomials with respect to a logarithmic weight function.

Quite another trend is given by known formulas for some multiple orthogonal polynomials, summarized here [44, § 12], [55, chap. 23], [76]: the polynomial $R_n = R_{\{n_1, \dots, n_p\}}$ of degree $n = n_1 + \dots + n_p$ is a multiple orthogonal polynomial with respect to the measures $d\mu_1, \dots, d\mu_p$ if R_n is orthogonal to polynomials of degree $< n_1$ with respect to $d\mu_1$, of degree $< n_2$ w.r.t. $d\mu_2, \dots$, of degree $< n_p$ w.r.t. to $d\mu_p$. This goes back to Hermite and Padé, and even to Jacobi (Jacobi-Perron algorithm), see [14]. An interesting recurrence relation (13) with $s = p$ occurs when⁵ $n_j = 1 + \lfloor (n-j)/p \rfloor, j = 1, \dots, p$.

⁵The floor $\lfloor x \rfloor =$ the largest integer $\leq x$.

Let $p = 2$, $d\mu_1(x) = x^{\alpha_1}dx$ and $d\mu_2(x) = x^{\alpha_2}dx$ on $(0, 1)$. The corresponding polynomials R_n are explicitly known [3, 58]. As they are orthogonal to polynomials of degree $< \min(n_1, n_2)$ with respect to any linear combination with constant coefficients of $d\mu_1$ and $d\mu_2$, let us take α_1 and $\alpha_2 \rightarrow 0$, then the orthogonality holds with respect to the constant weight and the limit of $\frac{x^{\alpha_2} - x^{\alpha_1}}{\alpha_2 - \alpha_1}$ which is $\log x$ [55, § 23.3.2.1], there we are: R_n does half of the job, as it is orthogonal with respect to the logarithmic weight to polynomials of degree $< n/2$ if n is even, of degree $< (n-1)/2$ if n is odd. We have $R_n(x) = \frac{1}{n_1!n_2!} \frac{d^{n_2}}{dx^{n_2}} \left[x^{n_2} \frac{d^{n_1}}{dx^{n_1}} x^{n_1} (x-1)^n \right]$ [3, §3.3], symmetric in n_1 and n_2 , $R_n(0) = (-1)^n$, $R_n(1) = \frac{n!}{n_1!n_2!}$, $R_0 = 1$, $R_1(x) = 2x - 1$, $R_2(x) = 9x^2 - 8x + 1$, $R_3(x) = 40x^3 - 54x^2 + 18x - 1$, $R_4(x) = 225x^4 - 400x^3 + 216x^2 - 36x + 1$, and the recurrence relation

$$\begin{aligned} xR_n(x) &= \frac{4(n+1)^2(n+2)}{(3n+2)^2(3n+4)}R_{n+1}(x) + \frac{4(n^2+19n/9+1)}{(3n+2)(3n+4)}R_n(x) \\ &\quad + \frac{4n(27n^2-16)}{9(3n-2)(3n+2)^2}R_{n-1}(x) + \frac{4n(n-1)}{3(3n-2)(3n+2)}R_{n-2}(x) \text{ if } n \text{ is even,} \\ &= \frac{4(n+1)}{9(3n+1)}R_{n+1}(x) + \frac{4(9n^2-n-1)}{9(3n-1)(3n+1)}R_n(x) + \frac{4n^2}{3(n+1)(3n+1)}R_{n-1}(x) \\ &\quad + \frac{4n(n-1)^2}{3(3n-1)(3n+1)(n+1)}R_{n-2}(x) \text{ if } n \text{ is odd.} \end{aligned} \quad (30)$$

The vectors of scalar products $\mathbf{v}^{(n)} = [(R_0, P_n), (R_1, P_n), \dots]$ have only a finite number of nonzero elements from (R_n, P_n) to (R_{2n+1}, P_n) .

$$\begin{aligned} \mathbf{v}^{\wedge}\{0\} &= [1 \quad -1/2 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad \dots] \\ \mathbf{v}^{\wedge}\{1\} &= [0 \quad 7/72 \quad -11/144 \quad -1/40 \quad 0 \quad 0 \quad 0 \quad \dots] \\ \mathbf{v}^{\wedge}\{2\} &= [0 \quad 0 \quad 647/25200 \quad -3/175 \quad -89/4900 \quad -1/504 \quad 0 \quad 0 \quad \dots] \end{aligned}$$

Unfortunately, numerical stability for large n is poor, the amplification of the effects of rounding errors is about $2^{n/2}$ after n steps. This may be related to the behavior of $|R_n(x)|$ on $[0, 1]$, increasing from 1 to about 2^n instead of keeping an approximately equal ripple, as orthogonal polynomials do.

Classical multiple orthogonal polynomials through the Sonine-Hahn criterium (the derivatives of the polynomials being multiple orthogonal polynomials themselves) have been studied by Douak & Maroni [25]. See also Van Assche & Coussement [106] for another approach.

7. ABOUT MATRICES IN SOLID-STATE PHYSICS.

7.1. Matrix approximation of the Hamiltonian operator. A solid-state system is a stable arrangement of atoms (whose positions are the *sites*) which may create, or at least amplify, interesting physical phenomena, such as electrical conductivity and even superconductivity, magnetization... The description of the behavior of such a system is achieved by a complex-valued function (called a wavefunction, which will simply be denoted as a *state* hereafter) of the time variable and d space variables, with $d = 1, 2$, or 3 according to our present knowledge of the world. The measure of some physical properties (*observables*), such as the magnetization, is the value of an appropriate linear functional or operator acting on the current state. At any given time, the states are to be found in a subset of $L^2(\mathbb{R}^d)$. This formidable set of functions⁶ is often approximated by the set of linear combinations of a finite set of simple "atomic functions" with a small support around each site, quite similar to finite elements constructions.

⁶And this is only for a single-electron operator, otherwise we should have to consider a power of $L^2(\mathbb{R}^d)$.

A state is therefore a vector \mathbf{u} of, say, N elements representing the relative weight of the sites in the decomposition of the wavefunction. Actually, the state depends on the time t and satisfies the Schrödinger equation

$$\frac{d\mathbf{u}(t)}{dt} = -\frac{i}{\hbar}\mathbf{H}\mathbf{u}(t), \quad (31)$$

where \mathbf{H} is a real symmetric matrix of order N approximating the relevant one-particle Hamiltonian operator of the system. The (m, n) element of \mathbf{H} describes the coupling of energy between sites m and n (self-coupling if $m = n$). This coupling decreases fast if m and n are far apart, and one often considers only nearby values (closest neighbor approximation, or tight-binding approximation see Economou's book [28, §5.2]) also the first pages of Giannozzi & al. [43] and Haydock [49, 50]).

Consider for instance a one-dimensional chain of sites $\{\dots, x_{n-1}, x_n, x_{n+1}, \dots\}$ at distance $x_{n+1} - x_n = \ell$ from each neighbor. If all the interactions between nearest neighbors are identical, we have

$$\mathbf{H} = \begin{bmatrix} \ddots & \ddots & \ddots & & & & \\ & \alpha & \beta & \alpha & & & \\ & & \alpha & \beta & \alpha & & \\ & & & \alpha & \beta & \alpha & \\ & & & & \ddots & \ddots & \ddots \end{bmatrix} \quad (32)$$

In this case like in more realistic systems, the Hamiltonian operator is therefore represented by a huge sparse symmetric matrix where each row is associated to a site and contains a small number of nonzero elements corresponding to neighboring sites (tight-binding approximation).

When the Hamiltonian operator is independent of time, the Schrödinger equation (31) is solved by

$$\mathbf{u}(t) = \exp(-it\mathbf{H}/\hbar)\mathbf{u}(0) = \sum_{r=0}^{\infty} \frac{(-it\mathbf{H})^r \mathbf{u}(0)}{r!\hbar^r}, \quad (33)$$

Starting with an initial state vector $\mathbf{u}(0)$.

Equation of motion method [49, §34]: we examine the diffusion of an electron initially localized on some site, say the m^{th} one, at $x = x_m$, therefore described by a vector $\mathbf{u}(0)$ with a single nonzero element. By applying the evolution operator (33) to this initial state, we have to look at the m^{th} column of a combination of powers of the matrix \mathbf{H} .

On the one-dimensional example given by (32) when $\beta = 0$, the elements of the powers of \mathbf{H} are Laurent coefficients of the same powers of $\alpha z^{-1} + \alpha z$ (symbol of a Toeplitz matrix [72, § 3 and 5]).

So, from the binomial theorem, the diagonal element of \mathbf{H}^{2r} is $\binom{2r}{r}\alpha^{2r} = \frac{(2r)!}{(r!)^2}\alpha^{2r}$ and (33) is

the Bessel function $\sum_{r=0}^{\infty} \frac{(-i\alpha t)^{2r}}{(r!)^2\hbar^{2r}} = J_0(2\alpha t/\hbar)$ at the diagonal element $n = m$, $i^{n-m} J_{n-m}(2\alpha t/\hbar)$

[57, Chap. IX] at the n^{th} element of the lattice, or, with $x = x_m + (n - m)\ell$, and reinstating β through multiplication by $e^{-i\beta t/\hbar}$,

$$\mathbf{u}(t)_n = e^{-i\beta t/\hbar} i^{n-m} J_{|x_n - x_m|/\ell}(2\alpha t/\hbar). \quad (34)$$

See [96] for what seems to be similar calculations, perhaps in [97] too.

Fig. 1 shows absolute values of components of a state vector $\mathbf{u}(t)_{m-4}, \dots, \mathbf{u}(t)_{m+4}$ at various time values, starting with $\mathbf{u}(0)_n = \delta_{n,m}$. The last time value is such that $2\alpha t/\hbar = 2.404825\dots$, the first zero of J_0 [1, p. 409] [57, IX, Table 32a]. Only the values on the lattice are relevant, the linear interpolation in Fig. 1 has no special meaning.




FIGURE 1. Diffusion of a single site occupation in a one-dimensional lattice.

7.2. **Density of states.** We now apply (33), using the eigenvalues and eigenstates $(E_p, \mathbf{v}^{(p)})$ of the simplified Hamiltonian operator \mathbf{H} . We now have

$$\mathbf{u}(t) = \sum_p \exp(-itE_p/\hbar) (\mathbf{v}^{(p)}, \mathbf{u}(0)) \mathbf{v}^{(p)}, \quad (35)$$

where $(\mathbf{v}^{(p)}, \mathbf{u}(0))$ is the usual (complex) scalar product of the two vectors $\mathbf{v}^{(p)}$ and $\mathbf{u}(0)$.

We consider the projection $\overline{\mathbf{u}(t)}_n$ on the n^{th} site starting from the m^{th} site, and rearrange the sum $\sum_p \exp(-itE_p/\hbar) \overline{\mathbf{v}_m^{(p)}} \mathbf{v}_n^{(p)}$, as $\int_a^b \exp(-itE/\hbar) d\mathcal{N}_{m,n}(E)$, where $\mathcal{N}_{m,n}(E)$ is a staircase function discontinuous at each eigenvalue. When $n = m$, $d\mathcal{N}_{n,n}(E)$ is the sum of the positive terms $|\mathbf{v}_n^{(p)}|^2$ for the eigenvalues E_p in an interval of length dE around E . The result $n_{n,n}(E)dE$, where $n_{n,n}(E)$ is called the (local) *density of states*.

By "local", it is meant that the result depends on the initial state $\mathbf{u}(0)$. If this state is concentrated on a single site, or on a small number of neighbors, the resulting $n(E)dE$ is obviously related to this starting site. For perfect crystals, as investigated in subsection 7.4, there is only a small number of non equivalent results. One often wants the average, or total, density of states of the whole system, and this asks for the trace of $(E\mathbf{I} - \mathbf{H})^{-1}$. For perfect crystals, the average density of states is a weighted sum of a small number of non equivalent local densities of states [49, § 23]. In the examples that will be considered in § 8, there is only one local density of states, therefore identical to the average one.

In the example (32), the eigenvalues are $E_p = \beta + 2\alpha \cos(p\pi/N)$ if the matrix has N rows & columns; normalized eigenvectors are $\mathbf{v}_n^{(p)} = \sqrt{2/N} \sin(np\pi/N)$ (average value of the sine squares is 1/2), see [46, chap. 7], [49, §12].

In our example, starting from the m^{th} site, at $x = x_m = m\ell$, (35) becomes at the n^{th} site at time t

$$\begin{aligned} \mathbf{u}(t)_n &= 2 \sum_p \exp(-it[\beta + 2\alpha \cos(p\pi/N)]/\hbar) \sin(mp\pi/N) \sin(np\pi/N)/N \\ &\sim (2/\pi) \int_0^\pi \exp(-it(\beta + 2\alpha \cos \theta)) \sin(\theta x_m/\ell) \sin(\theta x_n/\ell) d\theta \end{aligned} \quad (36)$$

when we let $N \rightarrow \infty$ (continuous spectrum), with $x = n\ell, \theta = p\pi/N$.

If $a = \beta - 2\alpha < E < b = \beta + 2\alpha$, let $E = \beta + 2\alpha \cos \theta_E$, then, between E and $E + dE$, there are $(N/\pi)|\theta_{E+dE} - \theta_E|$ eigenvalues, to multiply by the average of the squares of eigenvector elements, what remains is $\pi^{-1} \left| \arccos\left(\frac{E + dE - \beta}{2\alpha}\right) - \arccos\left(\frac{E - \beta}{2\alpha}\right) \right| \approx \frac{dE}{\pi \sqrt{4\alpha^2 - (E - \beta)^2}}$.

Nobody indulges in such awkward ways! Instead, one considers the *Green functions* [28, 43, 48, 49, 50, 87]

$$G_{m,n}(x) = \int_a^b \frac{n_{m,n}(t) dt}{x-t} = ((x\mathbf{I} - \mathbf{H})^{-1})_{m,n},$$

which, if $m = n$, have the properties of the Stieltjes functions of the first section!

In the example above, (36) becomes at $x = n\ell$,

$$\begin{aligned} \mathbf{u}(t)_n &= \frac{1}{\pi} \int_0^\pi \exp(-it(\beta + 2\alpha \cos \theta)/\hbar) [\cos(\theta(x_n - x_m)/\ell) - \cos(\theta(x_n + x_m)/\ell)] d\theta \\ &= e^{-it\beta/\hbar} [i^{m-n} J_{n-m}(-2\alpha t/\hbar) - i^{m+n} J_{n+m}(-2\alpha t/\hbar)] \end{aligned}$$

[1, §9.1.21] [89, §10.9.2], where we recover (34), considering that $J_{n+m} \rightarrow 0$ when n and m are large: we only consider $x = x_n$ close to x_m .

7.3. The recursion (Lanczos) method. Let $\mathbf{u}^{(0)}$ be a state represented by a vector of \mathbb{R}^N (only the space variables are considered now), and $\mu_n = (\mathbf{u}^{(0)}, \mathbf{H}^n \mathbf{u}^{(0)})$, where $(\ , \)$ is the usual scalar product of \mathbb{R}^N . From the expansion of $\mathbf{u}^{(0)}$ in the orthonormal set of eigenstates $\{\mathbf{v}^{(p)}\}$ as seen above, $\mu_n = \sum_p E_p^n |(\mathbf{u}^{(0)}, \mathbf{v}^{(p)})|^2 = \int_a^b t^n d\mu(t)$, where $d\mu(t)$ is the relevant density of states times dt . As \mathbf{H} is a very sparse matrix, the vectors $\mathbf{H}^n \mathbf{u}^{(0)}$ are easy to compute and they may be rearranged in an orthonormal sequence $\mathbf{u}^{(n)} = p_n(\mathbf{H})\mathbf{u}^{(0)}$ by linear algebra constructions. Of course, this means that $\delta_{m,n} = (\mathbf{u}^{(m)}, \mathbf{u}^{(n)}) = (p_m(\mathbf{H})\mathbf{u}^{(0)}, p_n(\mathbf{H})\mathbf{u}^{(0)}) = (\mathbf{u}^{(0)}, p_m(\mathbf{H})p_n(\mathbf{H})\mathbf{u}^{(0)})$ (from symmetry of \mathbf{H}) $= \int_a^b p_m(t)p_n(t)d\mu(t)$, so $p_n = \kappa_n P_n$ is the orthonormal polynomial of degree n with respect to $d\mu$. Therefore, from the recurrence relation $tp_n(t) = a_n p_{n-1}(t) + b_n p_n(t) + a_{n+1} p_{n+1}(t)$, $\mathbf{H}p_n(\mathbf{H}) = a_n p_{n-1}(\mathbf{H}) + b_n p_n(\mathbf{H}) + a_{n+1} p_{n+1}(\mathbf{H})$, or $\mathbf{H}\mathbf{u}^{(n)} = a_n \mathbf{u}^{(n-1)} + b_n \mathbf{u}^{(n)} + a_{n+1} \mathbf{u}^{(n+1)}$:

$$\mathbf{H}[\mathbf{u}^{(0)} \mid \mathbf{u}^{(1)} \mid \mathbf{u}^{(2)} \mid \dots] = [\mathbf{u}^{(0)} \mid \mathbf{u}^{(1)} \mid \mathbf{u}^{(2)} \mid \dots] \begin{bmatrix} b_0 & a_1 & & & \\ a_1 & b_1 & a_2 & & \\ & a_2 & b_2 & a_3 & \\ & & \ddots & \ddots & \ddots \end{bmatrix}, \quad (37)$$

so that the Hamiltonian matrix and the tridiagonal matrix of the recurrence coefficients have the same spectrum (should we be able to build N vectors $\mathbf{u}^{(n)}$ s). From $\mathbf{u}^{(n-1)}$ (if $n > 0$) and $\mathbf{u}^{(n)}$, one gets a_n and b_n by $a_n = (\mathbf{u}^{(n-1)}, \mathbf{H}\mathbf{u}^{(n)})$, $b_n = (\mathbf{u}^{(n)}, \mathbf{H}\mathbf{u}^{(n)})$ [65] [39, § 3.1.7.1] [45, chap. 9]. For the biography of Cornelius Lanczos (Kornél Löwy), see [88].

The recursion method has been, and still is, quite an inspiration in solid-state physics! [18, 36, 43, 49, 50, 51, 60]. The Hamiltonian operator of a given physical system is approximated by a matrix \mathbf{H} as above, and a set of recurrence coefficients is produced by the Lanczos method. The features of the weight function are then "read" from the asymptotic behavior of these recurrence coefficients.

The reverse procedure is used here: from the known densities of states of model systems, the recurrence coefficients are produced through modified moments, and asymptotic properties are investigated.

7.4. Perfect crystals. A perfect crystal is the repetition of a d -dimensional cell of atoms along d vectors $\mathbf{a}_1, \dots, \mathbf{a}_d$ [83].

If there is a large but finite number of atomic positions (sites), the Hamiltonian operator is a large matrix acting on a vector $v(x_1, \dots, x_d)$ as $\mathbf{H}v$ at the available site $(x_1, \dots, x_d) = \sum_m h_m v(x + \delta_m)$, where each δ_m is a vector relating x to one of its neighbors in its cell and neighboring cells. See the next section for two examples.

Let us try a vector $\exp(ik \cdot x) = \exp(i(k_1x_1 + \dots + k_dx_d))$. The product by \mathbf{H} reproduces the same vector times the scalar function $h(k) = \sum_m h_m \exp(ik \cdot \delta_m)$ which are therefore the eigenvalues of \mathbf{H} , for various real nonequivalent vectors k (Brillouin zone [43, §4]), i.e., such that each $k \cdot \delta_m \in [0, 2\pi)$ or $[-\pi, \pi)$. In mathematician's lingo, $h(k)$ is the symbol of the Toeplitz matrix \mathbf{H} (Grenander & Szegő [47, chap. 5,6, and notes of chap. 5])! More technically, \mathbf{H} of a perfect crystal is a block-Toeplitz matrix with a matrix symbol.

Assuming the eigenvalues to be distributed like the k -vectors (recall the simple 1D case where each k such that $\sin(kN\ell) = 0$ produces an eigenvalue), the number of eigenvalues less than some E is N times the volume $\mathcal{N}(E)$ in the Brillouin zone of the k -vectors such that $h(k) \leq E$, and the Green function of the average (total) density of states is $\text{trace}((x\mathbf{I} - \mathbf{H})^{-1}) = \sum \frac{1}{x - E_p} =$

$$N \int_{E=h(k)=t} \frac{d\mathcal{N}(t)}{x - t} = N \int_{k \in B} \frac{|dk|}{x - h(k)} \text{ with the Brillouin zone } B.$$

So, there is no need to estimate numerically the density of states of a perfect crystal, as the job has been done long ago. But recurrence coefficients found in this ideal case may be useful in later investigations of realistic models of true physical systems.

8. TWO FAMOUS 2-DIMENSIONAL LATTICES.

8.1. The square lattice. The four vectors relating a site to its neighbors are $(\pm\ell, 0), (0, \pm\ell)$, see fig. 2, so that $h(k_1, k_2) = 2 \cos(k_1\ell) + 2 \cos(k_2\ell)$ (multiplied by the relevant physical energy constant, and we also ignore the multiplications by 2 and ℓ).




FIGURE 2. Square lattice: nearest neighbors and density of states.

Then (Economou [28, §5.3.2]),

$$S(x) = G_{0,0}(x) = (\pi)^{-2} \int_0^\pi \int_0^\pi \frac{dk_1 dk_2}{x - \cos k_1 - \cos k_2} = \frac{2}{\pi x} \mathbf{K} \left(\frac{2}{x} \right), \quad (38)$$

where $\mathbf{K}(u) = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - u^2 \sin^2 \theta}} = \int_0^1 \frac{dr}{\sqrt{(1 - r^2)(1 - u^2 r^2)}}$ is the complete elliptic integral of the first kind of modulus u (the $(\pi)^{-2}$ factor is for convenience, so as to have $S(x) \sim 1/x$ for large x , unity as total weight).

Indeed, we integrate in k_2 for a given k_1 , seeing that the integral from 0 to π is half the integral on the circle of $\frac{d\zeta/(i\zeta)}{x - \cos k_1 - (\zeta + 1/\zeta)/2}$, where $\zeta = \exp(ik_2)$, so π times the residue of $-2/[\zeta^2 - 2(x - \cos k_1)\zeta + 1]$ at the pole in the unit disk, and this residue is $\frac{1}{\sqrt{(x - \cos k_1)^2 - 1}}$,

and $\frac{dk_1}{\sqrt{(x - \cos k_1)^2 - 1}} = \frac{\sqrt{1 - \alpha^2} d\xi}{\sqrt{(1 - \xi^2)[x + 1 - \alpha + ((x + 1)\alpha - 1)\xi][x - 1 - \alpha + ((x - 1)\alpha - 1)\xi]}}$

if $\cos k_1 = \frac{\xi + \alpha}{1 + \alpha\xi}$. If α is such that $\frac{(x + 1)\alpha - 1}{x + 1 - \alpha} = -\frac{(x - 1)\alpha - 1}{x - 1 - \alpha}$, we find

$\frac{dk_1}{\sqrt{(x - \cos k_1)^2 - 1}} = \frac{\alpha d\xi}{\sqrt{(1 - \xi^2)(1 - \alpha^4\xi^2)}}$ when $\alpha + \alpha^{-1} = x$, so the result is $2\alpha\pi^{-1}\mathbf{K}(\alpha^2) = \frac{2\alpha}{\pi(1 + \alpha^2)}\mathbf{K}\left(\frac{2\alpha}{1 + \alpha^2}\right)$, from the Gauss-Landen transformation formula (Borwein [10, §2.7],

Jahnke & Emde [57, chap. V, C, §2.2], NIST [89, §19.8]), whence the result (38).

When x is real outside $[-2, 2]$, \mathbf{K} has an argument in $(-1, 1)$ in (38) and is real there.

When $x \in [-2, 2]$, $\frac{2}{\pi x}\mathbf{K}\left(\frac{2}{x}\right) = \frac{1}{\pi} \int_0^{x/2} \frac{dr}{\sqrt{(1 - r^2)(x^2/4 - r^2)}} \pm \frac{i}{\pi} \int_{x/2}^1 \frac{dr}{\sqrt{(1 - r^2)(r^2 - x^2/4)}} = \frac{\text{sign } x}{\pi}\mathbf{K}(x/2) \pm \frac{i}{\pi}\mathbf{K}(\sqrt{1 - x^2/4})$ so⁷, the weight function is [28, eq. 5.39]

$$w(x) = (1/\pi^2)\mathbf{K}(\sqrt{1 - x^2/4}), \quad -2 \leq x \leq 2. \quad (39)$$

Near $x = 0$, using $\mathbf{K}(1 - \varepsilon) \sim (1/2)\log(8/\varepsilon)$ (from [1, 17.3.26] with $m_1 = 1 - (1 - \varepsilon)^2$), the density of states has a $(\log(\sqrt{32}/|x|))/\pi^2$ behavior near the origin.

The power moments are the coefficients of the expansion of $S(x) = \mu_0/x + \mu_1/x^2 + \dots$. From the known expansion of \mathbf{K} [89, §19.5.1] etc., $\mu_{2n+1} = 0$,

$$\mu_{2n} = \left(\frac{1 \times 3 \times \dots (2n - 1)}{n!}\right)^2 = \frac{1}{\pi} \left(\frac{2^n \Gamma(n + 1/2)}{\Gamma(n + 1)}\right)^2 = 1, 1, 9/4, 25/4, 1225/64, \dots$$

As the spectrum is $[-2, 2]$ (the extreme values of $\cos k_1 + \cos k_2$), the Chebyshev moments are here the moments of $T_n(x/2) = 1, x/2, (x^2 - 2)/2, (x^3 - 3x)/2, (x^4 - 4x^2 + 2)/2, \dots$, so $\nu_0 = 1, \nu_2 = -1/2, \nu_4 = 1/8, \nu_6 = -1/8, \dots$ which must of course be computed in a sensible way, as they seem fortunately to be much smaller than the μ_n s. We need an expansion of $S(x)$ in negative powers of $z = x/2 + \sqrt{x^2/4 - 1}$, from (16). By a stroke of luck, $z = 1/\alpha$ used in the proof of (38), so we return to an intermediate result $S(x) = 2/(\pi z)\mathbf{K}(z^{-2})$ and apply (16)

$$\nu_0 + \sum_1^{\infty} \frac{2\nu_n}{z^n} = \frac{2(z - z^{-1})}{\pi z}\mathbf{K}(z^{-2}), \text{ whence } \nu_0 = 1, \nu_2 = -1/2, \text{ and}$$

$$\nu_{4n} = -\nu_{4n+2} = \frac{1}{2^{2n+1}} \left(\frac{1 \times 3 \times \dots (2n - 1)}{n!}\right)^2, \quad n = 1, 2, \dots \quad (40)$$

It is then possible to compute tens of thousands recurrence coefficients with the algorithm of § 3.4. Some of them are given in Table 8. As $\theta_c = \pi/2$, we expect $a_n - a_\infty \approx f_n \cos(n\pi + \varphi_c) = (-1)^n f_n \cos \varphi_c$. There is also a $1/(8n^2)$ Legendre-Jacobi contribution from the endpoints. After subtraction of this $1/(8n^2)$, the $(-1)^n$ behavior is clear on the 10 first items of the table. The amplitude f_n of (17) being thought to decrease like $1/(n \log n)$ from conjecture 5.2, we consider $\rho_n = n \log n (a_n - 1 - 1/(8n^2))$, the limit is reached so slowly that values of ρ_n are shown on powers of 2. Assuming a $A + B/\log n$ behavior, the slope B is estimated from two successive values, and the limit A by $\rho_n - B/\log n$ (Neville extrapolation). With $\{a, b, c\} = \{-2, 2, 0\}$, the conjecture 5.2 expects the limit $1/2$. A second degree extrapolation makes this guess even more credible.

The graph of the weight function in Fig. 2 is established from $\mp \pi w(x) = \text{imaginary part of the limit of } S(x \pm \varepsilon i)$ (Sokhotskyi-Plemelj [53, §14.1]) for x in the spectrum. $S(x)$ is computed

$$\text{as } \frac{1}{x - b_0 - \frac{a_1^2}{x - b_1 - \dots}}$$

⁷use the change of variable $r^2 = 1 - (1 - x^2/4)s^2$ for the second integral.

n	a_n	ρ_n	extr1	slope	extr2
1	1.000000000				
2	1.118033989	0.1203081542			
3	0.974679435	-0.1292279653			
4	1.037456404	0.1643807059			
5	0.983962900	-0.1692895338			
6	1.020931612	0.1876981566			
7	0.988501032	-0.1913801051			
8	1.014170249	0.2032383673			
9	0.991116578	-0.2061868767			
10	1.010573403	0.2146792890			
16	1.005830221	0.2369760223	0.338188987	(-0.280621925)	0.395424284
32	1.002516216	0.2655192494	0.379692157	(-0.395693147)	0.441946913
64	1.001117839	0.2894108176	0.408868658	(-0.496811193)	0.467221659
128	1.000505800	0.3093937275	0.429291186	(-0.581746101)	0.480347508
256	1.000231686	0.3261864584	0.443735575	(-0.651830709)	0.487068739
512	1.000107052	0.3404045226	0.454149035	(-0.709575197)	0.490596148
1024	1.000049789	0.3525479246	0.461838543	(-0.757544840)	0.492596572
2048	1.000023277	0.3630130761	0.467664590	(-0.797927924)	0.493881804
4096	1.000010930	0.3721112300	0.472190923	(-0.832439487)	0.494822587
8192	1.000005151	0.3800864351	0.475788896	(-0.862366588)	0.495577749
16384	1.000002435	0.3871307257	0.478706503	(-0.888656889)	0.496212143
32768	1.000001155	0.3933961897	0.481112684	(-0.912006619)	0.496752864

TABLE 8. Square lattice: values of a_n , $\rho_n = n \log n[a_n - 1 - 1/(8n^2)]$, limit after 1st degree extrapolation, slope, and limit through 2nd degree extrapolation.

$$\frac{a_N^2}{x - b_N - \frac{a_{N+1}^2}{\ddots}} \text{ by } \frac{a_\infty^2}{x - b_\infty - \frac{a_\infty^2}{\ddots}} = a_\infty/z = [x - b_\infty - \sqrt{(x-a)(x-b)}]/2 \text{ which has a well-}$$

defined imaginary part on the two sides of the spectrum $[a, b]$. This is the termination method of Haydock & Nex [51, 87], and Lorentzen, Thron, and Waadeland [20, 68, 102] going back to Wynn [112, 113]. Máté, Nevai, and Totik [78, 85] introduced the use of Turán determinants $p_n^2(x) - p_{n-1}(x)p_{n+1}(x)$ as a way to recover the weight function when n is large. Indeed, when (21a) applies,

$$\begin{bmatrix} p_n(x) & p_{n+1}(x) \\ p_{n-1}(x) & p_n(x) \end{bmatrix} \approx \frac{1}{\sqrt{2\pi}} \begin{bmatrix} -1 & 1 \\ -z^{-1} & z \end{bmatrix} \begin{bmatrix} z^n \exp(\lambda(z^{-1})) & 0 \\ 0 & z^{-n} \exp(\lambda(z)) \end{bmatrix} \begin{bmatrix} -1 & -z \\ 1 & z^{-1} \end{bmatrix},$$

so that the determinant $\approx -(2\pi)^{-1}(z - z^{-1})^2 \exp(\lambda(z^{-1}) + \lambda(z)) = (2a_\infty^2 \pi)^{-1} \sqrt{(x-a)(b-x)}/w(x)$. The termination formula, and the Turán determinants formula are extended to spectra of several intervals = formulas for limit p -periodic continued fractions with $p > 1$ [20, 68, 105]. Turán determinants are also used in unbounded supports cases [100].

Of course, the formulas (38) and (39) are considered too, the complete elliptic integral \mathbf{K} is computed by the AGM method [1, 10, 89].

8.2. Hexagonal lattice: *graphene*.

Graphene is a two-dimensional hexagonal arrangement of carbon atoms, of extreme importance in theoretical and applied physics [16, 27, 59, 64].

One half of the sites (the "A" sites) of the hexagonal arrangement of fig. 3 are related to their neighbors through the three vectors $(1/2 \pm \sqrt{3}/2), (-1, 0)$, and these neighbors make the other half (the "B" sites) with $(-1/2 \pm \sqrt{3}/2), (1, 0)$ Horiguchi [54, §3], Katsnelson [59, §1.2], $h_{A \rightarrow B}(k_1, k_2) = 2e^{ik_1/2} \cos(k_2\sqrt{3}/2) + e^{-ik_1}$, $h_{B \rightarrow A}(k_1, k_2) = 2e^{-ik_1/2} \cos(k_2\sqrt{3}/2) + e^{ik_1}$.




FIGURE 3. Graphene: nearest neighbors and density of states.

Using a matrix symbol for a short while, we see that the Hamiltonian operator acts on a vector $\exp(ik \cdot x)$ where the A -sites and the B -sites are considered separately, as $\begin{bmatrix} 0 & h_{A \rightarrow B}(k) \\ h_{B \rightarrow A}(k) & 0 \end{bmatrix}$, so that the eigenvalues are $E(\xi) = \pm \sqrt{h_{A \rightarrow B}(k)h_{B \rightarrow A}(k)} = \pm \sqrt{4 \cos^2 \xi_2 + 4 \cos \xi_1 \cos \xi_2 + 1}$, where $\xi_1 = 3k_1/2, \xi_2 = k_2\sqrt{3}/2$, and the relevant Green function is

$$\begin{aligned} S(x) = G_{0,0}(x) &= \frac{1}{2\pi^2} \int_0^\pi \int_0^\pi \left[\frac{1}{x - E(\xi)} + \frac{1}{x + E(\xi)} \right] d\xi_1 d\xi_2 \\ &= \frac{x}{\pi^2} \int_0^\pi \int_0^\pi \frac{d\xi_1 d\xi_2}{x^2 - 4 \cos^2 \xi_2 - 4 \cos \xi_1 \cos \xi_2 - 1} \\ &= \frac{\sqrt{ux}}{\pi} \mathbf{K}(u), \text{ where } u = \frac{x^4 - 6x^2 - 3 - \sqrt{(x^2 - 1)^3(x^2 - 9)}}{8x}. \end{aligned} \quad (41)$$

The last formula [54] is established by a first integral in $\xi_1 = -i \log \zeta$ so that we integrate $-id\zeta/[(x^2 - 4 \cos^2 \xi_2 - 1)\zeta - 2(\zeta^2 + 1) \cos \xi_2]$ on the unit circle, and we integrate the residue on

$$\begin{aligned} \xi_2 \text{ as } &\frac{x}{\pi} \int_0^\pi \frac{d\xi_2}{\sqrt{(x^2 - 1 - 4 \cos^2 \xi_2)^2 - 16 \cos^2 \xi_2}} \\ &= \frac{2x}{\pi} \int_0^{\pi/2} \frac{d\xi_2}{\sqrt{x^4 - 6x^2 + 1 - 4(x^2 - 1) \cos(2\xi_2) + 4 \cos^2(2\xi_2)}} \\ &= \frac{x}{\pi} \int_{-1}^1 \frac{d \cos(2\xi_2)}{\sqrt{[1 - \cos^2(2\xi_2)][x^4 - 6x^2 + 1 - 4(x^2 - 1) \cos(2\xi_2) + 4 \cos^2(2\xi_2)]}}. \end{aligned}$$

As before, we change the variable $\cos(2\xi_2) = \frac{\eta + \alpha}{1 + \alpha\eta}$, resulting in

$$\frac{x}{\pi} \int_{-1}^1 \frac{d\eta}{\sqrt{[1 - \alpha^2 \eta^2] \sqrt{[1 - \eta^2][(x^4 - 6x^2 + 1)(1 + \alpha\eta)^2 - 4(x^2 - 1)(1 + \alpha\eta)(\eta + \alpha) + 4(\eta + \alpha)^2]}}}$$

We keep only even powers of η if $\alpha + \alpha^{-1} = (x^2 - 5)/2$, and we have then

$$\frac{\alpha x}{\pi \sqrt{1 + 2\alpha}} \int_{-1}^1 \frac{d\eta}{\sqrt{[1 - \eta^2][1 - u^2 \eta^2]}}, \text{ with } u^2 = \alpha^3 \frac{2 + \alpha}{1 + 2\alpha} = \left(\frac{\alpha(2 + \alpha)}{x} \right)^2 = \left(\frac{\alpha^2 x}{1 + 2\alpha} \right)^2. \text{ Note}$$

that $\alpha = (\sqrt{x^2 - 1} - \sqrt{x^2 - 9})^2/8 \sim 2/x^2$ and $u = (\alpha + 2\alpha^2)/x$

$= [x^4 - 6x^2 - 3 - \sqrt{(x^2 - 1)^3(x^2 - 9)}]/(8x) = 8x/[x^4 - 6x^2 - 3 + \sqrt{(x^2 - 1)^3(x^2 - 9)}] \sim 4/x^3$ when x is large.

The properties of the density of states in fig. 3 follow from the last line of (41) as we follow the imaginary part of $S(x - i\varepsilon)$: u is nonreal when $x \in (1, 3)$ and $(-3, -1)$. For x between -1 and 1 , u is real but outside $(-1, 1)$. Near $x = 1$, $u = -1 + i(x - 1)^{3/2} + \dots$, $\mathbf{K}(u) \sim \log(4/\sqrt{1 - u^2}) \sim (-3/4) \log|x - 1| + \text{constant}$, [1, 17.3.26][57, chap. V, §C.1 and 3], $S(x) \sim -(3i)/(4\pi) \log|x - 1| + \text{constant}$. Near $x = 3$, the limit imaginary part of \mathbf{K} is $\pi/4$, so $\sqrt{3}/4 = 0.433013\dots$ for S . Near $x = 0$, $u \sim -3/(4x)$, $\mathbf{K}(u) \sim \pi/(2u)$ (from the Gauss-Landen formula seen above), and $\text{Im}(S(x)) \sim |x|/\sqrt{3}$.

Check of first power moments: μ_n is the constant Fourier coefficient of the power $(4 \cos^2 \xi_2 + 4 \cos \xi_1 \cos \xi_2 + 1)^n$, $S(x) = \frac{1}{x} + \frac{3}{x^3} + \frac{15}{x^5} + \frac{93}{x^7} + \frac{639}{x^9} + \dots = \frac{1}{x} \left| \frac{3}{x} \right| \frac{2}{x} \left| \frac{3}{x} \right| \frac{5/3}{x} \left| \frac{44/15}{x} \right| \frac{393/220}{x} \dots$. The Chebyshev moments are the moments of $T_n(x/3) = 1, x/3, 2x^2/9 - 1, 4x^3/27 - x, 8x^4/81 - 8x^2/9 + 1, \dots$ as the spectrum is $[-3, 3]$ (extreme real values of \pm the square root of $4 \cos^2 \xi_2 + 4 \cos \xi_1 \cos \xi_2 + 1 = (\cos \xi_1 + 2 \cos \xi_2)^2 + \sin^2 \xi_1$), so $\nu_0 = 1, \nu_2 = -1/3, \nu_4 = -5/27, \dots$. More instances are $\nu_6 = 47/243, \nu_8 = -167/729, \nu_{10} = 1013/6561$.

From (16), $\nu_0/2$ and $\nu_n, n > 0$, = the coefficients of z^{-n} of $\frac{3(z - z^{-1})}{4} S \left(x = \frac{3(z + z^{-1})}{2} \right) = \frac{3(z - z^{-1})}{4} \left[\frac{2}{3(z + z^{-1})} + 3 \left(\frac{2}{3(z + z^{-1})} \right)^3 + \dots \right] = \frac{1}{2} - \frac{1}{3z} + \dots$

How to compute accurately a very large set of these coefficients? A recurrence relation is an invaluable tool for efficient and economical computation of a sequence. Of course, one must be lucky, or clever, enough to find such a relation. For instance, Piessens & *al.* [91, 92], Milovanović & Cvetković [82] find recurrence relations for examples of Chebyshev modified moments.

An important family of recurrence relations is found for sequences of Taylor (or Laurent, or Frobenius) coefficients of solutions of linear differential equations with rational coefficients (Laplace method, see Milne-Thomson [81, chap. 15], Bender & Orszag [9, § 3.2, 3.3], Andrews & *al.* [2, App. F]). Let $F(x) = \sum_0^\delta c_n x^n$ be a solution of the differential equation $\sum_{m=0}^\delta X_m(x) d^m F(x)/dx^m = \sum_0^\infty \alpha_n x^n$, where $X_m(x)$ is the polynomial $\sum_{p=0}^d \chi_{m,p} x^p$, and where the right-hand side is a known expansion. Then, substituting the unknown expansion $\sum c_r x^r$ of $F(x)$ into the differential equation,

$\sum_{m=0}^\delta X_m(x) [\sum_{r=m}^\infty r(r-1) \dots (r-m+1) c_r x^{r-m}] = \sum_0^\infty \alpha_n x^n$, and we gather the terms contributing to the x^n power:

$$\sum_{m=0}^\delta \sum_{p=0}^d \chi_{m,p} (n+m-p)(n+m-p-1) \dots (n-p-1) c_{n+m-p} = \alpha_n,$$

$n = 0, 1, \dots$, which is the sought recurrence relation involving $c_{n+\delta}, \dots, c_{n-d}$. Chebyshev expansions themselves are considered by Fox & Parker [35, Chap. 5], Paszkowski [90], see also the 'Chebfun' package [26].

We apply this programme to the coefficients of the expansion

$$3(z - z^{-1})S(x)/4 = 3(z - z^{-1})\sqrt{ux}\mathbf{K}(u)/(4\pi) = \frac{\nu_0}{2} + \sum_{n=1}^{\infty} \frac{\nu_{2n}}{z^{2n}} \quad (42)$$

where u is the algebraic function $\frac{x^4 - 6x^2 - 3 - \sqrt{(x^2 - 1)^3(x^2 - 9)}}{8x}$, and where $x = 3(z + 1/z)/2$.

First, $u(1 - u^2)d^2\mathbf{K}/du^2 + (1 - 3u^2)d\mathbf{K}/du - u\mathbf{K} = 0$ [89, § 19.4.8] turns into the more beautiful $\frac{d^2(\sqrt{u}\mathbf{K})}{du^2} - 2\frac{u}{1 - u^2}\frac{d(\sqrt{u}\mathbf{K})}{du} + \frac{\sqrt{u}\mathbf{K}}{4u^2} = 0$.

Next, u is the root of $u + 1/u = (x^4 - 6x^2 - 3)/(4x)$ behaving as $4/x^3$ for large x , we translate in z from $x = 3(z + 1/z)/2$, leading to the rather formidable

$$u^{\pm 1} = \frac{27z^4 + 36z^2 + 2 + 36z^{-2} + 27z^{-4} \mp (z - z^{-1})(9z^2 + 14 + 9z^{-2})^{3/2}}{64(z + z^{-1})} \quad (43)$$

So, we differentiate $u + u^{-1} = \frac{27z^4 + 36z^2 + 2 + 36z^{-2} + 27z^{-4}}{32(z + z^{-1})}$ as

$$(1 - u^{-2})du = \frac{81z^5 + 171z^3 + 106z - 106z^{-1} - 171z^{-3} - 81z^{-5}}{32z(z + z^{-1})^2} dz, \text{ or} \quad (44)$$

$$\frac{du}{u dz} = \frac{(z^2 - 1)(9z^2 + 14 + 9/z^2)^2}{32(z^2 + 1)^2(u - u^{-1})} = -\frac{(9z^2 + 14 + 9/z^2)^{1/2}}{z^2 + 1},$$

which is already better looking than before, and we build the differential equation for $\sqrt{u} \mathbf{K}$ in the slightly modified form $u \frac{d}{du} \left[u \frac{d\sqrt{u} \mathbf{K}(u)}{du} \right] - \frac{u + u^{-1}}{u - u^{-1}} u \frac{d\sqrt{u} \mathbf{K}(u)}{du} + \frac{\sqrt{u} \mathbf{K}(u)}{4} = 0$ with respect to the variable z :

$$\frac{z^2 + 1}{(9z^2 + 14 + 9/z^2)^{1/2}} \frac{d}{dz} \left[\frac{z^2 + 1}{(9z^2 + 14 + 9/z^2)^{1/2}} \frac{d\sqrt{u} \mathbf{K}}{dz} \right]$$

$$+ \frac{27z^4 + 36z^2 + 2 + 36z^{-2} + 27z^{-4}}{(z - z^{-1})(9z^2 + 14 + 9/z^2)^{3/2}} \frac{z^2 + 1}{(9z^2 + 14 + 9/z^2)^{1/2}} \frac{d\sqrt{u} \mathbf{K}}{dz} + \frac{\sqrt{u} \mathbf{K}}{4} = 0.$$

Finally, we turn to the full function of (16) and (41), say $F(z) = (z - z^{-1})S(x) = \text{constant}$ times $(z - z^{-1})\sqrt{ux} \mathbf{K}(u)$ by substituting $\sqrt{u} \mathbf{K}(u)$ into constant $(z - z^{-1})^{-1}(z + z^{-1})^{-1/2} S$

$$\frac{z^2 + 1}{(9z^2 + 14 + 9/z^2)^{1/2}} \frac{d}{dz} \left[\frac{z^2 + 1}{(9z^2 + 14 + 9/z^2)^{1/2}} \frac{d(z - z^{-1})^{-1}(z + z^{-1})^{-1/2} F}{dz} \right]$$

$$+ \frac{(z^2 + 1)(27z^4 + 36z^2 + 2 + 36z^{-2} + 27z^{-4})}{(z - z^{-1})(9z^2 + 14 + 9/z^2)^2} \frac{d(z - z^{-1})^{-1}(z + z^{-1})^{-1/2} F}{dz}$$

$$+ \frac{(z - z^{-1})^{-1}(z + z^{-1})^{-1/2} F}{4} = 0, \text{ so}$$

$$(z^2 + 1)^2(9z^2 + 14 + 9z^{-2})(z - z^{-1})^2 d^2 F/dz^2$$

$$+ (z^2 + 1)(9z^4 - 14z^2 - 72 - 42z^{-2} - 9z^{-4})(z - z^{-1}) dF/dz$$

$$+ 8(3z^4 + 18z^2 + 22 + 18z^{-2} + 3z^{-4})F = 0$$

Now, put $F(z) = \nu_0/2 + \sum_1^\infty \frac{\nu_{2n}}{z^{2n}}$ in the differential equation $(9z^8 + 14z^6 - 9z^4 - 28z^2 - 9 + 14z^{-2} + 9z^{-4}) \sum 2n(2n+1)\nu_{2n}z^{-2n-2} - (9z^7 - 14z^5 - 81z^3 - 28z + 63z^{-1} + 42z^{-3} + 9z^{-5}) \sum 2n\nu_{2n}z^{-2n-1} + 8(3z^4 + 18z^2 + 22 + 18z^{-2} + 3z^{-4})[\nu_0/2 + \sum \nu_{2n}z^{-2n}] = 0$, and consider the contributions to z^{-2n}

$$9(n+3)^2\nu_{2n+6} + 2(7n^2 + 35n + 45)\nu_{2n+4} - 9(n^2 - 2n - 7)\nu_{2n+2} - 4(7n^2 - 11)\nu_{2n}$$

$$- 9(n^2 + 2n - 7)\nu_{2n-2} + 2(7n^2 - 35n + 45)\nu_{2n-4} + 9(n-3)^2\nu_{2n-6} = 0, n = 0, 1, \dots \quad (45)$$

starting at $n = -2$ with $\nu_0 = 1/2$ (which is actually $\nu_0/2$) and $\nu_n = 0$ for $n < 0$ (the $\nu_0/2$ anomaly can be relieved if we define $\nu_{-n} = \nu_n$, remark indeed that the coefficient of ν_{2n+k} is the coefficient of ν_{2n-k} with $n \rightarrow -n$). The next items are $-1/3, -5/27, 47/243, -167/729, 1013/6561, \dots$ as already seen before, but the recurrence relation (45) allows now to compute incredibly easily any number of these coefficients, one could get one million of them if needed!

Actual computation of a large number of moments by using (45) in the obvious way may lead to numerical disaster, should the wanted solution ν_{2n} be *dominated* by another solution, say, ω_{2n} of the same recurrence (45), i.e., if $\nu_{2n}/\omega_{2n} \rightarrow 0$ when $n \rightarrow \infty$ [110, § 2.6]. Numerical approximations of starting values ν_0 , etc. lead to the actual production of some combination $\nu_{2n} + \varepsilon\omega_{2n}$ and this becomes hopeless for large n , however small ε is. As the starting values are rational numbers in our case, computations with rational numbers avoid the problem, but calculations are space and time consuming for large n . See also the discussions of minimal solutions by Gautschi [38], and in the preface⁸ of Abramowitz and Stegun's handbook [1].

Any solution ω_{2n} of the seven terms recurrence relation (45) can be investigated through its generating function $G(z)$ satisfying the same differential equation as the equation for $F(z)$ seen above. This equation has singular points at $\pm 1, \pm i$, and $(-7 \pm 4\sqrt{2}i)/9 = \exp(\pm i\theta_c)$, all of them on the unit circle. The Taylor expansion of G has therefore a unit convergence radius, and all the solutions ω_{2n} have a less than exponential growth. This can also be seen through contour integral formulas for the Taylor coefficients (Laplace's method, [81, chap. XV]). More details can be found from asymptotic estimation of integrals [9, chap. 6].

n	a_n	ρ_n	slope	extrap.	φ/π	ρ_{n-1}/ρ_n
1	1.732050807568877	0				
2	1.414213562373095	0.33595				
3	1.732050807568877	-0.12782				
4	1.290994448735806	-0.18423				
5	1.712697677155351	0.44419				
6	1.336549152243806	-0.46936				
7	1.628436152438179	0.23792				
8	1.419330149577324	0.16886				
9	1.556750628851699	-0.57145				
10	1.460678158360451	0.77835				
11	1.544107839921112	-0.67587				
12	1.448901939699682	0.30117				
13	1.564500085917823	0.19001				
14	1.431783794658055	-0.57642				
15	1.567701045607340	0.68518				
135	1.507049663059332	0.98260	(0)	0.98260	0.7932	-0.7722
1383	1.500654249446685	1.11873	(-2.07548)	1.40571	0.7922	-0.7751
2007	1.500299317783996	-1.13581	(-2.11734)	1.41425	0.7918	-0.7767
4087	1.500149457578103	-1.15763	(-2.09351)	1.40939	0.7902	-0.7795
8210	1.499924544322330	1.17608	(-2.08240)	1.40713	0.7920	-0.7760
16077	1.500038998496210	-1.19159	(-2.07718)	1.40606	0.7910	-0.7780
34062	1.499974587460581	-1.20647	(-2.07210)	1.40503	0.7920	-0.7757
61370	1.499985980208882	-1.21710	(-2.07231)	1.40507	0.7910	-0.7778

TABLE 9. Graphene: values of $n, a_n, \rho_n = n \log n [a_n - 3/2 + (3/4)(-1)^n/n - (3/16)/n^2]$, (slope), extrap., $\varphi/\pi, \rho_{n-1}/\rho_n$.

The main asymptotic behavior of our own ν_n follows from $S(x) \sim -(3i)/(4\pi) \log|x-1| + \text{constant}$ near $x = \pm 1$, seen above, so, near $z = \pm z_0^{\pm 1} = \pm \exp(\pm i\theta_0)$, where $\cos\theta_0 = 1/3$, $z_0 = (1 + 2i\sqrt{2})/3$, from (16): $(6/8)(z - 1/z)S(x) \sim \pm(3\sqrt{2}/4\pi) \log(z \pm z_0^{\pm 1})$ whence $\nu_n = 0$

⁸Masterpieces deserve to be read entirely, including the preface!

when n is odd, $\nu_n \sim 2 \frac{3\sqrt{2} z_0^n + z_0^{-n}}{4\pi n} = \frac{3\sqrt{2} \cos(n\theta_0)}{\pi n}$ when n is even. For instance, $\nu_{1000000} = -0.2197681875531559 \cdot 10^{-6}$ and $(3\sqrt{2}/\pi) \cos(10^6 \theta_0) = -0.2197654658865520$.




FIGURE 4. Graphene: 100 first $\rho_n = n \log n [a_n - 3/2 + (3/4)(-1)^n/n - (3/16)/n^2]$.

Do we really need the recursion method? The plain Chebyshev, or Fourier, coefficients are already so efficient that they can be preferred, as discussed by Weisse & al. [108, §V.B.2, Table II]. See also Prevost [93] on weight reconstruction with Chebyshev moments.

Anyhow, computing recurrence coefficients from modified moments yields the main behavior $a_n \rightarrow a_\infty = (b-a)/4 = 1.5$ as expected from (8). The next correction is given by the $|x|^\alpha$ behavior near $x = 0$, with $\alpha = 1$, and is $a_n \sim a_\infty - ((b-a) \cos(n\pi)|\alpha|)/(8n) = 1.5 - 0.75(-1)^n/n$. The Jacobi-Legendre effect of the endpoints ± 3 is $(b-a)/(32n^2) = 3/(16n^2)$. The remaining behavior times $n \log n$ believed to be the right factor $\rho_n = n \log n [a_n - 1.5 + 0.75(-1)^n/n - 0.1875/n^2]$ is shown in table 9, and is expected to behave like a constant times $\cos(2n\theta_c + \varphi_c)$. Starting with a large index (here, 135), only crest values are retained, i.e., such that $2n\theta_c + \varphi_c$ happens to be very close to an integer multiple of π . The neighboring values ρ_{n-1} and ρ_{n+1} are then very close together, that's how the interesting values of n are selected. Moreover, the almost common value of ρ_{n-1}/ρ_n and ρ_{n+1}/ρ_n must then be almost $\cos(2\theta_c) = -7/9 = -0.7777\dots$, checked in the last column of table 9 (the first approximated crest is at $n = 10$, see also Fig. 4), whereas φ/π is estimated through the fractional part of $2n\theta_c/\pi$ for these very particular values of n , of which only those in approximate geometric progression have been selected, in the hope to have a better view of the limit of $|\rho_n|$. Although the phase φ is very stable, the evolution of $|\rho_n|$ towards its limit is again excruciatingly slow. An approximate law $A + B/\log n$ is again assumed with the slope B estimated from two successive values, and A as $|\rho_n| - B/\log n$ (extrapolated value). With two contributions at ± 1 in the spectrum $[-3, 3]$, the formula from 5.2 amounts to expecting $\sqrt{2} = 1.414\dots$ which is neither close nor far from the numerical estimate $1.405\dots$

The first 69999 recurrence coefficients a_1, \dots, a_{69999} are given in the file <http://perso.uclouvain.be/alphonse.magnus/graphene69999.txt> of size about 2M, with a precision of 25 digits, in the following format:

```
1.7320508075688772935274463,
1.4142135623730950488016887,
1.7320508075688772935274463,
1.2909944487358056283930885,
1.7126976771553505360155865,
...
1.4999899549568860858094207,
1.5000093070617496121273346,
```


9. ACKNOWLEDGEMENTS.

Many thanks to P. Deift, F. Ducastelle, J.P. Gaspard, Ph. Lambin, P. Nevai, L. Paquet, A. Ronveaux, the referees, and to the people of the Université catholique de Louvain who still accept retired people in the Institute.

REFERENCES

- [1] M. Abramowitz, I.A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards Applied Mathematics Series, **55**, 1964 = Dover.
- [2] G.E. Andrews, R. Askey, R. Roy, *Special Functions*, Cambridge U.P., 1999.
- [3] A.I. Aptekarev, A. Branquinho, W. Van Assche, Multiple orthogonal polynomials for classical weights, *Trans. A.M.S.* **355** (2003) 3887-3914
- [4] A.I. Aptekarev, W. Van Assche, Scalar and matrix Riemann-Hilbert approach to the strong asymptotics of Padé approximants and complex orthogonal polynomials with varying weight, *Journal of Approximation Theory* **129** (2004) 129-166
- [5] I. Area, E. Godoy, A. Ronveaux, A. Zarzo, Hypergeometric-type differential equations: second kind solutions and related integrals *Journal of Computational and Applied Mathematics* **157** (2003) 93-106
- [6] I. Area, E. Godoy, A. Ronveaux, A. Zarzo, Hypergeometric type q -difference equations: Rodrigues type representation for the second kind solution, *Journal of Computational and Applied Mathematics* **173** (2005) 81-92
- [7] W. Barrett, An asymptotic formula relating to orthogonal polynomials, *J. London Math. Soc. (2)* **6** (1973), 701-704.
- [8] B. Beckermann, E. Bourreau, How to choose modified moments? *J. Comp. Appl. Math.* **98** (1998) 81-98.
- [9] C. M. Bender, S. A. Orszag, *Advanced mathematical methods for scientists and engineers*, McGraw-Hill, 1978.
- [10] J.M. Borwein, P.B. Borwein, *Pi and the AGM*, Wiley, 1987.
- [11] J.P. Boyd, *Chebyshev and Fourier Spectral Methods*, 2nd ed., Dover, 2001.
- [12] C. Brezinski, The life and work of André Cholesky. Numer. Algorithms 43 (2006), no. 3, 279-288.
- [13] C. Brezinski, D. Tournès, *André-Louis Cholesky, Mathematician, Topographer and Army Officer*, Birkhäuser, 2014.
- [14] M.G. de Bruin, Simultaneous Padé approximation and orthogonality, pp. 74-83 in C. Brezinski & *al.*, eds: *Polynômes Orthogonaux et Applications Proceedings of the Laguerre Symposium held at Bar-le-Duc, October 15-18, 1984*, Lecture Notes in Mathematics **1171**, Springer, 1985.
- [15] A. Bultheel, M. Van Barel, *Linear algebra, rational approximation and orthogonal polynomials*, Studies in Computational Mathematics, **6**, North-Holland Publishing Co., Amsterdam, 1997.
- [16] A.H. Castro Neto, F. Guinea, N.M.R. Peres, K.S. Novoselov, A.K. Geim, The electronic properties of graphene, *Rev. Mod. Phys.*, **81** (2009) 109-162.
- [17] T. S. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon and Breach, 1978.
- [18] A. Cordelli, G. Grosso, G. Pastori Parravicini, Physical and numerical aspects in Lanczos and modified Lanczos calculations, *Computer Physics Communications* **83** (1994) 255-265.
- [19] J.R. Cruz, Z. Banjanin, Applications of the quotient-difference algorithm to modern spectral estimation, *in Linear Algebra in Signals, Systems, and Control (Boston, MA, 1986)*, organized by B.N. Datta, 343-354, SIAM, Philadelphia, PA, 1988.
- [20] A. Cuyt, V. Petersen, B. Verdonk, H. Waadeland, W. B. Jones, *Handbook of Continued Fractions for Special Functions*, Springer 2008, softcover reprint Dec. 2010.
- [21] B. Danloy, Numerical construction of Gaussian quadrature formulas for $\int_0^1 (-\log x) \cdot x^\alpha \cdot f(x) \cdot dx$ and $\int_0^\infty E_m(x) \cdot f(x) \cdot dx$. *Math. Comp.* **27** (1973), 861-869.
- [22] Ph. J. Davis, Ph. Rabinowitz, *Methods of Numerical Integration*, 2nd ed., Ac. Press 1984.
- [23] P. Deift, *Orthogonal Polynomials and Random Matrices: A Riemann-Hilbert Approach*, in *Courant Lecture Notes*, Vol. **3**, New York University, New York, 1999.
- [24] P. Deift and O. Conway, On the recurrence coefficients for polynomials orthogonal with respect to a log weight, in preparation.
- [25] K. Douak, P. Maroni, Une caractérisation des polynômes d -orthogonaux "classiques", *Journal of Approximation Theory*, Volume **82**, Issue 2, August 1995, Pages 177-204.
- [26] T. A. Driscoll, N. Hale, and L. N. Trefethen, editors, *Chebfun Guide*, Pafnuty Publications, Oxford, 2014.
- [27] F. Ducastelle: Electronic structure of vacancy resonant states in graphene: a critical review of the single vacancy case, *Phys. Rev. B* **88** 075413 10 pages (2013), arXiv:1305.2690v1 [cond-mat.mtrl-sci] 13 May 2013, <http://arxiv.org/abs/1305.2690>
- [28] E.N. Economou, *Green's Functions in Quantum Physics*, Springer, Second Edition 1983, Third Ed. 2006.

- [29] A. Erdélyi *et al.*, *Higher Transcendental Functions*, vol. **2**, McGraw-Hill, 1953.
- [30] A. Erdélyi, *Asymptotic Expansions*, Dover, 1956.
- [31] A.S. Fokas, A.R., Its, A. V. Kitaev, : Discrete Painlevé equations and their appearance in quantum gravity, *Comm. Math. Phys.* **142** (1991), 313–344.
- [32] B. Fornberg, *A Practical Guide to Pseudospectral Methods*, Cambridge University Press, 1996.
- [33] A. Foulquié Moreno, A. Martínez-Finkelshtein, V.L. Sousa, On a conjecture of A. Magnus concerning the asymptotic behavior of the recurrence coefficients of the generalized Jacobi polynomials, *J. Approx. Theory* **162** (2010) 807-831.
- [34] A. Foulquié Moreno, A. Martínez-Finkelshtein, V. L. Sousa, Asymptotics of Orthogonal Polynomials for a Weight with a Jump on $[-1, 1]$ *Constr. Approx.* **33**, No. 2, 219-263 (2011)
- [35] L. Fox, I. B. Parker, *Chebyshev Polynomials in Numerical Analysis*, Oxford U.P. 1968.
- [36] J.P. Gaspard, F. Cyrot-Lackmann, Density of states from moments. Application to the impurity band, *J. Phys. C: Solid State Phys.* **6** (1973) 3077-3096.
- [37] W. Gautschi, On the Construction of Gaussian Quadrature Rules from Modified Moments, *Mathematics of Computation*, Vol. **24**, No. 110 (Apr., 1970), pp. 245-260+s1-s52 = §20.2 [GA40] in *Selected Works of W.Gautschi*, vol. 2 (C.Brezinski & A. Sameh, editors), Birkhäuser, 2014.
- [38] W. Gautschi, Minimal Solutions of Three-Term Recurrence Relations and Orthogonal Polynomials, *Mathematics of Computation* **36**, 1981, 547-554 = §16.2 [GA75] in *Selected Works of W.Gautschi*, vol. 2 (C.Brezinski & A. Sameh, editors), Birkhäuser, 2014.
- [39] W. Gautschi, *Orthogonal Polynomials: Computation and Approximation*, Oxford University Press, 2004.
- [40] W. Gautschi, Gauss quadrature routines for two classes of logarithmic weight functions, *Numer Algor* (2010) **55**, 265–277 = §20.19 [GA198] in *Selected Works of W.Gautschi*, vol. 2 (C.Brezinski & A. Sameh , editors), Birkhäuser, 2014.
- [41] W. Gautschi, *Orthogonal polynomials in MATLAB: Exercises and Solutions*, SIAM, 2016.
- [42] F. Gesztesy, B. Simon, m -functions and inverses spectral analysis for finite and semi-infinite Jacobi matrices, *Journal d'Analyse Mathématique* **73**(1997) 267-297.
- [43] P. Giannozzi, G. Grosso and G. Pastori Parravicini, Theory of electronic states in lattices and superlattices, *Riv. Nuovo Cimento* **13** nr 3 (1990), 1–80.
- [44] L. Golinskii, V. Totik, Orthogonal polynomials: from Jacobi to Simon, pp. 821-874 *in Spectral theory and mathematical physics: a Festschrift in honor of Barry Simon's 60th birthday, Proc. Sympos. Pure Math.*, **76**, Part 2, Amer. Math. Soc., Providence, RI, 2007.
- [45] G.H. Golub, C.F. Van Loan, *Matrix computations*, Johns Hopkins Studies in the Mathematical Sciences, Johns Hopkins University Press, Baltimore, MD, third edition, 1996; fourth edition 2013.
- [46] R.T. Gregory, D.L. Karney, *A Collection of Matrices for Testing Computational Algorithms*, Wiley, 1969.
- [47] U. Grenander, G. Szegő, *Toeplitz Forms and their Applications*, University of California Press, Berkeley, 1958 = Chelsea, 1984.
- [48] G. Grosso, G.P. Parravicini, *Solid state physics*, Academic Press 2000, 2nd printing 2003.
- [49] R. Haydock, Recursive solution of Schrödinger's equation, *Solid State Physics* **35**, Academic P., New York, 1980, 216-294.
- [50] R. Haydock, The recursion method and the Schroedinger equation, pp. 217-228. *in* P. Nevai (ed.), *Orthogonal Polynomials: Theory and Practice*, Proceedings of the NATO Advanced Study Institute on Orthogonal Polynomials and Their Applications Columbus, Ohio, U.S.A. May 22 - June 3, 1989, NATO ASI Series, Series C: Mathematical and Physical Sciences Vol. **294** Kluwer, 1990.
- [51] R Haydock and C M M Nex, A general terminator for the recursion method, *J. Phys. C: Solid State Phys.* **18** (1985) 2235
- [52] P. Henrici, *Applied and Computational Complex Analysis II. Special Functions–Integral Transforms – Asymptotics – Continued Fractions.* , Wiley, N.Y., 1977.
- [53] P. Henrici *Applied and Computational Complex Analysis III. Discrete Fourier Analysis – Cauchy Integrals – Construction of Conformal Maps – Univalent Functions.*, Wiley, N.Y., 1986.
- [54] T. Horiguchi, Lattice Green's Functions for the Triangular and Honeycomb Lattices, *J. Math. Phys.* **13**, 1411-1419 (1972); <http://dx.doi.org/10.1063/1.1666155> (9 pages).
- [55] M. E. H. Ismail, *Classical and Quantum Orthogonal Polynomials in One Variable*, Encyclopedia of Mathematics and its Applications, vol. **98**, Cambridge University Press, 2005 (paperback edition 2009).
- [56] Alexander R. Its, The Riemann-Hilbert Problem and Integrable Systems, *Notices AMS* **50** , N. 11, Dec. 2003, 1389-1400.
- [57] E. Jahnke, F. Emde, F. Lösch, *Tables of Higher Functions, Tafeln höherer Funktionen*, 6th ed., Teubner & McGraw-Hill, 1960.
- [58] V. Kaliaguine, A. Ronveaux, On a system of "classical" polynomials of simultaneous orthogonality, *Journal of Computational and Applied Mathematics* **67** (1996) 207-217.

- [59] M. I. Katsnelson, *Graphene: Carbon in Two Dimensions*, Cambridge University Press 2012.
- [60] E. Koch, The Lanczos method, chap. 10 in E. Pavarini, E. Koch, P. Coleman (Eds.), *Many-Body Physics: From Kondo to Hubbard. Modeling and Simulation*, Vol. 5, Verlag des Forschungszentrum Jülich, 2015.
- [61] A. B.J. Kuijlaars, Riemann-Hilbert Analysis for Orthogonal Polynomials, pp 167-210 in E. Koelink, W. Van Assche (Eds.) : *Orthogonal Polynomials and Special Functions, Lecture Notes in Mathematics 1817*, Springer, 2003.
- [62] A.B.J. Kuijlaars, K.T.-R. McLaughlin, W. Van Assche, M. Vanlessen, The Riemann-Hilbert approach to strong asymptotics for orthogonal polynomials on $[-1, 1]$, *Advances in Mathematics 188* (2004) 337-398.
- [63] Ph. Lambin, J.P. Gaspard, Continued-fraction technique for tight-binding systems. A generalized-moments approach, *Phys. Rev. B 26* (1982) 4356-4368.
- [64] Ph. Lambin, H. Amara, F. Ducastelle, and L. Henrard: Long-range interactions between substitutional nitrogen dopants in graphene: Electronic properties calculations, *Physical Review B 86*, 045448 (2012), <http://hal.archives-ouvertes.fr/hal-00702460/>;
- [65] C. Lanczos, Iteration method for the solution of the eigenvalue problem of linear differential and integral operators, *J. Res. Natl. Bur. Stand. 45*, 255-282 (1950) <http://nvlpubs.nist.gov/nistpubs/jres/045/4/V45.N04.A01.pdf>
- [66] L. Lefèvre, D. Dochain, S. Feye de Azevedo, A. Magnus, Optimal selection of orthogonal polynomials applied to the integration of chemical reactor equations by collocation methods, *Computers & Chemical Engineering* Vol. 24, no. 12, p. 2571-2588 (2000).
- [67] M. J. Lighthill, *Inroduction to Fourier Analysis and Generalised Functions*, Cambridge U.P., 1958.
- [68] L. Lorentzen, H. Waadeland, *Continued fractions. Vol. 1. Convergence theory*. Second edition. Atlantis Studies in Mathematics for Engineering and Science 1, Atlantis Press/World Scientific, 2008.
- [69] D. S. Lubinsky, A survey of general orthogonal polynomials for weights on finite and infinite intervals, *Acta Applicandæ Math. 10* (1987) 237-296.
- [70] D. S. Lubinsky, Asymptotics of Orthogonal Polynomials: Some Old, Some New, Some Identities, *Acta Applicandæ Mathematicæ 61* 207-256, 2000
- [71] A.P. Magnus, Recurrence coefficients for orthogonal polynomials on connected and non connected sets, in *Padé Approximation and its Applications, Proceedings, Antwerp 1979* (L. Wuytack, editor), pp. 150-171, Springer-Verlag (*Lecture Notes Math. 765*), Berlin, 1979.
- [72] A.P.Magnus, Toeplitz matrix techniques and convergence of complex weight Padé approximants, *J.Comp. Appl. Math. 19* (1987) 23-38.
- [73] A.P. Magnus, Asymptotics for the simplest generalized Jacobi polynomials recurrence coefficients from Freud's equations: numerical explorations, *Annals of Numerical Mathematics 2* (1995) 311-325.
- [74] A.P. Magnus, Painlevé-type differential equations for the recurrence coefficients of semi-classical orthogonal polynomials, *J. Comp. Appl. Math. 57* (1-2) (1995) 215-237.
- [75] G. Mansell, W. Merryfield, B. Shizgal, U. Weinert, A comparison of differential quadrature methods for the solution of partial differential equations, *Comput. Methods Appl. Mech. Eng. 104*, 295-316 (1993).
- [76] A. Martínez-Finkelshtein, W. Van Assche, What is A Multiple Orthogonal Polynomial?, *Notices of the AMS 63*, Nr 9, 2016, 1029-1031.
- [77] J.C. Mason, D.C. Handscomb, *Chebyshev polynomials*, Chapman & Hall/CRC, 2003.
- [78] A. Máté, P. Nevai, V. Totik, Extensions of Szegő's theory of orthogonal polynomials., II , *Constr. Approx. 3* (1987),51-72.
- [79] A. Máté, P. Nevai, V. Totik, Twisted difference operators and perturbed Chebyshev polynomials, *Duke Math. J. 57*, No.1, 301-331 (1988).
- [80] R. J. Mathar, Gaussian quadrature of $\int_0^1 f(x) \log^m(x) dx$ and $\int_{-1}^1 f(x) \cos(\pi x/2) dx$, arXiv:1303.5101v1 [math.CA] 20 Mar 2013.
- [81] L. M. Milne-Thomson, *The Calculus Of Finite Differences*, Macmillan And Company., Limited, 1933. <http://www.archive.org/details/calculusoffinite032017mbp>
- [82] G.V. Milovanović & A.S. Cvetković, Logarithmic modification of the Jacobi weight function, *Studia Univ. "Babeş-Bolyai", Mathematica* , Vol. LII, Nr 4, 143-153, Dec. 2007.
- [83] J.W. Morris, Jr., *The Structure and Properties of Materials*, McGraw Hill, 2005.
- [84] P. Nevai, Géza Freud, orthogonal polynomials and Christoffel functions. A case study. *J. Approx. Theory*, 48 (1986), pp. 3-167.
- [85] P. Nevai, Orthogonal Polynomials, Recurrences, Jacobi Matrices, and Measures, pp. 79-104 in A.A. Gonchar and E.B. Saff, eds., *Progress in Approximation Theory*, Springer, 1992.
- [86] P. Nevai, W. Van Assche, Compact perturbations of orthogonal polynomials, *Pacific J. Math. 153* (1992) 163-184.

- [87] C. M. M. Nex, Computing Greenians: Quadrature and Termination, pp. 52-60 in D.G. Pettifor & D.L. Weaire, editors: *The Recursion Method and Its Applications*, Springer Series in Solid-State Sciences **58**, Springer, 1985.
- [88] D.P. O’Leary, Historical comment on [65], <http://nvlpubs.nist.gov/nistpubs/sp958-lide/077-080.pdf>
- [89] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, C. W. Clark, editors: *NIST Handbook of Mathematical Functions*, NIST (National Institute of Standards and Technology) and Cambridge University Press, 2010. Also digital library of mathematical functions <http://dlmf.nist.gov/>
- [90] S. Paszkowski, *Numerical Applications of Chebyshev Polynomials and Series* (Polish), Państwowe Wydawnictwo Naukowe, Warsaw, 1975. [Russian translation in: "Nauka", Fiz.-Mat. Lit., Moscow, 1983], *Polynômes et séries de Tchebichev*, Report ANO **140**, Univ. Lille1, Juillet 1984.
- [91] R. Piessens, M. Branders, The evaluation and application of some modified moments, *BIT* **13** (1973), 443-450.
- [92] R. Piessens, M.M. Chawla, N. Jayarajan, Gaussian quadrature formulas for the numerical calculation of integrals with logarithmic singularity, *J. Computational Physics* **21** 1976, 356-360.
- [93] M. Prevost, Approximation of weight function and approached Padé approximants, *Journal of Computational and Applied Mathematics* **32** (1990) 237-252.
- [94] R. A. Sack, A. F. Donovan, An Algorithm for Gaussian Quadrature Given Generalized Moments, Department of Mathematics, University of Salford, Salford, England, 1969.
- [95] R.A. Sack, A.F. Donovan, An algorithm for Gaussian quadrature given modified moments, *Numer. Math.* **18** (1971/72), 465-478.
- [96] E. Sadurní, Exact propagators on the lattice with applications to diffractive effects, *Journal of Physics A: Mathematical and Theoretical*, Volume **45** (2012), Number 46, article 465302, DOI <http://dx.doi.org/10.1088/1751-8113/45/46/465302>
- [97] R. Sasaki, M. Znojil, One-dimensional Schrödinger equation with non-analytic potential $V(x) = -g^2 \exp(-|x|)$ and its exact Bessel-function solvability, <http://arxiv.org/abs/1605.07310>.
- [98] B. Shizgal, *Spectral Methods in Chemistry and Physics. Applications to Kinetic Theory and Quantum Mechanics*, Springer 2015.
- [99] A. Sidi, Numerical Quadrature and Nonlinear Sequence Transformations; Unified Rules for Efficient Computation of Integrals With Algebraic and Logarithmic Endpoint Singularities, *Math. Comp.*, **35**, N. 151, July 1980, pp. 851-874.
- [100] G. Świdorski, Periodic perturbations of unbounded Jacobi matrices II: Formulas for density, *Journal of Approximation Theory* **216** (2017) 67-85.
- [101] G. Szegő, *Orthogonal Polynomials*, Amer. Math. Soc., 4th edition, 1975.
- [102] W.J. Thron, H. Waadeland, Accelerating convergence of limit periodic continued fractions $K(a_n/1)$, *Numer. Math.* **34** (1980), no. 2, 155-170.
- [103] Walter Van Assche, *Asymptotics for Orthogonal Polynomials*, Springer Lecture Notes in Mathematics **1265**, Springer-Verlag 1987.
- [104] W. Van Assche, Asymptotics for orthogonal polynomials and three-term recurrences, p. 435-462. in P. Nevai, editor: *Orthogonal Polynomials: Theory and Practice*, Proceedings of the NATO Advanced Study Institute on Orthogonal Polynomials and Their Applications Columbus, Ohio, U.S.A. May 22 - June 3, 1989, NATO ASI Series C: Mathematical and Physical Sciences Vol. **294**, Kluwer, 1990.
- [105] W. Van Assche, Christoffel functions and Turán determinants on several intervals. Proceedings of the Seventh Spanish Symposium on Orthogonal Polynomials and Applications (VII SPOA) (Granada, 1991). *J. Comput. Appl. Math.* **48** (1993), no. 1-2, 207-223.
- [106] W. Van Assche, E. Coussement, Some classical multiple orthogonal polynomials, *Journal of Computational and Applied Mathematics* **127** (2001) 317-347
- [107] H. S. Wall, *Analytic Theory of Continued Fractions*, New York, Van Nostrand, 1948 (= Chelsea Publishing, 1973).
- [108] A. Weisse, G. Wellein, A. Alvermann, and H. Fehske, The kernel polynomial method, *Rev. Modern Phys.*, **78** (2006) 275-306.
- [109] J.H. Wilkinson, *The algebraic eigenvalue problem*, Clarendon Press, Oxford 1965.
- [110] J. Wimp, *Computation with Recurrence Relations*, Pitman, 1984.
- [111] R. Wong, J. F. Lin, Asymptotic Expansions of Fourier Transforms of Functions with Logarithmic Singularities, *J. Math. An. Appl.* **64**, 173-180 (1978)
- [112] P. Wynn, Converging Factors for Continued Fractions, *Numerische Mathematik* **1**, 272-307 (1959)
- [113] P. Wynn, Note on a Converging Factor for a Certain Continued Fraction, *Numerische Mathematik* **5**, 332-352 (1963)
- [114] A. Zygmund, *Trigonometric Series*, Vol. I & II, Third edition, Cambridge University Press 2002.