

A higher dimensional homotopy sequence

M. Grandis E.M. Vitale

Abstract. We associate to a continuous map between pointed spaces a long 2-exact sequence of homotopy pointed groupoids. The usual homotopy sequence of a map follows from this 2-exact sequence taking, for each groupoid, the set of connected components. We also study a condition of strong 2-exactness for a sequence of cat-groups and pointed groupoids.

MSC 2000 : 18G55, 20L05, 55Q05.

Introduction

Let us start with an old story. The simplest homotopy invariant of a pointed topological space Y is the pointed set $\pi_0(Y)$ of its path-connected components. Using the loop functor Ω , we have a family of pointed sets $\pi_n(Y) = \pi_0(\Omega^n Y)$ which are (abelian) groups for $n \geq 1$ ($n \geq 2$). If $f: X \rightarrow Y$ is a continuous map preserving the base point, these homotopy invariants fit into a long exact sequence of groups and pointed sets

$$\dots \rightarrow \pi_n(Kf) \rightarrow \pi_n(X) \rightarrow \pi_n(Y) \rightarrow \dots$$

$$\dots \rightarrow \pi_1(Kf) \rightarrow \pi_1(X) \rightarrow \pi_1(Y) \rightarrow \pi_0(Kf) \rightarrow \pi_0(X) \rightarrow \pi_0(Y)$$

where $Kf \rightarrow X$ is the homotopy kernel of f .

The aim of this short note is to show that this sequence is a kind of “projection” on the category of pointed sets of an exact sequence of higher dimensional homotopy invariants. In fact the first two invariants of Y , $\pi_0(Y)$ and $\pi_1(Y)$, can be interpreted, respectively, as the set of connected components and the group of automorphisms at the base point of the fundamental groupoid $\Pi_1(Y)$. Using once again the loop functor, one obtains a family of pointed groupoids $\Pi_{n+1}(Y) = \Pi_1(\Omega^n Y)$. We will show that these homotopy invariants fit into a sequence of pointed groupoids and pointed functors

$$\dots \rightarrow \Pi_n(Kf) \rightarrow \Pi_n(X) \rightarrow \Pi_n(Y) \rightarrow \dots$$

$$\dots \rightarrow \Pi_2(Kf) \rightarrow \Pi_2(X) \rightarrow \Pi_2(Y) \rightarrow \Pi_1(Kf) \rightarrow \Pi_1(X) \rightarrow \Pi_1(Y)$$

which is “2-exact”, i.e. exact in a suitable categorical sense (see Definition 1). Taking, for each pointed groupoid of this sequence, the pointed set of its

connected components, one comes back to the classical exact sequence of the map f .

The second part of this paper is devoted to study a condition of strong 2-exactness which can be stated when a cat-group acts on a pointed groupoid. It is well-known that the classical homotopy sequence has a strong exactness at the transition point between groups and pointed sets

$$\pi_1(Y) \rightarrow \pi_0(Kf) \rightarrow \pi_0(X)$$

which is stated in terms of the action of the group $\pi_1(Y)$ on the pointed set $\pi_0(Kf)$. We show that the higher dimensional homotopy sequence satisfies a similar condition. We state our condition using a suitable 2-dimensional colimit, which is nothing but the cokernel (in the sense of bilimits) when the sequence is a sequence of symmetric cat-groups. This provide also a new interpretation of the strong exactness for a sequence of groups and pointed sets.

A warning: the composite of two arrows $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ in a category is denoted by $f \cdot g$.

1 Preliminaries

In this section we recall all the ingredients we need for the sequence of groupoids: the kernel of a morphism of pointed groupoids, the definition of 2-exactness, the homotopy equivariance of the fundamental groupoid of a space.

A pointed groupoid $\mathbb{G} = (\mathbb{G}, 0)$ is a groupoid \mathbb{G} (that is a category in which each arrow is an isomorphism) together with a chosen object 0 . A morphism of pointed groupoids $F: \mathbb{G} \rightarrow \mathbb{H}$ is a functor with a specified arrow $f_0: 0 \rightarrow F(0)$ in \mathbb{H} (a morphism $F: \mathbb{G} \rightarrow \mathbb{H}$ is strict if the arrow f_0 is the identity); a pointed natural transformation $\varphi: F \Rightarrow G: \mathbb{G} \rightarrow \mathbb{H}$ between morphisms of pointed groupoids is a natural transformation (necessarily a natural isomorphism) such that $f_0 \cdot \varphi_0 = g_0$. In this way we obtain a 2-category Gpd_* .

Given a morphism $F: \mathbb{G} \rightarrow \mathbb{H}$ in Gpd_* , its (homotopy) kernel $kF: KF \rightarrow \mathbb{G}$ can be described in the following way :

- an object of KF is a pair (X, x) with X an object of \mathbb{G} and $x: 0 \rightarrow F(X)$ an arrow in \mathbb{H} ;
- an arrow $f: (X, x) \rightarrow (X', x')$ of KF is an arrow $f: X \rightarrow X'$ in \mathbb{G} such that $x \cdot F(f) = x'$;
- the base object of KF is $(0, f_0)$;
- the functor kF sends $f: (X, x) \rightarrow (X', x')$ to $f: X \rightarrow X'$; it is a strict morphism.

There is a pointed natural transformation $\kappa F: 0 \Rightarrow kF \cdot F$ (where 0 is the constant morphism which sends each arrow to the identity of the base object of

\mathbb{H}) given, at the point (X, x) , by $x: 0 \rightarrow F(X)$.

$$\begin{array}{ccc}
 & \mathbb{G} & \\
 kF \nearrow & & \searrow F \\
 KF & \xrightarrow{\quad 0 \quad} & \mathbb{H} \\
 & \kappa F \uparrow & \\
 & &
 \end{array}
 \qquad
 \begin{array}{ccc}
 & \mathbb{G} & \\
 G \nearrow & & \searrow F \\
 \mathbb{K} & \xrightarrow{\quad 0 \quad} & \mathbb{H} \\
 & \varphi \uparrow & \\
 & &
 \end{array}$$

The triple $(KF, kF, \kappa F)$ has the following universal property : given any other triple (\mathbb{K}, G, φ) in Gpd_* as in the previous diagram, there is a unique morphism $G': \mathbb{K} \rightarrow KF$ such that $G' \cdot kF = G$ and $G' \cdot \kappa F = \varphi$. The functor G' is defined by $G'(g: Y \rightarrow Y') = G(g): (G(Y), \varphi_Y) \rightarrow (G(Y'), \varphi_{Y'})$. The kernel $(KF, kF, \kappa F)$ has also a “biuniversal” property, studied in [11, 16], which characterizes it up to equivalence.

A suitable notion of exactness in Gpd_* , introduced in [11, 16] to study some examples coming from ring theory, is given in the following definition, related to a notion of “homotopical exactness” studied in [9].

Definition 1 *A triple (G, φ, F) in Gpd_**

$$\begin{array}{ccc}
 & \mathbb{G} & \\
 G \nearrow & & \searrow F \\
 \mathbb{K} & \xrightarrow{\quad 0 \quad} & \mathbb{H} \\
 & \varphi \uparrow & \\
 & &
 \end{array}$$

is 2-exact if the comparison morphism $G': \mathbb{K} \rightarrow KF$ is full and essentially surjective on objects.

If $\mathbb{G} = (\mathbb{G}, 0)$ is a pointed groupoid, we write $\pi_0(\mathbb{G})$ for the pointed set of isomorphism classes of objects, and $\pi_1(\mathbb{G})$ for the group of automorphisms $\mathbb{G}(0, 0)$. If \mathbb{G} is a (braided) categorical group, $\pi_0\mathbb{G}$ is a (abelian) group and $\pi_1(\mathbb{G})$ is abelian (see [5, 10] for the notion of (braided) categorical group). Both π_0 and π_1 give rise to functors on the underlying category of Gpd_*

$$\pi_0: Gpd_* \rightarrow Set_* \qquad \pi_1: Gpd_* \rightarrow Groups$$

which are homotopy invariants, in the sense that if there is a 2-cell $\varphi: F \Rightarrow G: \mathbb{G} \rightarrow \mathbb{H}$ in Gpd_* , then $\pi_0(F) = \pi_0(G)$ and $\pi_1(F) = \pi_1(G)$. In particular, if F is an equivalence in Gpd_* , then $\pi_0(F)$ and $\pi_1(F)$ are isomorphisms. Finally, let us observe that if a triple (G, φ, F) as in Definition 1 is 2-exact, then

$$\pi_0(\mathbb{K}) \xrightarrow{\pi_0(G)} \pi_0(\mathbb{G}) \xrightarrow{\pi_0(F)} \pi_0(\mathbb{H}) \qquad \pi_1(\mathbb{K}) \xrightarrow{\pi_1(G)} \pi_1(\mathbb{G}) \xrightarrow{\pi_1(F)} \pi_1(\mathbb{H})$$

are exact sequences of pointed sets and groups.

Consider now a pointed topological space Y and its fundamental groupoid $\Pi_1(Y)$, i.e. the pointed groupoid having points of Y as objects and homotopy

rel end-points classes of paths as arrows (we use the additive notation for the concatenation of paths). This construction gives rise to a functor

$$\Pi_1: Top_* \rightarrow Gpd_*$$

between the category of pointed topological spaces and the category of pointed groupoids. Recall, from [3], the following lemma.

Lemma 2 *The functor Π_1 is homotopy equivariant. In particular, if $f: X \rightarrow Y$ is a homotopy equivalence in Top_* , then $\Pi_1(f)$ is an equivalence in Gpd_* .*

In fact, $\Pi_1: Top_* \rightarrow Gpd_*$ is a 2-functor, when we take as 2-cells in Top_* homotopy classes of homotopies. Moreover, for any f in Top_* , $\Pi_1(f)$ is a strict morphism in Gpd_* .

2 The exact sequence

Consider a map in Top_* together with its homotopy kernel

$$Kf \xrightarrow{kf} X \xrightarrow{f} Y$$

Recall that Kf is the subspace of the product space $X \times Y^I$ given by the pairs $(x, \eta: * \rightarrow f(x))$, with x a point of X , η a path in Y and $*$ the base point.

Proposition 3 *The sequence*

$$\Pi_1(Kf) \xrightarrow{\Pi_1(kf)} \Pi_1(X) \xrightarrow{\Pi_1(f)} \Pi_1(Y),$$

with the pointed natural transformation

$$\varphi: 0 \Rightarrow \Pi_1(kf) \cdot \Pi_1(f) \quad \varphi_{(x,\eta)} = [\eta]: * \rightarrow f(x),$$

is 2-exact.

Proof: Consider the following commutative diagram in Gpd_* , where f' is the comparison morphism

$$\begin{array}{ccc} \Pi_1(Kf) & \xrightarrow{\Pi_1(kf)} & \Pi_1(X) \\ & \searrow f' & \uparrow \\ & & K\Pi_1(f) \end{array}$$

Following the general description given in the first section, we obtain the following explicit description for $\Pi_1(Kf)$, $K\Pi_1(f)$ and f' :

- an object of $\Pi_1(Kf)$ is a pair $(x, \eta: * \rightarrow f(x))$ in $X \times Y^I$;

- an arrow $[h, H]: (x, \eta) \rightarrow (x', \eta')$ in $\Pi_1(Kf)$ is a class of pairs with $h: x \rightarrow x'$ a path in X and $H: I \rightarrow Y^I$ such that $H(0) = \eta, H(1) = \eta'$ and, for all t in I , $H(t): * \rightarrow f(h(t))$ a path in Y ;
- an object of $K\Pi_1(f)$ is a pair $(x, [\eta]: * \rightarrow f(x))$, where $[\eta]$ is a map in $\Pi_1(Y)$, i.e. a class of paths $\eta: * \rightarrow f(x)$ in Y ;
- an arrow $[h]: (x, [\eta]) \rightarrow (x', [\eta'])$ in $K\Pi_1(f)$ is a class of paths $h: x \rightarrow x'$ in X such that the following diagram in $\Pi_1(Y)$ commutes

$$\begin{array}{ccc}
 * & \xrightarrow{[\eta]} & f(x) \\
 & \searrow^{[\eta']} & \downarrow [h \cdot f] \\
 & & f(x')
 \end{array}$$

- the functor f' sends $[h, H]: (x, \eta) \rightarrow (x', \eta')$ to $[h]: (x, [\eta]) \rightarrow (x', [\eta'])$.

Clearly, f' is (essentially) surjective on objects. Moreover, consider an arrow $[h]: f'(x, \eta) \rightarrow f'(x', \eta')$ in $K\Pi_1(f)$. The commutativity of the previous diagram gives us a continuous map $L: I \times I \rightarrow Y$ such that $L(0, t) = (\eta + h \cdot f)(t), L(1, t) = \eta'(t), L(s, 0) = *, L(s, 1) = f(h(s))$ for all s, t in I . By a well-known transformation (studied abstractly in [8], under the name of “lens conversion”) one can derive from L a map $N: I \times I \rightarrow Y$ such that $N(0, t) = \eta(t), N(1, t) = \eta'(t), N(s, 0) = *, N(s, 1) = f(h(s))$ for all s, t in I . Finally, put $H: I \rightarrow Y^I$ $H(s) = N(s, -)$. In this way, we have an arrow $[h, H]: (x, \eta) \rightarrow (x', \eta')$ in $\Pi_1(Kf)$ such that $f'([h, H]) = [h]$, that is f' is full. \square

In the next corollary, we write $\Omega: Top_* \rightarrow Top_*$ for the loop-space endofunctor and $\Pi_{n+1}(Y)$ for the pointed groupoid $\Pi_1(\Omega^n Y)$.

Corollary 4 *Let $f: X \rightarrow Y$ be a map in Top_* : there is a long 2-exact sequence of pointed groupoids*

$$\begin{aligned}
 & \dots \rightarrow \Pi_n(Kf) \rightarrow \Pi_n(X) \rightarrow \Pi_n(Y) \rightarrow \dots \\
 & \dots \rightarrow \Pi_2(Kf) \rightarrow \Pi_2(X) \rightarrow \Pi_2(Y) \rightarrow \Pi_1(Kf) \rightarrow \Pi_1(X) \rightarrow \Pi_1(Y)
 \end{aligned}$$

Proof: It is enough to recall that the dual Puppe sequence [15]

$$\begin{aligned}
 & \dots \rightarrow \Omega^n(Kf) \rightarrow \Omega^n(X) \rightarrow \Omega^n(Y) \rightarrow \dots \\
 & \dots \rightarrow \Omega(Kf) \rightarrow \Omega(X) \rightarrow \Omega(Y) \rightarrow Kf \rightarrow X \rightarrow Y
 \end{aligned}$$

is homotopy equivalent to the sequence of iterated homotopy kernels

$$\dots \longrightarrow K(k(Kf)) \xrightarrow{k(k(Kf))} K(kf) \xrightarrow{k(kf)} Kf \xrightarrow{kf} X \xrightarrow{f} Y$$

(as proved in a general, abstract setting in [7]). Since to be full and essentially surjective is stable under composition with equivalences, we can apply Proposition 3 to each point of the kernel sequence and, by Lemma 2, we obtain the required long 2-exact sequence. \square

Remarks :

1) Clearly, $\pi_0(\Pi_1(Y)) = \pi_0(Y)$ and $\pi_1(\Pi_1(Y)) = \pi_1(Y)$; more generally, $\pi_0(\Pi_{n+1}(Y)) = \pi_n(Y)$ and $\pi_1(\Pi_{n+1}(Y)) = \pi_{n+1}(Y)$. As a consequence, applying the functor $\pi_0: Gpd_* \rightarrow Set_*$ to the 2-exact sequence of Corollary 4, we obtain the usual homotopy exact sequence

$$\dots \rightarrow \pi_1(Kf) \rightarrow \pi_1(X) \rightarrow \pi_1(Y) \rightarrow \pi_0(Kf) \rightarrow \pi_0(X) \rightarrow \pi_0(Y).$$

Applying the functor $\pi_1: Gpd_* \rightarrow Groups$, we obtain the same sequence, but we miss the three terms of degree zero.

2) In [6], A. Garzon, J. Miranda and A. del Río show that the groupoid $\Pi_n(Y)$ is a cat-group for $n \geq 2$, a braided cat-group for $n \geq 3$ and a symmetric cat-group for $n \geq 4$. Moreover, if $f: X \rightarrow Y$ is in Top_* and $n \geq 2$, $\Pi_n(f)$ is a monoidal functor (compatible with the braiding if $n \geq 3$). Since the definition of 2-exactness remains unchanged passing from pointed groupoids to (eventually braided or symmetric) cat-groups, the 2-exact sequence of Corollary 4 is in fact a 2-exact sequence of cat-groups for $n \geq 2$.

3 Strong exactness

Let A and B be two groups. To give a homomorphism $f: A \rightarrow B$ is equivalent to giving an action of the group A on the underlying set B which also satisfies a supplementary condition (iii), namely a mapping $+: A \times B \rightarrow B$ such that

- (i) $0_A + b = b$
- (ii) $(a_1 + a_2) + b = a_1 + (a_2 + b)$
- (iii) $a + (b_1 + b_2) = (a + b_1) + b_2$

for all $a, a_1, a_2 \in A$ and for all $b, b_1, b_2 \in B$. Indeed, given the action $+: A \times B \rightarrow B$, we get $f: A \rightarrow B$ by $f(a) = a + 0_B$. Conversely, given $f: A \rightarrow B$, we put $a + b = f(a) + b$. Moreover, given a morphism $g: B \rightarrow C$ in $Groups$, the following conditions are equivalent:

- (1) the composite $f \cdot g$ is equal to the zero morphism;
- (2) the following diagram, where p_B is the projection, commutes

$$\begin{array}{ccc} A \times B & \xrightarrow{+} & B \\ p_B \downarrow & & \downarrow g \\ B & \xrightarrow{g} & C \end{array}$$

Finally, the following conditions are equivalent:

- (I) if $g(b) = 0_C$, then there is $a \in A$ such that $b = f(a)$;

(II) if $g(b_1) = g(b_2)$, then there is $a \in A$ such that $b_1 = a + b_2$.

Assume now that A is a group and B and C are just pointed sets. Still, given a map $+: A \times B \rightarrow B$ such that (i) and (ii) hold, we obtain a morphism $f: A \rightarrow B$ in Set_* by $f(a) = a + 0_B$ (the opposite construction does not make sense). But now condition (2) is stronger than condition (1) and condition (II) is stronger than condition (I). The strong conditions can be expressed by means of the pointed orbit set B/A , a sort of ‘‘cokernel’’ of the action $+: A \times B \rightarrow B$. In fact, the projection $P_+: B \rightarrow B/A$ is the coequalizer of $+$ and p_B (within pointed sets). Now, a mapping $g: B \rightarrow C$ satisfies (2) iff it factors through P_+ ; it also satisfies (II) iff the comparison mapping $B/A \rightarrow C$ is surjective.

The interest of the *strong exactness* condition (II) comes from the homotopy sequence: if $f: X \rightarrow Y$ is in Top_* , there is a well-known action of the H -space ΩY on Kf

$$+: \Omega Y \times Kf \rightarrow Kf; \quad \omega + (x, \eta) = (x, \omega + \eta)$$

which induces a map $+: \pi_0(\Omega Y) \times \pi_0(Kf) \rightarrow \pi_0(Kf)$ such that conditions (i), (ii), (2) and (II) hold (see [2] for a detailed discussion).

The rest of this section is devoted to study the 2-dimensional analogue of strong exactness. Let \mathbb{A} be a cat-group and \mathbb{B} a pointed groupoid and consider a functor $\mu: \mathbb{A} \times \mathbb{B} \rightarrow \mathbb{B}$ together with two natural isomorphisms $m_B^0: B \rightarrow \mu(0, B)$ and $m_{A_1, A_2, B}: \mu(A_1 \otimes A_2, B) \rightarrow \mu(A_1, \mu(A_2, B))$, coherent with respect to the cat-group structure of \mathbb{A} . (This is equivalent to giving a monoidal functor from \mathbb{A} to the monoidal category of endofunctors of \mathbb{B} .) Starting from (μ, m^0, m) : $\mathbb{A} \times \mathbb{B} \rightarrow \mathbb{B}$, we can construct a new pointed groupoid $Cok\mu$ in the following way:

- the objects of $Cok\mu$ are those of \mathbb{B} ;
- a pre-morphism $(A, f): B_1 \rightarrow B_2$ in $Cok\mu$ is a pair with A in \mathbb{A} and $f: B_1 \rightarrow \mu(A, B_2)$ in \mathbb{B} ;
- a morphism $[A, f]: B_1 \rightarrow B_2$ is an equivalence class of pre-morphisms: two pre-morphisms $(A, f), (A', f'): B_1 \rightarrow B_2$ are equivalent if there is $\alpha: A \rightarrow A'$ such that $f \cdot \mu(\alpha, B_2) = f'$;
- the base point of $Cok\mu$ is that of \mathbb{B} .

There is a morphism of pointed groupoids $P_\mu: \mathbb{B} \rightarrow Cok\mu$ which sends $g: B_1 \rightarrow B_2$ to $[0, g \cdot m_{B_2}^0]: B_1 \rightarrow B_2$. There is also a natural transformation

$$\begin{array}{ccc} \mathbb{A} \times \mathbb{B} & \xrightarrow{\mu} & \mathbb{B} \\ p_{\mathbb{B}} \downarrow & \pi_\mu \Rightarrow & \downarrow P_\mu \\ \mathbb{B} & \xrightarrow{P_\mu} & Cok\mu \end{array}$$

given by $\pi_\mu(A, B) = [A^*, m_B^0 \cdot m_{A^*, A, B}: B \rightarrow \mu(0, B) \simeq \mu(A^* \otimes A, B) \rightarrow \mu(A^*, \mu(A, B))]: B \rightarrow \mu(A, B)$ (where A^* is a dual of A). Moreover, the following

diagrams in $Cok\mu$ commute

$$\begin{array}{ccc}
& B & \xrightarrow{\pi_\mu(A_1 \otimes A_2, B)} \mu(A_1 \otimes A_2, B) \\
\pi_\mu(0, B) \swarrow & \downarrow \pi_\mu(A_2, B) & \downarrow P_\mu(m_{A_1, A_2, B}) \\
& \mu(0, B) & \xrightarrow{\pi_\mu(A_1, \mu(A_2, B))} \mu(A_1, \mu(A_2, B)) \\
& \uparrow P_\mu(m_B^0) & \\
& B &
\end{array}$$

The previous construction is universal in the following sense ($Cok\mu$ is the iso-coserter of μ and $p_{\mathbb{B}}$): given

$$\begin{array}{ccc}
\mathbb{A} \times \mathbb{B} & \xrightarrow{\mu} & \mathbb{B} \\
p_{\mathbb{B}} \downarrow & \beta \Rightarrow & \downarrow G \\
\mathbb{B} & \xrightarrow{G} & \mathbb{C}
\end{array}$$

with $G: \mathbb{B} \rightarrow \mathbb{C}$ in Gpd_* and β a natural transformation such that

$$\begin{array}{ccc}
G(B) & \xrightarrow{\beta_{A_1 \otimes A_2, B}} G(\mu(A_1 \otimes A_2, B)) \\
\beta_{0, B} \swarrow & \downarrow \beta_{A_2, B} & \downarrow G(m_{A_1, A_2, B}) \\
& G(\mu(0, B)) & \xrightarrow{\beta_{A_1, \mu(A_2, B)}} G(\mu(A_1, \mu(A_2, B))) \\
& \uparrow G(m_B^0) & \\
& G(B) &
\end{array}$$

commute, there is a unique $G': Cok\mu \rightarrow \mathbb{C}$ in Gpd_* such that $P_\mu \cdot G' = G$ and $\pi_\mu \cdot G' = \beta$.

Proof : One has to define $G': Cok\mu \rightarrow \mathbb{C}$ by

$$G': [A, f]: B_1 \rightarrow B_2 \mapsto G(f) \cdot \beta_{A, B_2}^{-1}: G(B_1) \rightarrow G(\mu(A, B_2)) \rightarrow G(B_2)$$

The uniqueness follows from the commutativity of the following diagram

$$\begin{array}{ccc}
B_1 & \xrightarrow{[A, f]} & B_2 \\
P_\mu(f) \searrow & & \swarrow \pi_\mu(A, B_2) \\
& \mu(A, B_2) &
\end{array}$$

□

This universal property characterizes $Cok\mu$ up to isomorphism. The triple $(Cok\mu, P_\mu, \pi_\mu)$ has also a “biuniversal” property (it is the bi-coequaliser of μ and $p_{\mathbb{B}}$) which characterizes it up to equivalence. It is similar to those of the kernel and of the cokernel (see [11, 16]).

Let us explain the notation $Cok\mu$. Starting from $(\mu, m^0, m): \mathbb{A} \times \mathbb{B} \rightarrow \mathbb{B}$, we get a morphism $F: \mathbb{A} \rightarrow \mathbb{B}$ in Gpd_* by $F(A) = \mu(A, 0)$ and $f_0 = m_0^0: 0 \rightarrow \mu(0, 0) = F(0)$. If it is the case that \mathbb{A} and \mathbb{B} are symmetric cat-groups and F is a monoidal functor compatible with the symmetry, then $Cok\mu$ is exactly the cokernel of F as described in [11, 16].

In order to state strong exactness for pointed groupoids, observe that, given

$$\begin{array}{ccc} \mathbb{A} \times \mathbb{B} & \xrightarrow{\mu} & \mathbb{B} \\ p_{\mathbb{B}} \downarrow & \beta \Rightarrow & \downarrow G \\ \mathbb{B} & \xrightarrow{G} & \mathbb{C} \end{array}$$

as before, we get a pointed natural transformation

$$\begin{array}{ccc} & \mathbb{B} & \\ F \nearrow & & \searrow G \\ \mathbb{A} & \xrightarrow{0} & \mathbb{C} \\ & \alpha \uparrow & \end{array}$$

by $\alpha_A = g_0 \cdot \beta_{A,0}: 0 \rightarrow G(O) \rightarrow G(\mu(A, 0)) = G(F(A))$. Recall now that, if (F, α, G) is a sequence of symmetric cat-groups, its 2-exactness can be equivalently stated by asking that the canonical comparison from the cokernel of F to G is full and faithful (Proposition 6.2 in [11]). With this fact in mind, we give the following definition.

Definition 5 *Consider*

$$\begin{array}{ccc} \mathbb{A} \times \mathbb{B} & \xrightarrow{\mu} & \mathbb{B} \\ p_{\mathbb{B}} \downarrow & \beta \Rightarrow & \downarrow G \\ \mathbb{B} & \xrightarrow{G} & \mathbb{C} \end{array}$$

as before. The sequence (μ, β, G) is strongly 2-exact if the comparison functor $G': Cok\mu \rightarrow \mathbb{C}$ is full and faithful.

Here is the expected link between strong 2-exactness (Definition 5) and 2-exactness (Definition 1). The proof is a direct calculation.

Proposition 6 *Consider*

$$\begin{array}{ccc} \mathbb{A} \times \mathbb{B} & \xrightarrow{\mu} & \mathbb{B} \\ p_{\mathbb{B}} \downarrow & \beta \Rightarrow & \downarrow G \\ \mathbb{B} & \xrightarrow{G} & \mathbb{C} \end{array} \quad \text{and} \quad \begin{array}{ccc} & \mathbb{B} & \\ F \nearrow & & \searrow G \\ \mathbb{A} & \xrightarrow{0} & \mathbb{C} \\ & \alpha \uparrow & \end{array}$$

as before. Consider also the factorization F' of F through the kernel of G and the factorization G' of G through $\text{Cok}\mu$

$$F' : \mathbb{A} \rightarrow KG \quad G' : \text{Cok}\mu \rightarrow \mathbb{C}.$$

- a) If G' is faithful, then F' is full;
b) If G' is full, then F' is essentially surjective.

The interested reader can verify that, if $\mathbb{A}, \mathbb{B}, \mathbb{C}, F, G$ and α belong to the 2-category of cat-groups and if β is compatible with the cat-group structure of \mathbb{B} and \mathbb{C} , then both the implications of Proposition 6 can be reversed.

Let us write down explicitly the condition of fullness for $G' : \text{Cok}\mu \rightarrow \mathbb{C}$: given two objects B_1 and B_2 in \mathbb{B} and an arrow $c : G(B_1) \rightarrow G(B_2)$ in \mathbb{C} , there is A in \mathbb{A} and $f : B_1 \rightarrow \mu(A, B_2)$ in \mathbb{B} such that $G'[A, f] = c$.

The analogy between fullness of G' and strong exactness for a sequence of groups and pointed sets, i.e. condition (II), is now clear. This analogy is made more precise in the next remark.

Remark :

1) Assume that the categories \mathbb{A}, \mathbb{B} and \mathbb{C} of Proposition 6 are discrete (so that \mathbb{A} is a group, and \mathbb{B} and \mathbb{C} are pointed sets, but $\text{Cok}\mu$ is not discrete). Then the fullness of G' is exactly condition (II) (moreover, the faithfulness of G' implies that F is injective).

2) Assume that the categories \mathbb{A}, \mathbb{B} and \mathbb{C} of Proposition 6 have a unique object (so that \mathbb{A} is an abelian group and \mathbb{B} and \mathbb{C} are groups). Then the faithfulness of G' is exactly condition (II) (moreover, the fullness of G' is equivalent to the surjectivity of G).

Finally, we come back to the higher dimensional homotopy sequence. Let $f : X \rightarrow Y$ be an arrow in Top_* and consider the action $\Omega Y \times Kf \rightarrow Kf$ as at the beginning of this section. It induces an action $\mu : \Pi_1(\Omega Y) \times \Pi_1(Kf) \rightarrow \Pi_1(Kf)$ of the cat-group $\Pi_2(Y) = \Pi_1(\Omega Y)$ on the pointed groupoid $\Pi_1(Kf)$. Moreover, the diagram

$$\begin{array}{ccc} \Pi_2(Y) \times \Pi_1(Kf) & \xrightarrow{\mu} & \Pi_1(Kf) \\ p_{\Pi_1(Kf)} \downarrow & & \downarrow \Pi_1(kf) \\ \Pi_1(Kf) & \xrightarrow{\Pi_1(kf)} & \Pi_1(X) \end{array}$$

is strictly commutative.

Proposition 7 *Let*

$$\begin{array}{ccc} \Pi_2(Y) \times \Pi_1(Kf) & \xrightarrow{\mu} & \Pi_1(Kf) \\ p_{\Pi_1(Kf)} \downarrow & = & \downarrow \Pi_1(kf) \\ \Pi_1(Kf) & \xrightarrow{\Pi_1(kf)} & \Pi_1(X) \end{array}$$

be as before. The sequence $(\mu, =, \Pi_1(kf))$ is strongly 2-exact.

We leave the proof as an exercise for the reader.

References

- [1] F. BORCEUX: *Handbook of categorical algebra*, Cambridge University Press (1994).
- [2] R. BROWN: *Fibrations of groupoids*, Journal of Algebra 15 (1970) 103-132.
- [3] R. BROWN: *Topology: A geometric account of general topology, homotopy types and the fundamental groupoid*, Ellis Horwood Ltd. (1988).
- [4] R. BROWN: *Groupoids and crossed objects in algebraic topology*, Homology, Homotopy and Applications 1 (1999) 1-78 (available at <http://www.rmi.acnet.ge/hha/>).
- [5] P. CARRASCO, A.M. CEGARRA: *(Braided) tensor structures on homotopy groupoids and nerves of (braided) categorical groups*, Communications in Algebra 24 (1996) 3995-4058.
- [6] A.R. GARZON, J.G. MIRANDA, A. DEL RÍO: *Tensor structures on homotopy groupoids of topological spaces*, Intern. Math. Journal 2 (2002) 407-431.
- [7] M. GRANDIS: *Homotopical algebra in homotopical categories*, Appl. Categ. Struct. 2 (1994) 351-406.
- [8] M. GRANDIS: *Categorically algebraic foundations for homotopical algebra*, Appl. Categ. Struct. 5 (1997) 363-413.
- [9] M. GRANDIS: *A note on exactness and stability in homotopical algebra*, Theory and Applications of Categories 9 (2001) 17-42 (available at <http://www.tac.mta.ca/tac/>).
- [10] A. JOYAL, R. STREET: *Braided tensor categories*, Adv. Math. 102 (1993) 20-78.
- [11] S. KASANGIAN, E.M. VITALE: *Factorization systems for symmetric cat-groups*, Theory and Applications of Categories 7 (2000) 47-70 (available at <http://www.tac.mta.ca/tac/>).
- [12] R.W. KIEBOOM, E.M. VITALE: *On the exact sequence associated to a fibration of 2-groupoids*, preprint (2000).
- [13] M. MATHER: *Pull-backs in homotopy theory*, Can. J. Math. 28 (1976) 225-263.

- [14] R.A. PICCININI: *Lectures on homotopy theory*, North-Holland (1992).
- [15] D. Puppe: *Homotopiemengen und ihre induzierten Abbildungen, I*, Math. Z. 69 (1958) 299-344, 395-417.
- [16] E.M. VITALE: *A Picard-Brauer exact sequence of categorical groups*, J. Pure Appl. Algebra (to appear).
- [17] G.W. WHITEHEAD: *Elements of homotopy theory*, Springer Verlag (1978).

Marco Grandis
Dipartimento di Matematica, Università di Genova,
Via Dodecaneso 35, 16146 Genova, Italy
grandis@dima.unige.it

Enrico M. Vitale
Département de Mathématique, Université catholique de Louvain,
Chemin du Cyclotron 2, 1348 Louvain-la-Neuve, Belgium
vitale@math.ucl.ac.be