

$\mathfrak{sl}(N)$ –link homology ($N \geq 4$) using foams and the Kapustin–Li formula

MARCO MACKAAY

MARKO STOŠIĆ

PEDRO VAZ

We use foams to give a topological construction of a rational link homology categorifying the $\mathfrak{sl}(N)$ link invariant, for $N \geq 4$. To evaluate closed foams we use the Kapustin–Li formula adapted to foams by Khovanov and Rozansky [9]. We show that for any link our homology is isomorphic to the Khovanov–Rozansky [11] homology.

57M27; 57M25, 81R50, 18G60

1 Introduction

In [14], Murakami, Ohtsuki and Yamada (MOY) developed a graphical calculus for the $\mathfrak{sl}(N)$ link polynomial. In [7], Khovanov categorified the $\mathfrak{sl}(3)$ polynomial using singular cobordisms between webs called foams. Mackaay and Vaz [12] generalized Khovanov’s results to obtain the universal $\mathfrak{sl}(3)$ integral link homology, following an approach similar to the one adopted by Bar-Natan [1] for the original $\mathfrak{sl}(2)$ integral Khovanov homology. In [11] Khovanov and Rozansky defined a rational link homology which categorifies the $\mathfrak{sl}(N)$ link polynomial using the theory of matrix factorizations.

In this paper we use foams, as in [1; 7; 12], for an almost completely combinatorial topological construction of a rational link homology categorifying the $\mathfrak{sl}(N)$ link polynomial. Our theory is functorial under link cobordisms up to nonzero scalars. Khovanov had to modify considerably his original setting for the construction of $\mathfrak{sl}(2)$ link homology in order to produce his $\mathfrak{sl}(3)$ link homology. It required the introduction of singular cobordisms with a particular type of singularity. The jump from $\mathfrak{sl}(3)$ to $\mathfrak{sl}(N)$, for $N > 3$, requires the introduction of a new type of singularity. The latter is needed for proving invariance under the third Reidemeister move and is qualitatively different from the other singularities. The old singularities, which are the same as for $\mathfrak{sl}(3)$, are part of seams where three facets meet. The new singularities occur at points where such seams intersect. This is new, because for $\mathfrak{sl}(3)$ the seams are disjoint arcs. Furthermore the combinatorics involved in establishing certain identities gets much

harder for arbitrary N . The theory of symmetric polynomials, in particular Schur polynomials, is used to handle that problem.

Our aim was to find a combinatorial topological definition of Khovanov–Rozansky link homology. Such a definition is desirable for several reasons, the main one being that it might help to find a good way to compute the Khovanov–Rozansky link homology. Unfortunately the construction that we present in this paper is not completely combinatorial. The introduction of the new singularities makes it much harder to evaluate closed foams and we do not know how to do it combinatorially. Instead we use the Kapustin–Li formula [6], adapted by Khovanov and Rozansky [9]¹. A positive side-effect is that it allows us to show that for any link our homology is isomorphic to Khovanov and Rozansky’s.

Although we have not completely achieved our final goal, we believe that we have made good progress towards it. In Propositions 6.2 and 6.9 we derive a small set of relations on foams which we show to be sufficient to guarantee that our link homology is homotopy invariant under the Reidemeister moves. By deriving these relations from the Kapustin–Li formula we prove that these relations are consistent. However, in order to get a purely combinatorial construction we would have to show that they generate the kernel of the Kapustin–Li formula on closed foams. We conjecture that this holds true, but so far our attempts to prove it have failed. It would be very interesting to have a proof of this conjecture. So far we can only conclude that any other way of evaluating closed foams which satisfies the same relations as in Propositions 6.2 and 6.9 gives rise to a functorial link homology which categorifies the $\mathfrak{sl}(N)$ link polynomial. We conjecture that such a link homology is equivalent to the one presented in this paper and therefore to Khovanov and Rozansky’s.

In Section 2 we recall some basic facts about the $\mathfrak{sl}(N)$ –link polynomials. In Section 3 we recall some basic facts about Schur polynomials and the cohomology of partial flag varieties. In Section 4 we define prefoams and their grading. In Section 5 we explain the Kapustin–Li formula for evaluating closed prefoams and compute the spheres and the theta-foams. In Section 6 we derive a set of basic relations in the category \mathbf{Foam}_N , which is the quotient of the category of prefoams by the kernel of the Kapustin–Li evaluation. In Section 7 we show that our link homology complex is homotopy invariant under the Reidemeister moves. In Section 8 we show that our link homology complex extends to a link homology projective functor. In Section 9 we show that our link homology, obtained from our link homology complex using the tautological functor, categorifies the $\mathfrak{sl}(N)$ –link polynomial and that it is isomorphic to the Khovanov–Rozansky link homology.

¹We thank M Khovanov for suggesting that we try to use the Kapustin–Li formula.

2 Graphical calculus for the $\mathfrak{sl}(N)$ polynomial

In this section we recall some facts about the graphical calculus for $\mathfrak{sl}(N)$. The $\mathfrak{sl}(N)$ link polynomial is defined by the skein relation

$$q^N P_N(\text{positive}) - q^{-N} P_N(\text{negative}) = (q - q^{-1}) P_N(\text{flattened}),$$

and its value for the unknot, which we take to be equal to $[N] = (q^N - q^{-N}) / (q - q^{-1})$. Let D be a diagram of a link $L \in S^3$ with n_+ positive crossings and n_- negative crossings. Following an approach based on MOY’s state sum model [14] we can




Figure 1: Positive and negative crossings and their 0- and 1-flattening

compute $P_N(D)$ by flattening each crossing of D in two possible ways, as shown in Figure 1, where we also show our convention for positive and negative crossings. Each complete flattening of D is an example of a *web*: a trivalent graph with three types of edges: *simple*, *double* and *marked* edges. A vertex has at most one marked incident edge. Only the simple edges are equipped with an orientation, which fixes the orientations on the other edges according to the following conventions: given a vertex with two simple ingoing (outgoing) edges, then it has one double edge which is outgoing (ingoing). There are no vertices with one ingoing simple edge and one outgoing simple edge. If a vertex has only one simple ingoing (outgoing) edge, then it has one double edge which is ingoing (outgoing) and one marked edge which is outgoing (ingoing). Other vertices are not allowed (see Figure 2). In a web the orientations of all edges have to match up according to the rules above. Note that a complete flattening of D never has marked edges, but we will need the latter for webs that show up in the proof of invariance under the third Reidemeister move.




Figure 2: Vertices

Simple edges correspond to edges labelled 1, double edges to edges labelled 2 and marked simple edges to edges labelled 3 in [14], where edges carry labels from 1

to $N - 1$ and label j is associated to the j -th exterior power of the fundamental representation of $\mathfrak{sl}(N)$.

We call a planar trivalent graph generated by the vertices and edges defined above a *web*. Webs can contain closed plane loops (simple, double or marked). The *MOY web moves* in Figure 3 provide a recursive way of assigning to each web Γ that only contains simple and double edges a polynomial in $\mathbb{Z}[q, q^{-1}]$ with positive coefficients, which we call $P_N(\Gamma)$. There are more general web moves [13], which allow for the evaluation of arbitrary webs, but we do not need them here. Note that a complete flattening of a link diagram only contains simple and double edges.




Figure 3: MOY web moves

Consistency of the relations in Figure 3 is shown in [14].

Finally let us define the $\mathfrak{sl}(N)$ link polynomial. For any i let Γ_i denote a complete flattening of D . Then

$$P_N(D) = \sum_{\Gamma} q^{\alpha(\Gamma)} P_N(\Gamma),$$

where $\alpha(\Gamma)$ is determined by the coefficients of the resolutions in Figure 1 and the sum is over all possible flattenings Γ of D .

3 Schur polynomials and the cohomology of partial flag varieties

In this section we recall some basic facts about Schur polynomials and the cohomology of partial flag varieties which we need in the rest of this paper.

3.1 Schur polynomials

A nice basis for homogeneous symmetric polynomials is given by the Schur polynomials. If $\lambda = (\lambda_1, \dots, \lambda_k)$ is a partition such that $\lambda_1 \geq \dots \geq \lambda_k \geq 0$, then the Schur polynomial $\pi_\lambda(x_1, \dots, x_k)$ is given by the following expression:

$$(1) \quad \pi_\lambda(x_1, \dots, x_k) = \frac{|x_i^{\lambda_j+k-j}|}{\Delta},$$

where $\Delta = \prod_{i < j} (x_i - x_j)$, and by $|x_i^{\lambda_j+k-j}|$, we have denoted the determinant of the $k \times k$ matrix whose (i, j) entry is equal to $x_i^{\lambda_j+k-j}$. Note that the elementary symmetric polynomials are given by $\pi_{1,0,0,\dots,0}, \pi_{1,1,0,\dots,0}, \dots, \pi_{1,1,1,\dots,1}$. There are multiplication rules for the Schur polynomials which show that any $\pi_{\lambda_1, \lambda_2, \dots, \lambda_k}$ can be expressed in terms of the elementary symmetric polynomials.

If we do not specify the variables of the Schur polynomial π_λ , we will assume that these are exactly x_1, \dots, x_k , with k being the length of λ , ie

$$\pi_{\lambda_1, \dots, \lambda_k} := \pi_{\lambda_1, \dots, \lambda_k}(x_1, \dots, x_k).$$

In this paper we only use Schur polynomials of two and three variables. In the case of two variables, the Schur polynomials are indexed by pairs of nonnegative integers (i, j) , such that $i \geq j$, and (1) becomes

$$\pi_{i,j} = \sum_{\ell=j}^i x_1^\ell x_2^{i+j-\ell}.$$

Directly from *Pieri’s formula* (see Fulton [3] and Fulton and Harris [4]) we obtain the following multiplication rule for the Schur polynomials in two variables:

$$(2) \quad \pi_{i,j} \pi_{a,b} = \sum \pi_{x,y},$$

where the sum on the right-hand side is over all indices x and y such that $x + y = i + j + a + b$ and $a + i \geq x \geq \max(a + j, b + i)$. Note that this implies $\min(a + j, b + i) \geq y \geq b + j$. Also, we shall write $\pi_{x,y} \in \pi_{i,j} \pi_{a,b}$ if $\pi_{x,y}$ belongs to the sum on the right-hand side of (2). Hence, we have that $\pi_{x,x} \in \pi_{i,j} \pi_{a,b}$ if and only if $a + j = b + i = x$ and $\pi_{x+1,x} \in \pi_{i,j} \pi_{a,b}$ if and only if $a + j = x + 1, b + i = x$ or $a + j = x, b + i = x + 1$.

We shall need the following combinatorial result which expresses the Schur polynomial in three variables as a combination of Schur polynomials of two variables.

For $i \geq j \geq k \geq 0$, and the triple (a, b, c) of nonnegative integers, we define

$$(a, b, c) \sqsubset (i, j, k),$$

if $a + b + c = i + j + k$, $i \geq a \geq j$ and $j \geq b \geq k$. We note that this implies that $i \geq c \geq k$, and hence $\max\{a, b, c\} \leq i$.

Lemma 3.1

$$\pi_{i,j,k}(x_1, x_2, x_3) = \sum_{(a,b,c) \sqsubset (i,j,k)} \pi_{a,b}(x_1, x_2)x_3^c.$$

Proof From the definition of the Schur polynomial, we have

$$\pi_{i,j,k}(x_1, x_2, x_3) = \frac{(x_1x_2x_3)^k}{(x_1 - x_2)(x_1 - x_3)(x_2 - x_3)} \begin{vmatrix} x_1^{i-k+2} & x_1^{j-k+1} & 1 \\ x_2^{i-k+2} & x_2^{j-k+1} & 1 \\ x_3^{i-k+2} & x_3^{j-k+1} & 1 \end{vmatrix}.$$

After subtracting the last row from the first and the second one of the last determinant, we obtain

$$\pi_{i,j,k} = \frac{(x_1x_2x_3)^k}{(x_1 - x_2)(x_1 - x_3)(x_2 - x_3)} \begin{vmatrix} x_1^{i-k+2} - x_3^{i-k+2} & x_1^{j-k+1} - x_3^{j-k+1} & 0 \\ x_2^{i-k+2} - x_3^{i-k+2} & x_2^{j-k+1} - x_3^{j-k+1} & 0 \\ 0 & 0 & 1 \end{vmatrix},$$

and so

$$\pi_{i,j,k} = \frac{(x_1x_2x_3)^k}{x_1 - x_2} \begin{vmatrix} \sum_{m=0}^{i-k+1} x_1^m x_3^{i-k+1-m} & \sum_{n=0}^{j-k} x_1^n x_3^{j-k-n} \\ \sum_{m=0}^{i-k+1} x_2^m x_3^{i-k+1-m} & \sum_{n=0}^{j-k} x_2^n x_3^{j-k+n} \end{vmatrix}.$$

Finally, after expanding the last determinant we obtain

$$(3) \quad \pi_{i,j,k} = \frac{(x_1x_2x_3)^k}{x_1 - x_2} \sum_{m=0}^{i-k+1} \sum_{n=0}^{j-k} (x_1^m x_2^n - x_1^n x_2^m) x_3^{i+j-2k+1-m-n}.$$

We split the last double sum into two: the first one when m goes from 0 to $j - k$, denoted by S_1 , and the other one when m goes from $j - k + 1$ to $i - k + 1$, denoted by S_2 . To show that $S_1 = 0$, we split the double sum further into three parts: when $m < n$, $m = n$ and $m > n$. Obviously, each summand with $m = n$ is equal to 0, while the summands of the sum for $m < n$ are exactly the opposite of the summands of the sum for $m > n$. Thus, by replacing only S_2 instead of the double sum in (3) and after rescaling the indices $a = m + k - 1$, $b = n + k$, we get

$$\begin{aligned} \pi_{i,j,k} &= \frac{(x_1x_2x_3)^k}{x_1 - x_2} \sum_{m=j-k+1}^{i-k+1} \sum_{n=0}^{j-k} (x_1^m x_2^n - x_1^n x_2^m) x_3^{i+j-2k+1-m-n} \\ &= \sum_{a=j}^i \sum_{b=k}^j \pi_{a,b} x_3^{i+j+k-a-b} = \sum_{(a,b,c) \sqsubset (i,j,k)} \pi_{a,b} x_3^c. \quad \square \end{aligned}$$

Of course there is a multiplication rule for three-variable Schur polynomials which is compatible with (2) and the lemma above, but we do not want to discuss it here. For details see Fulton [3] and Fulton and Harris [4].

3.2 The cohomology of partial flag varieties

In this paper the rational cohomology rings of partial flag varieties play an essential role. The partial flag variety $\text{Fl}_{d_1, d_2, \dots, d_l}$ for $1 \leq d_1 < d_2 < \dots < d_l = N$ is defined by

$$\text{Fl}_{d_1, d_2, \dots, d_l} = \{V_{d_1} \subset V_{d_2} \subset \dots \subset V_{d_l} = \mathbb{C}^N \mid \dim(V_i) = i\}.$$

A special case is $\text{Fl}_{k, N}$, the Grassmannian variety of all k -planes in \mathbb{C}^N , also denoted $\mathcal{G}_{k, N}$. The dimension of the partial flag variety is given by

$$\dim \text{Fl}_{d_1, d_2, \dots, d_l} = N^2 - \sum_{i=1}^{l-1} (d_{i+1} - d_i)^2 - d_1^2.$$

The rational cohomology rings of the partial flag varieties are well known and we only recall those facts that we need in this paper. For more details and proofs see Fulton [3] and Hiller [5].

Lemma 3.2 $H^*(\mathcal{G}_{k, N})$ is isomorphic to the vector space generated by all $\pi_{i_1, i_2, \dots, i_k}$ modulo the relations

$$(4) \quad \pi_{N-k+1, 0, \dots, 0} = 0, \quad \pi_{N-k+2, 0, \dots, 0} = 0, \quad \dots, \quad \pi_{N, 0, \dots, 0} = 0,$$

where there are exactly $k - 1$ zeros in the multi-indices of the Schur polynomials.

A consequence of the multiplication rules for Schur polynomials is:

Corollary 3.3 The Schur polynomials $\pi_{i_1, i_2, \dots, i_k}$, for $N - k \geq i_1 \geq i_2 \geq \dots \geq i_k \geq 0$, form a basis of $H^*(\mathcal{G}_{k, N})$.

Thus, the dimension of $H^*(\mathcal{G}_{k, N})$ is $\binom{N}{k}$, and up to a degree shift, its quantum dimension (or graded dimension) is $\left[\begin{smallmatrix} N \\ k \end{smallmatrix} \right]$.

Another consequence of the multiplication rules is:

Corollary 3.4 The Schur polynomials $\pi_{1, 0, 0, \dots, 0}, \pi_{1, 1, 0, \dots, 0}, \dots, \pi_{1, 1, 1, \dots, 1}$ (the elementary symmetric polynomials) generate $H^*(\mathcal{G}_{k, N})$ as a ring.

Furthermore, we can introduce a nondegenerate trace form on $H^*(\mathcal{G}_{k,N})$ by giving its values on the basis elements

$$(5) \quad \epsilon(\pi_\lambda) = \begin{cases} (-1)^{\lfloor k/2 \rfloor}, & \lambda = (N-k, \dots, N-k) \\ 0, & \text{else.} \end{cases}$$

This makes $H^*(\mathcal{G}_{k,N})$ into a commutative Frobenius algebra. One can compute the basis dual to $\{\pi_\lambda\}$ in $H^*(\mathcal{G}_{k,N})$, with respect to ϵ . It is given by

$$(6) \quad \hat{\pi}_{\lambda_1, \dots, \lambda_k} = (-1)^{\lfloor k/2 \rfloor} \pi_{N-k-\lambda_k, \dots, N-k-\lambda_1}.$$

We can also express the cohomology rings of the partial flag varieties $\text{Fl}_{1,2,N}$ and $\text{Fl}_{2,3,N}$ in terms of Schur polynomials. Indeed, we have

$$\begin{aligned} H^*(\text{Fl}_{1,2,N}) &= \mathbb{Q}[x_1, x_2] / \langle \pi_{N-1,0}, \pi_{N,0} \rangle, \\ H^*(\text{Fl}_{2,3,N}) &= \mathbb{Q}[x_1 + x_2, x_1 x_2, x_3] / \langle \pi_{N-2,0,0}, \pi_{N-1,0,0}, \pi_{N,0,0} \rangle. \end{aligned}$$

The natural projection map $p_1: \text{Fl}_{1,2,N} \rightarrow \mathcal{G}_{2,N}$ induces

$$p_1^*: H^*(\mathcal{G}_{2,N}) \rightarrow H^*(\text{Fl}_{1,2,N}),$$

which is just the inclusion of the polynomial rings. Analogously, the natural projection $p_2: \text{Fl}_{2,3,N} \rightarrow \mathcal{G}_{3,N}$, induces

$$p_2^*: H^*(\mathcal{G}_{3,N}) \rightarrow H^*(\text{Fl}_{2,3,N}),$$

which is also given by the inclusion of the polynomial rings.

4 Prefoams

In this section we begin to define the foams we will work with. The philosophy behind these foams will be explained in Section 5. To categorify the $\mathfrak{sl}(N)$ link polynomial we need singular cobordisms with two types of singularities. The basic examples are given in Figure 4. These foams are composed of three types of facets: simple, double and triple facets. The double facets are coloured and the triple facets are marked to show the difference. Intersecting such a foam with a plane results in a web, as long as the plane avoids the singularities where six facets meet, such as on the right in Figure 4.

We adapt the definition of a world-sheet foam given in [15] to our setting.

Definition 4.1 Let s_γ be a finite closed oriented 4-valent graph, which may contain disjoint circles. We assume that all edges of s_γ are oriented. A cycle in s_γ is defined to be a circle or a closed sequence of edges which form a piece-wise linear circle. Let Σ be a compact orientable possibly disconnected surface, whose connected components are




Figure 4: Some elementary foams

white, coloured or marked, also denoted by simple, double or triple. Each component can have a boundary consisting of several disjoint circles and can have additional decorations which we discuss below. A closed *prefoam* u is the identification space Σ/s_γ obtained by gluing boundary circles of Σ to cycles in s_γ such that every edge and circle in s_γ is glued to exactly three boundary circles of Σ and such that for any point $p \in s_\gamma$,

- (1) if p is an interior point of an edge, then p has a neighborhood homeomorphic to the letter Y times an interval with exactly one of the facets being double, and at most one of them being triple. For an example see Figure 4;
- (2) if p is a vertex of s_γ , then it has a neighborhood as shown in Figure 4.

We call s_γ the *singular graph*, its edges and vertices *singular arcs* and *singular vertices*, and the connected components of $u - s_\gamma$ the *facets*.

Furthermore the facets can be decorated with dots. A simple facet can only have black dots (\bullet), a double facet can also have white dots (\circ), and a triple facet besides black and white dots can have double dots (\odot). Dots can move freely on a facet but are not allowed to cross singular arcs. See Figure 5 for an example of a prefoam.




Figure 5: A prefoam

Note that the cycles to which the boundaries of the simple and the triple facets are glued are always oriented, whereas the ones to which the boundaries of the double facets are

glued are not. Note also that there are two types of singular vertices. Given a singular vertex v , there are precisely two singular edges which meet at v and bound a triple facet: one oriented toward v , denoted e_1 , and one oriented away from v , denoted e_2 . If we use the “left hand rule”, then the cyclic ordering of the facets incident to e_1 and e_2 is either $(3, 2, 1)$ and $(3, 1, 2)$ respectively, or the other way around. We say that v is of type I in the first case and of type II in the second case. When we go around a triple facet we see that there have to be as many singular vertices of type I as there are of type II for the cyclic orderings of the facets to match up. This shows that for a closed prefoam the number of singular vertices of type I is equal to the number of singular vertices of type II.

We can intersect a prefoam u generically by a plane W in order to get a web, as long as the plane avoids the vertices of s_γ . The orientation of s_γ determines the orientation of the simple edges of the web according to the convention in Figure 6.




Figure 6: Orientations near a singular arc

Suppose that for all but a finite number of values $i \in]0, 1[$, the plane $W \times i$ intersects u generically. Suppose also that $W \times 0$ and $W \times 1$ intersect u generically and outside the vertices of s_γ . We call $W \times I \cap u$ an *open prefoam*. Interpreted as morphisms we read open prefoams from bottom to top, and their composition consists of placing one prefoam on top of the other, as long as their boundaries are isotopic and the orientations of the simple edges coincide.

Definition 4.2 Let **Prefoam** be the category whose objects are closed webs and whose morphisms are \mathbb{Q} -linear combinations of isotopy classes of prefoams with the obvious identity prefoams and composition rule.

We now define the q -degree of a prefoam. Let u be a prefoam, u_1 , u_2 and u_3 the disjoint union of its simple and double and marked facets respectively and $s_\gamma(u)$ its

singular graph. Define the partial q -gradings of u as

$$q_i(u) = \chi(u_i) - \frac{1}{2}\chi(\partial u_i \cap \partial u), \quad i = 1, 2, 3$$

$$q_{s_\gamma}(u) = \chi(s_\gamma(u)) - \frac{1}{2}\chi(\partial s_\gamma(u)),$$

where χ is the Euler characteristic and ∂ denotes the boundary.

Definition 4.3 Let u be a prefoam with d_\bullet dots of type \bullet , d_\circ dots of type \circ and d_\odot dots of type \odot . The q -grading of u is given by

$$(7) \quad q(u) = -\sum_{i=1}^3 i(N-i)q_i(u) - 2(N-2)q_{s_\gamma}(u) + 2d_\bullet + 4d_\circ + 6d_\odot.$$

The following result is a direct consequence of the definitions.

Lemma 4.4 $q(u)$ is additive under the gluing of prefoams.

5 The Kapustin–Li formula and the evaluation of closed prefoams

Let us briefly recall the philosophy behind the prefoams. Recall that we want to associate a number to each closed prefoam. To do this, we could first associate an element in the homology ring of a configuration space of planes in some big \mathbb{C}^M to a prefoam. To each i -facet we can associate an arbitrary i -plane in \mathbb{C}^M . We consider the subspace of all such planes with orthogonality conditions imposed by the singular graph. The set of dots could be interpreted as a cocycle on the same big configuration space. The evaluation of a prefoam with dots should then correspond to the pairing of the fundamental cycle of this subspace and the cocycle corresponding to the dots. However, one encounters difficult technical problems when working out the details of this philosophy. Without explaining these details, we can say that one problem consists of figuring out what to associate to the singular vertices. Ideally we would like to find a combinatorial solution to this problem, but so far it has eluded us. That is the reason why we are forced to use the Kapustin–Li formula.

We denote a simple facet with i dots by:



Recall that $\pi_{k,m}$ can be expressed in terms of $\pi_{1,0}$ and $\pi_{1,1}$. In the philosophy explained above, the latter should correspond to \bullet and \circ on a double facet respectively. We can then define

$$(k,m)$$

as being the linear combination of dotted double facets corresponding to the expression of $\pi_{k,m}$ in terms of $\pi_{1,0}$ and $\pi_{1,1}$. Analogously we expressed $\pi_{p,q,r}$ in terms of $\pi_{1,0,0}$, $\pi_{1,1,0}$ and $\pi_{1,1,1}$ (see Section 3). The latter correspond to \bullet , \circ and \odot on a triple facet respectively, so we can make sense of:

$$\frac{*}{(p,q,r)}$$

Our dot conventions and the results in Proposition 6.2 will allow us to use decorated facets in exactly the same way as we did Schur polynomials in the cohomology rings of partial flag varieties.

In the sequel, we shall give a working definition of the Kapustin–Li formula for the evaluation of prefoams and state some of its basic properties. The Kapustin–Li formula was introduced by A Kapustin and Y Li [6] in the context of the evaluation of 2-dimensional TQFTs and extended to the case of prefoams by M Khovanov and L Rozansky in [9].

5.1 The general framework

Let $u = \Sigma/s_\gamma$ be a closed prefoam with singular graph s_γ and without any dots on it. Let F denote an arbitrary i -facet, $i \in \{1, 2, 3\}$, with a 1-facet being a simple facet, a 2-facet being a double facet and a 3-facet being a triple facet.

Recall that to each i -facet we associated the rational cohomology ring of the Grassmannian $\mathcal{G}_{i,N}$, ie $H^*(\mathcal{G}_{i,N}, \mathbb{Q})$. Alternatively, we can associate to every i -facet F , i variables x_1^F, \dots, x_i^F , with $\deg x_i^F = 2i$, and the potential $W(x_1^F, \dots, x_i^F)$, which is the polynomial defined such that

$$W(\sigma_1, \dots, \sigma_i) = y_1^{N+1} + \dots + y_i^{N+1},$$

where σ_j is the j -th elementary symmetric polynomial in the variables y_1, \dots, y_i . The Jacobi algebra J_W , which is given by

$$J_W = \mathbb{Q}[x_1^F, \dots, x_i^F]/\langle \partial W \rangle$$

where we mod out by the ideal generated by the partial derivatives of W , is isomorphic to $H^*(\mathcal{G}_{i,N}, \mathbb{Q})$. Note that the top degree nonvanishing element in this Jacobi algebra is $\pi_{N-i, \dots, N-i}$ (multi-index of length i), ie the polynomial in variables x_1^F, \dots, x_i^F which gives $\pi_{N-i, \dots, N-i}$ after replacing the variable x_j^F by $\pi_{1, \dots, 1, 0, \dots, 0}$ with exactly j 1's, $1 \leq j \leq i$ (see also Section 3.1). We define the trace (volume) form, ϵ , on the cohomology ring of the Grassmannian, by giving it on the basis of the Schur polynomials:

$$\epsilon(\pi_{j_1, \dots, j_i}) = \begin{cases} (-1)^{\lfloor i/2 \rfloor} & \text{if } (j_1, \dots, j_i) = (N-i, \dots, N-i) \\ 0 & \text{else.} \end{cases}$$

The Kapustin–Li formula associates to u an element in the product of the cohomology rings of the Grassmannians (or Jacobi algebras), J , over all the facets in the prefoam. Alternatively, we can see this element as a polynomial, $\text{KL}_u \in J$, in all the variables associated to the facets. Now, let us put some dots on u . Recall that a dot corresponds to an elementary symmetric polynomial. So a linear combination of dots on u is equivalent to a polynomial, f , in the variables of the dotted facets. The value of this dotted prefoam we define to be

$$(8) \quad \langle u \rangle_{\text{KL}} := \epsilon \left(\prod_F \frac{\det(\partial_i \partial_j W_F)^{g(F)}}{(N+1)^{g'(F)}} \text{KL}_u f \right).$$

The product is over all facets F and W_F is the potential associated to F . For any i -facet F , $i = 1, 2, 3$, the symbol $g(F)$ denotes the genus of F and $g'(F) = ig(F)$.

Having explained the general idea, we are left with defining the element KL_u for a dotless prefoam. For that we have to explain Khovanov and Rozansky’s extension of the Kapustin–Li formula to prefoams [9], which uses the theory of matrix factorizations.

5.2 Matrix factorizations

Let $R = \mathbb{Q}[x_1, \dots, x_k]$ be a polynomial ring, and $W \in R$. By a matrix factorization over ring R with the potential W we mean a triple (M, D, W) , where $M = M^0 \oplus M^1$ ($\text{rank } M^0 = \text{rank } M^1$) is a finite-dimensional $\mathbb{Z}/2\mathbb{Z}$ -graded free R -module, while the (twisted) differential $D \in \text{End}(M)$ is such that $\text{deg } D = 1$ and

$$(9) \quad D^2 = W \text{Id}.$$

In other words, a matrix factorization is given by the following square matrix with polynomial entries

$$D = \begin{bmatrix} 0 & D_0 \\ D_1 & 0 \end{bmatrix},$$

such that $D_0 D_1 = D_1 D_0 = W \text{Id}$. Matrix factorizations are also represented in the following form:

$$M^0 \xrightarrow{D_0} M^1 \xrightarrow{D_1} M^0.$$

The tensor product of two matrix factorizations with potentials W_1 and W_2 is a matrix factorization with potential $W_1 + W_2$.

The dual of the matrix factorization (M, D, W) is given by

$$(M, D, W)^* = (M^*, D^*, -W),$$

where

$$D^* = \begin{bmatrix} 0 & D_1^* \\ -D_0^* & 0 \end{bmatrix}$$

and D_i^* , $i = 0, 1$, is the dual map (transpose matrix) of D_i .

Throughout the paper we shall use a particular type of matrix factorizations – namely the tensor products of Koszul factorizations. For two elements $a, b \in R$, the Koszul factorization $\{a, b\}$ is defined as the matrix factorization

$$R \xrightarrow{a} R \xrightarrow{b} R.$$

Moreover if $a = (a_1, \dots, a_m) \in R^m$ and $b = (b_1, \dots, b_m) \in R^m$, then the tensor product of the Koszul factorizations $\{a_i, b_i\}$, $i = 1, \dots, m$, is denoted by

$$(10) \quad \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \\ \vdots & \vdots \\ a_m & b_m \end{pmatrix} := \bigotimes_{i=1}^m \{a_i, b_i\}.$$

Sometimes we also write $\{a, b\} = \bigotimes_{i=1}^m \{a_i, b_i\}$. If $\sum_{i=1}^m a_i b_i = 0$ then $\{a, b\}$ is a 2-periodic complex, and its homology is an $R/\langle a_1, \dots, a_m, b_1, \dots, b_m \rangle$ -module.

5.3 Decoration of prefoams

As we said, to each facet we associate certain variables (depending on the type of facet), a potential and the corresponding Jacobi algebra. If the variables associated to a facet F are x_1, \dots, x_i , then we define $R_F = \mathbb{Q}[x_1, \dots, x_i]$.

Now we pass to the edges. To each edge, we associate a matrix factorization whose potential is equal to the signed sum of the potentials of the facets that are glued along this edge. We define it to be a certain tensor product of Koszul factorizations. In the cases we are interested in, there are always three facets glued along an edge, with two

possibilities: either two simple facets and one double facet, or one simple, one double and one triple facet.

In the first case, we denote the variables of the two simple facets by x and y and the potentials by x^{N+1} and y^{N+1} respectively. To the double facet we associate the variables s and t and the potential $W(s, t)$. To the edge we associate the matrix factorization which is the tensor product of Koszul factorizations given by

$$(11) \quad \text{MF}_1 = \begin{pmatrix} x + y - s & A' \\ xy - t & B' \end{pmatrix},$$

where A' and B' are given by

$$A' = \frac{W(x + y, xy) - W(s, xy)}{x + y - s},$$

$$B' = \frac{W(s, xy) - W(s, t)}{xy - t}.$$

Note that $(x + y - s)A' + (xy - t)B' = x^{N+1} + y^{N+1} - W(s, t)$.

In the second case, the variable of the simple facet is x and the potential is x^{N+1} , the variables of the double facet are s and t and the potential is $W(s, t)$, and the variables of the triple face are p, q and r and the potential is $W(p, q, r)$. Define the polynomials

$$(12) \quad A = \frac{W(x + s, xs + t, xt) - W(p, xs + t, xt)}{x + s - p},$$

$$(13) \quad B = \frac{W(p, xs + t, xt) - W(p, q, xt)}{xs + t - q},$$

$$(14) \quad C = \frac{W(p, q, xt) - W(p, q, r)}{xt - r},$$

so that

$$(x + s - p)A + (xs + t - q)B + (xt - r)C = x^{N+1} + W(s, t) - W(p, q, r).$$

To such an edge we associate the matrix factorization given by the following tensor product of Koszul factorizations:

$$(15) \quad \text{MF}_2 = \begin{pmatrix} x + s - p & A \\ xs + t - q & B \\ xt - r & C \end{pmatrix}.$$

In both cases, to an edge with the opposite orientation we associate the dual matrix factorization.

Next we explain what we associate to a singular vertex. First of all, for each vertex v , we define its local graph γ_v to be the intersection of a small sphere centered at v with the prefoam. Then the vertices of γ_v correspond to the edges of u that are incident to v , to which we had associated matrix factorizations.

In this paper all local graphs are in fact tetrahedrons. However, recall that there are two types of vertices (see the remarks below Definition 4.1). Label the six facets that are incident to a vertex v by the numbers 1, 2, 3, 4, 5 and 6. Furthermore, denote the edge along which are glued the facets i , j and k by (ijk) . Denote the matrix factorization associated to the edge (ijk) by M_{ijk} , if the edge points toward v , and by M_{ijk}^* , if the edge points away from v . Note that M_{ijk} and M_{ijk}^* are both defined over $R_i \otimes R_j \otimes R_k$.

Now we can take the tensor product of these four matrix factorizations, over the polynomial rings of the facets of the prefoam, that correspond to the vertices of γ_v . This way we obtain the matrix factorization M_v , whose potential is equal to 0, and so it is a 2-periodic chain complex and we can take its homology. To each vertex v we associate an element $O_v \in H^*(M_v)$.

More precisely, if v is of type I, then

$$(16) \quad H^*(M_v) \cong \text{Ext} \left(\text{MF}_1(x, y, s_1, t_1) \otimes_{s_1, t_1} \text{MF}_2(z, s_1, t_1, p, q, r), \right. \\ \left. \text{MF}_1(y, z, s_2, t_2) \otimes_{s_2, t_2} \text{MF}_2(x, s_2, t_2, p, q, r) \right).$$

If v is of type II, then

$$(17) \quad H^*(M_v) \cong \text{Ext} \left(\text{MF}_1(y, z, s_2, t_2) \otimes_{s_2, t_2} \text{MF}_2(x, s_2, t_2, p, q, r), \right. \\ \left. \text{MF}_1(x, y, s_1, t_1) \otimes_{s_1, t_1} \text{MF}_2(z, s_1, t_1, p, q, r) \right).$$

Both isomorphisms hold up to a global shift in q . Note that

$$\text{MF}_1(x, y, s_1, t_1) \otimes_{s_1, t_1} \text{MF}_2(z, s_1, t_1, p, q, r) \\ \simeq \text{MF}_1(y, z, s_2, t_2) \otimes_{s_2, t_2} \text{MF}_2(x, s_2, t_2, p, q, r),$$

because both tensor products are homotopy equivalent to

$$\begin{pmatrix} x + y + z - p & * \\ xy + xz + yz - q & * \\ xyz - r & * \end{pmatrix}.$$

We have not specified the right-hand side of the latter Koszul factorizations, because by [10, Theorem 2.1] we have $\{a, b\} \simeq \{a, b'\}$ if $\sum a_i b_i = \sum a_i b'_i$ and if the sequence

$\{a_i\}$ is regular. If v is of type I, we take O_v to be the cohomology class of a fixed degree 0 homotopy equivalence

$$w_v: \text{MF}_1(x, y, s_1, t_1) \otimes_{s_1, t_1} \text{MF}_2(z, s_1, t_1, p, q, r) \rightarrow \text{MF}_1(y, z, s_2, t_2) \otimes_{s_2, t_2} \text{MF}_2(x, s_2, t_2, p, q, r).$$

The choice of O_v is unique up to a scalar, because the q -dimension of the Ext-group in (16) is equal to

$$q^{3N-6} \text{qdim}(H^*(M_v)) = q^{3N-6} [N][N-1][N-2] = 1 + q(\dots),$$

where (\dots) is a polynomial in q . Note that M_v is homotopy equivalent to the matrix factorization which corresponds to the closure of Υ in [11], which allows one to compute the q -dimension above using the results in the latter paper. If v is of type II, we take O_v to be the cohomology class of the homotopy inverse of w_v . Note that a particular choice of w_v fixes O_v for both types of vertices and that the value of the Kapustin–Li formula for a closed prefoam does not depend on that choice because there are as many singular vertices of type I as there are of type II (see the remarks below Definition 4.1). We do not know an explicit formula for O_v . Although such a formula would be very interesting to have, we do not need it for the purposes of this paper.

5.4 The Kapustin–Li derivative and the evaluation of closed prefoams

From the definition, every boundary component of each facet F is either a circle or a cyclicly ordered finite sequence of edges, such that the beginning of the next edge corresponds to the end of the previous edge. For every boundary component we choose an edge e - the value of the Kapustin–Li formula does not depend on this choice. Denote the differential of the matrix factorization associated to this edge by D_e .

The associated Kapustin–Li derivative of D_e in the variables x_1, \dots, x_k associated to the facet F , is an element from $\text{End}(M) \cong M \otimes M^*$, given by

$$(18) \quad O_{F,e} = \partial D_e^\wedge = \frac{1}{k!} \sum_{\sigma \in S_k} (\text{sgn } \sigma) \partial_{\sigma(1)} D_e \partial_{\sigma(2)} D_e \dots \partial_{\sigma(k)} D_e,$$

where S_k is the set of all permutations of the set $\{1, \dots, k\}$, and $\partial_i D$ is the partial derivative of D with respect to the variable x_i . Note that e can be the preferred edge for more than one facet. In general, let O_e be the product of $O_{F,e}$ over all facets F for which e is the preferred edge. The order of the factors in this product is irrelevant, because they commute [9]. If e is not the preferred edge for any F , we take O_e to be the identity.

Finally, around each boundary component of ∂F , for each facet F , we contract all tensor factors O_e and O_v . Note that one has to use supercontraction in order to get the right signs.

For a better understanding of the Kapustin–Li formula, consider the special case of a theta prefoam Θ . There are three facets F_1, F_2, F_3 which are glued along a common circle c , which is the preferred edge for all three. We associated a certain matrix factorization M to c with differential D . Let $\partial D_1^\wedge, \partial D_2^\wedge$ and ∂D_3^\wedge be the Kapustin–Li derivatives of D with respect to the variables of the facets F_1, F_2 and F_3 , respectively. Then we have

$$(19) \quad \text{KL}_\Theta = \text{Tr}^s(\partial D_1^\wedge \partial D_2^\wedge \partial D_3^\wedge).$$

As a matter of fact we will see that we have to *normalize* the Kapustin–Li formula in order to get “nice values”.

5.5 Dot conversion and dot migration

The pictures related to the computations in this subsection and the next three can be found in Proposition 6.2.

Since KL_u takes values in the tensor product of the Jacobian algebras of the potentials associated to the facets of u , we see that for a simple facet we have $x^N = 0$, for a double facet $\pi_{i,j} = 0$ if $i \geq N - 1$, and for a triple facet $\pi_{p,q,r} = 0$ if $p \geq N - 2$. We call these the *dot conversion relations*.

To each edge along which two simple facets with variables x and y and one double facet with the variables s and t are glued, we associated the matrix factorization MF_1 with entries $x + y - s$ and $xy - t$. Therefore $\text{Ext}(\text{MF}_1, \text{MF}_1)$ is a module over $R/\langle x + y - s, xy - t \rangle$. Hence, we obtain the *dot migration relations* along this edge.

Analogously, to the other type of singular edge along which are glued a simple facet with variable x , a double facet with variable s and t , and a triple facet with variables p, q and r , we associated the matrix factorization MF_2 and $\text{Ext}(\text{MF}_2, \text{MF}_2)$ is a module over $R/\langle x + s - p, xs + t - q, xt - r \rangle$, and hence we obtain the *dot migration relations* along this edge.

5.6 (1, 1, 2)–Theta

In this subsection we prove the first Θ –foam relation in Proposition 6.2, for the theta-foam with two simple and one double facet. Recall that $W(s, t)$ is the polynomial such

that $W(x + y, xy) = x^{N+1} + y^{N+1}$. More precisely, we have

$$(20) \quad W(s, t) = \sum_{i+2j=N+1} a_{ij} s^i t^j,$$

with $a_{N+1,0} = 1$, $a_{N+1-2j,j} = ((-1)^j/j)(N + 1) \binom{N-j}{j-1}$, for $2 \leq 2j \leq N + 1$, and $a_{ij} = 0$ otherwise. In particular $a_{N-1,1} = -(N + 1)$. Then we have

$$(21) \quad W'_1(s, t) = \sum_{i+2j=N+1} i a_{ij} s^{i-1} t^j,$$

$$(22) \quad W'_2(s, t) = \sum_{i+2j=N+1} j a_{ij} s^i t^{j-1}.$$

By $W'_1(s, t)$ and $W'_2(s, t)$, we denote the partial derivatives of $W(s, t)$ with respect to the first and the second variable, respectively.

To the singular circle of a standard theta prefoam with two simple facets, with variables x and y respectively, and one double facet, with variables s and t , we assign the matrix factorization MF_1 :

$$(23) \quad \text{MF}_1 = \begin{pmatrix} x + y - s & A' \\ xy - t & B' \end{pmatrix}.$$

Recall that

$$(24) \quad A' = \frac{W(x + y, xy) - W(s, xy)}{x + y - s},$$

$$(25) \quad B' = \frac{W(s, xy) - W(s, t)}{xy - t}.$$

Hence, the differential of this matrix factorization is given by the 4-by-4 matrix

$$(26) \quad D = \begin{bmatrix} 0 & D_0 \\ D_1 & 0 \end{bmatrix},$$

where
$$D_0 = \begin{bmatrix} x + y - s & -B' \\ xy - t & A' \end{bmatrix}, \quad D_1 = \begin{bmatrix} A' & B' \\ t - xy & x + y - s \end{bmatrix}.$$

The Kapustin–Li formula assigns the polynomial, $\text{KL}_{\Theta_1}(x, y, s, t)$, which is given by the supertrace of the twisted differential of D

$$(27) \quad \text{KL}_{\Theta_1} = \text{Tr}^s \left(\partial_x D \partial_y D \frac{1}{2} (\partial_s D \partial_t D - \partial_t D \partial_s D) \right).$$

Straightforward computation gives

$$(28) \quad \text{KL}_{\Theta_1} = B'_s(A'_x - A'_y) + (A'_x + A'_s)(B'_y + xB'_t) - (A'_y + A'_s)(B'_x + yB'_t),$$

where by A'_i and B'_i we have denoted the partial derivatives with respect to the variable i . From the definitions (24) and (25) we have

$$\begin{aligned} A'_x - A'_y &= (y-x) \frac{W'_2(x+y, xy) - W'_2(s, xy)}{x+y-s}, \\ A'_x + A'_s &= \frac{W'_1(x+y, xy) - W'_1(s, xy) + y(W'_2(x+y, xy) - W'_2(s, xy))}{x+y-s}, \\ A'_y + A'_s &= \frac{W'_1(x+y, xy) - W'_1(s, xy) + x(W'_2(x+y, xy) - W'_2(s, xy))}{x+y-s}, \\ B'_s &= \frac{W'_1(s, xy) - W'_1(s, t)}{xy-t}, \\ B'_x + yB'_t &= y \frac{W'_2(s, xy) - W'_2(s, t)}{xy-t}, \\ B'_y + xB'_t &= x \frac{W'_2(s, xy) - W'_2(s, t)}{xy-t}. \end{aligned}$$

After substituting this back into (28), we obtain

$$(29) \quad \text{KL}_{\Theta_1} = (y-x) \begin{vmatrix} \alpha & \beta \\ \gamma & \delta \end{vmatrix},$$

where

$$\begin{aligned} \alpha &= \frac{W'_1(x+y, xy) - W'_1(s, xy)}{x+y-s}, \\ \beta &= \frac{W'_2(x+y, xy) - W'_2(s, xy)}{x+y-s}, \\ \gamma &= \frac{W'_1(s, xy) - W'_1(s, t)}{xy-t}, \\ \delta &= \frac{W'_2(s, xy) - W'_2(s, t)}{xy-t}. \end{aligned}$$

From this formula we see that KL_{Θ_1} is homogeneous of degree $4N - 6$ (remember that $\deg x = \deg y = \deg s = 2$ and $\deg t = 4$).

Since the evaluation is in the product of the Grassmannians corresponding to the three disks, ie in the ring

$$\mathbb{Q}[x]/x^N \times \mathbb{Q}[y]/y^N \times \mathbb{Q}[s, t]/\langle W'_1(s, t), W'_2(s, t) \rangle,$$

we have $x^N = y^N = 0 = W'_1(s, t) = W'_2(s, t)$. Also, we can express the monomials in s and t as linear combinations of the Schur polynomials $\pi_{k,l}$ (writing $s = \pi_{1,0}$ and $t = \pi_{1,1}$), and we have $W'_1(s, t) = (N + 1)\pi_{N,0}$ and $W'_2(s, t) = -(N + 1)\pi_{N-1,0}$. Hence, we can write KL_{Θ_1} as

$$\text{KL}_{\Theta_1} = (y - x) \sum_{N-2 \geq k \geq l \geq 0} \pi_{k,l} p_{kl}(x, y),$$

with p_{kl} being a polynomial in x and y . We want to determine which combinations of dots on the simple facets give rise to nonzero evaluations, so our aim is to compute the coefficient of $\pi_{N-2,N-2}$ in the sum on the right-hand side of the above equation (ie in the determinant in (29)). For degree reasons, this coefficient is of degree zero, and so we shall only compute the parts of α , β , γ and δ which do not contain x and y . We shall denote these parts by putting a bar over the Greek letters. Thus we have

$$\begin{aligned} \bar{\alpha} &= (N + 1)s^{N-1}, \\ \bar{\beta} &= -(N + 1)s^{N-2}, \\ \bar{\gamma} &= \sum_{i+2j=N+1, j \geq 1} i a_{ij} s^{i-1} t^{j-1}, \\ \bar{\delta} &= \sum_{i+2j=N+1, j \geq 2} j a_{ij} s^i t^{j-2}. \end{aligned}$$

Note that we have

$$t\bar{\gamma} + (N + 1)s^N = W'_1(s, t),$$

and

$$t\bar{\delta} - (N + 1)s^{N-1} = W'_2(s, t),$$

and so in the cohomology ring of the Grassmannian $\mathcal{G}_{2,N}$, we have $t\bar{\gamma} = -(N + 1)s^N$ and $t\bar{\delta} = (N + 1)s^{N-1}$. On the other hand, by using $s = \pi_{1,0}$ and $t = \pi_{1,1}$, we obtain that in

$$H^*(\mathcal{G}_{2,N}) \cong \mathbb{Q}[s, t] / \langle \pi_{N-1,0}, \pi_{N,0} \rangle,$$

the following holds

$$s^{N-2} = \pi_{N-2,0} + tq(s, t),$$

for some polynomial q , and so

$$s^{N-1} = s^{N-2}s = \pi_{N-1,0} + \pi_{N-2,1} + stq(s, t) = t(\pi_{N-3,0} + sq(s, t)).$$

Thus, we have

$$\begin{aligned}
 (30) \quad \left| \frac{\bar{\alpha}}{\bar{\gamma}} \frac{\bar{\beta}}{\bar{\delta}} \right| &= (N + 1)(\pi_{N-3,0} + sq(s, t))t\bar{\delta} + (N + 1)\pi_{N-2,0}\bar{\gamma} + (N + 1)q(s, t)t\bar{\gamma} \\
 &= (N + 1)^2(\pi_{N-3,0} + sq(s, t))s^{N-1} + (N + 1)\pi_{N-2,0}\bar{\gamma} \\
 &\qquad\qquad\qquad - (N + 1)^2q(s, t)s^N \\
 &= (N + 1)^2\pi_{N-3,0}s^{N-1} + (N + 1)\pi_{N-2,0}\bar{\gamma}.
 \end{aligned}$$

Since
$$\bar{\gamma} = (N - 1)a_{N-1,1}s^{N-2} + tr(s, t)$$

holds in the cohomology ring of the Grassmannian, for some polynomial $r(s, t)$, we have

$$\pi_{N-2,0}\bar{\gamma} = \pi_{N-2,0}(N - 1)a_{N-1,1}s^{N-2} = -\pi_{N-2,0}(N - 1)(N + 1)s^{N-2}.$$

Also, we have that for every $k \geq 2$,

$$s^k = \pi_{k,0} + (k - 1)\pi_{k-1,1} + t^2w(s, t),$$

for some polynomial w . Replacing this in (30) and bearing in mind that $\pi_{i,j} = 0$, for $i \geq N - 1$, we get

$$\begin{aligned}
 (31) \quad \left| \frac{\bar{\alpha}}{\bar{\gamma}} \frac{\bar{\beta}}{\bar{\delta}} \right| &= (N + 1)^2s^{N-2}(\pi_{N-2,0} + \pi_{N-3,1} - (N - 1)\pi_{N-2,0}) \\
 &= (N + 1)^2(\pi_{N-2,0} + (N - 3)\pi_{N-3,1} + \pi_{2,2}w(s, t)) \\
 &\qquad\qquad\qquad \times (\pi_{N-3,1} - (N - 2)\pi_{N-2,0}) \\
 &= -(N + 1)^2\pi_{N-2,N-2}.
 \end{aligned}$$

Hence, we have

$$\text{KL}_{\Theta_1} = (N + 1)^2(x - y)\pi_{N-2,N-2} + \sum_{\substack{N-2 \geq k \geq l \geq 0 \\ N-2 > l}} c_{i,j,k,l}\pi_{k,l}x^i y^j.$$

Recall that in the product of the Grassmannians corresponding to the three disks, ie in the ring $\mathbb{Q}[x]/x^N \times \mathbb{Q}[y]/y^N \times \mathbb{Q}[s, t]/\langle \pi_{N-1,0}, \pi_{N,0} \rangle$, we have

$$\epsilon(x^{N-1}y^{N-1}\pi_{N-2,N-2}) = -1.$$

Therefore the only monomials f in x and y such that $\langle \text{KL}_{\Theta_1} f \rangle_{\text{KL}} \neq 0$ are $f_1 = x^{N-2}y^{N-1}$ and $f_2 = x^{N-1}y^{N-2}$, and $\langle \text{KL}_{\Theta_1} f_1 \rangle_{\text{KL}} = -(N + 1)^2$ and $\langle \text{KL}_{\Theta_1} f_2 \rangle_{\text{KL}} = (N + 1)^2$. Thus, we have that the value of the theta prefoam with unlabelled 2-facet is nonzero only when the first 1-facet has $N - 2$ dots and the second one has $N - 1$ dots (and has the value $-(N + 1)^2$) and when the first 1-facet has $N - 1$ dots and the

second one has $N - 2$ dots (and has the value $(N + 1)^2$). The evaluation of this theta foam with other labellings can be obtained from the result above by dot migration.

5.7 (1, 2, 3)-Theta

In this subsection we prove the second Θ -foam relation in Proposition 6.2, for the theta-foam with one simple, one double and one marked facet. For this theta the method is the same as in the previous case, just the computations are more complicated. In this case, we have one 1-facet, to which we associate the variable x , one 2-facet, with variables s and t and the 3-facet with variables p, q and r . Recall that the polynomial $W(p, q, r)$ is such that $W(a + b + c, ab + bc + ac, abc) = a^{N+1} + b^{N+1} + c^{N+1}$. We denote by $W'_i(p, q, r)$, $i = 1, 2, 3$, the partial derivative of W with respect to i -th variable. Also, let A, B and C be the polynomials such that

$$(32) \quad A = \frac{W(x + s, xs + t, xt) - W(p, xs + t, xt)}{x + s - p},$$

$$(33) \quad B = \frac{W(p, xs + t, xt) - W(p, q, xt)}{xs + t - q},$$

$$(34) \quad C = \frac{W(p, q, xt) - W(p, q, r)}{xt - r}.$$

To the singular circle of this theta prefoam, we associated the matrix factorization (see (12)–(15)):

$$\text{MF}_2 = \begin{pmatrix} x + s - p & A \\ xs + t - q & B \\ xt - r & C \end{pmatrix}.$$

The differential of this matrix factorization is the 8 by 8 matrix

$$D = \begin{bmatrix} 0 & D_0 \\ D_1 & 0 \end{bmatrix},$$

where

$$D_0 = \left[\begin{array}{c|c} d_0 & -C I_2 \\ \hline (xt - r) I_2 & d_1 \end{array} \right],$$

$$D_1 = \left[\begin{array}{c|c} d_1 & C I_2 \\ \hline (r - xt) I_2 & d_0 \end{array} \right].$$

Here d_0 and d_1 are the differentials of the matrix factorization

$$\begin{pmatrix} x + s - p & A \\ xs + t - q & B \end{pmatrix},$$

that is,
$$d_0 = \begin{bmatrix} x + s - p & -B \\ xs + t - q & A \end{bmatrix}, \quad d_1 = \begin{bmatrix} A & B \\ q - xs - t & x + s - p \end{bmatrix}.$$

The Kapustin–Li formula assigns to this prefoam the polynomial $\text{KL}_{\Theta_2}(x, s, t, p, q, r)$ given as the supertrace of the twisted differential of D , ie

$$\text{KL}_{\Theta_2} = \text{Tr}^s \left(\partial_x D \frac{1}{2} (\partial_s D \partial_t D - \partial_t D \partial_s D) \partial_3 D^\wedge \right),$$

where

$$\partial_3 D^\wedge = \frac{1}{3!} (\partial_p D \partial_q D \partial_r D - \partial_p D \partial_r D \partial_q D + \partial_q D \partial_r D \partial_p D - \partial_q D \partial_p D \partial_r D + \partial_r D \partial_p D \partial_q D - \partial_r D \partial_q D \partial_p D).$$

After straightforward computations and some grouping, we obtain

$$\begin{aligned} \text{KL}_{\Theta_2} = & -(A_p + A_s)((B_t + B_q)(C_x + tC_r) - (B_x + sB_q)(C_t + xC_r) \\ & \quad - (B_x - sB_t)C_q) \\ & - (A_p + A_x)((B_s + xB_q)(C_t + xC_r) + (B_s - xB_t)C_q) \\ & - (A_x - A_s)(B_p(C_t + xC_r) - (B_t + B_q)C_p + B_pC_q) \\ & + A_t(((B_s + xB_q) + B_p)(C_x + tC_r) + ((B_s + xB_q) \\ & \quad + (B_x + sB_q))C_p + ((sB_s - xB_x) + (s - x)B_p)C_q). \end{aligned}$$

In order to simplify this expression, we introduce the following polynomials:

$$\begin{aligned} a_{1i} &= \frac{W'_i(x + s, xs + t, xt) - W'_i(p, xs + t, xt)}{x + s - p}, & i = 1, 2, 3, \\ a_{2i} &= \frac{W'_i(p, xs + t, xt) - W'_i(p, q, xt)}{xs + t - q}, & i = 1, 2, 3, \\ a_{3i} &= \frac{W'_i(p, q, xt) - W'_i(p, q, r)}{xt - r}, & i = 1, 2, 3. \end{aligned}$$

Then from (32)–(34), we have

$$\begin{aligned} A_x + A_p &= a_{11} + sa_{12} + ta_{13}, & A_p + A_s &= a_{11} + xa_{12}, \\ A_x - A_s &= (s - x)a_{12} + ta_{13}, & A_t &= a_{12} + xa_{13}, \\ B_p &= a_{21}, & B_s - xB_t &= -x^2a_{23}, \\ sB_s - xB_x &= xta_{23}, & B_x - sB_t &= (t - sx)a_{23}, \\ B_t + B_q &= a_{22} + xa_{23}, & B_x + sB_q &= sa_{22} + ta_{23}, & B_s + xB_q &= xa_{22}, \\ C_p &= a_{31}, & C_q &= a_{32}, & C_x + tC_r &= ta_{33}, & C_t + xC_r &= xa_{33}. \end{aligned}$$

Using this KL_{Θ_2} becomes

$$\text{KL}_{\Theta_2} = -(t - sx + x^2) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}.$$

Now the last part follows analogously as in the case of the $(1, 1, 2)$ -theta prefoam. For degree reasons the coefficient of $\pi_{N-3, N-3, N-3}$ in the latter determinant is of degree zero, and one can obtain that it is equal to $(N + 1)^3$. Thus, the coefficient of $\pi_{N-3, N-3, N-3}$ in KL_{Θ_2} is $-(N + 1)^3(t - sx + x^2)$ from which we obtain the value of the theta prefoam when the 3-facet is undotted. For example, we see that

$$\epsilon(\text{KL}_{\Theta_2} \pi_{1,1}(s, t)^{N-3} x^{N-1}) = -(N + 1)^3.$$

It is then easy to obtain the values when the 3-facet is labelled by

$$\pi_{N-3, N-3, N-3}(p, q, r)$$

using dot migration. The example above implies that

$$\epsilon(\text{KL}_{\Theta_2} \pi_{N-3, N-3, N-3}(p, q, r)x^2) = -(N + 1)^3.$$

5.8 Spheres

The values of dotted spheres are easy to compute. Note that for any sphere with dots f the Kapustin–Li formula gives

$$\epsilon(f).$$

Therefore for a simple sphere we get 1 if $f = x^{N-1}$, for a double sphere we get -1 if $f = \pi_{N-2, N-2}$ and for a triple sphere we get -1 if $f = \pi_{N-3, N-3, N-3}$.

5.9 Normalization

It will be convenient to normalize the Kapustin–Li evaluation. Let u be a closed prefoam with graph Γ . Note that Γ has two types of edges: the ones incident to two simple facets and one double facet and the ones incident to one simple, one double and one triple facet. Edges of the same type form cycles in Γ . Let $e_{112}(u)$ be the total number of cycles in Γ with edges of the first type and $e_{123}(u)$ the total number of cycles with edges of the second type. We normalize the Kapustin–Li formula by dividing KL_u by

$$(N + 1)^{2e_{112} + 3e_{123}}.$$

In the sequel we only use this normalized Kapustin–Li evaluation keeping the same notation $\langle u \rangle_{\text{KL}}$. Note that the numbers $e_{112}(u)$ and $e_{123}(u)$ are invariant under the

relation (MP). Note also that with this normalization the KL–evaluation in the examples above always gives 0, -1 or 1 .

5.10 The gluing property

If u is an open prefoam whose boundary consists of two parts Γ_1 and Γ_2 , then the Kapustin–Li formula associates to u an element from $\text{Ext}(M_1, M_2)$, where M_1 and M_2 are matrix factorizations associated to Γ_1 and Γ_2 respectively. If u' is another prefoam whose boundary consists of Γ_2 and Γ_3 , then it corresponds to an element from $\text{Ext}(M_2, M_3)$, while the element associated to the prefoam uu' , which is obtained by gluing the prefoams u and u' along Γ_2 , is equal to the composite of the elements associated to u and u' .

On the other hand, we can see u as a morphism from the empty set to its boundary $\Gamma = \Gamma_2 \cup \overline{\Gamma_1}$, where $\overline{\Gamma_1}$ is equal to Γ_1 but with the opposite orientation. In that case, the Kapustin–Li formula associates to it an element from

$$\text{Ext}(\emptyset, M_{\Gamma_2} \otimes M_{\Gamma_1}^*) \cong H^*(\Gamma).$$

Of course both ways of applying the Kapustin–Li formula are equivalent up to a global q –shift by [11, Corollary 6].

In the case of a prefoam u with corners, ie a prefoam with two horizontal boundary components Γ_1 and Γ_2 which are connected by vertical edges, one has to “pinch” the vertical edges. This way one can consider u to be a morphism from the empty set to $\Gamma_2 \cup_v \overline{\Gamma_1}$, where \cup_v means that the webs are glued at their vertices. The same observations as above hold, except that $M_{\Gamma_2} \otimes M_{\Gamma_1}^*$ is now the tensor product over the polynomial ring in the variables associated to the horizontal edges with corners.

6 The category \mathbf{Foam}_N

Recall that $\langle u \rangle_{\text{KL}}$ denotes the Kapustin–Li evaluation of a closed prefoam u .

Definition 6.1 The category \mathbf{Foam}_N is the quotient of the category $\mathbf{Prefoam}$ by the kernel of $\langle \rangle_{\text{KL}}$, ie by the following identifications: for any webs Γ, Γ' and finite sets $f_i \in \text{Hom}_{\mathbf{Prefoam}}(\Gamma, \Gamma')$ and $c_i \in \mathbb{Q}$ we impose the relations

$$\sum_i c_i f_i = 0 \iff \sum_i c_i \langle g' f_i g \rangle_{\text{KL}} = 0,$$

for all $g \in \text{Hom}_{\mathbf{Prefoam}}(\emptyset, \Gamma)$ and $g' \in \text{Hom}_{\mathbf{Prefoam}}(\Gamma', \emptyset)$. The morphisms of \mathbf{Foam}_N are called *foams*.

In the next two propositions we prove the “principal” relations in \mathbf{Foam}_N . All other relations that we need are consequences of these and will be proved in subsequent lemmas and corollaries.

Proposition 6.2 *The following identities hold in \mathbf{Foam}_N :*


The dot conversion relations:

$$\boxed{i} = 0 \quad \text{if } i \geq N.$$

$$\boxed{(k,m)} = 0 \quad \text{if } k \geq N - 1.$$

$$\boxed{^*(p,q,r)} = 0 \quad \text{if } p \geq N - 2.$$

The dot migration relations:



The cutting neck relations:

$$(CN_1) \quad \text{Cylinder} = \sum_{i=0}^{N-1} \left(\text{Cap } (N-1-i) + \text{Cap } i \right)$$

$$(CN_2) \quad \text{Cylinder} = - \sum_{0 \leq j \leq i \leq N-2} \left(\text{Cap } (i,j) + \text{Cap } (i,j) \right)$$

$$(CN_*) \quad \text{Cylinder with } * = - \sum_{0 \leq k \leq j \leq i \leq N-3} \begin{matrix} \text{Sphere with } * \\ (i,j,k) \end{matrix}$$

The sphere relations:

$$(S_1) \quad \text{Sphere with } i = \begin{cases} 1, & i = N - 1 \\ 0, & \text{else.} \end{cases}$$

$$(S_2) \quad \text{Sphere with } (i,j) = \begin{cases} -1, & i = j = N - 2 \\ 0, & \text{else.} \end{cases}$$

$$(S_*) \quad \text{Sphere with } (i,j,k) = \begin{cases} -1, & i = j = k = N - 3 \\ 0, & \text{else.} \end{cases}$$

The Θ -foam relations:

$$(\Theta) \quad \begin{matrix} \text{Theta-foam with } N-1 \\ \text{facets} \end{matrix} = -1 = - \begin{matrix} \text{Theta-foam with } N-2 \\ \text{facets} \end{matrix}$$

$$(\Theta_*) \quad \begin{matrix} \text{Theta-foam with } (N-3, N-3, N-3) \\ \text{facets and } * \end{matrix} = -1 = - \begin{matrix} \text{Theta-foam with } (N-3, N-3, N-3) \\ \text{facets and } * \end{matrix}$$

Inverting the orientation of the singular circle of (Θ_*) inverts the sign of the corresponding foam. A theta-foam with dots on the double facet can be transformed into a theta-foam with dots only on the other two facets, using the dot migration relations.

The Matveev–Piergalini relation:

$$(MP) \quad \text{Foam with } * = \text{Foam with } * \text{ and } \text{cutting neck} = \text{Foam with } * \text{ and } \text{cutting neck} = \text{Foam with } * \text{ and } \text{cutting neck}$$

Proof The dot conversion and migration relations, the sphere relations, the theta foam relations have already been proved in Section 5.

The cutting neck relations are special cases of Formula (5.68) in [9], where O_j and O_j^* can be read off from our equations (6).

The Matveev–Piergalini (MP) relation is an immediate consequence of the choice of input for the singular vertices. Note that in this relation there are always two singular

vertices of different type. The elements in the Ext-groups associated to those two types of singular vertices are inverses of each other, which implies exactly the (MP) relation by the gluing properties explained in Section 5.10. \square

The following identities are a consequence of the dot and the theta relations.

Lemma 6.3

$$\begin{array}{c} \text{Diagram of a theta foam with vertices } (p,q,r), (j,k), i \end{array} = \begin{cases} -1 & \text{if } (p, q, r) = (N - 3 - i, N - 2 - k, N - 2 - j) \\ -1 & \text{if } (p, q, r) = (N - 3 - k, N - 3 - j, N - 1 - i) \\ 1 & \text{if } (p, q, r) = (N - 3 - k, N - 2 - i, N - 2 - j) \\ 0 & \text{else.} \end{cases}$$

Note that the first three cases only make sense if

$$\begin{aligned} N - 2 &\geq j \geq k \geq i + 1 \geq 1, \\ N - 1 &\geq i \geq j + 2 \geq k + 2 \geq 2, \\ N - 2 &\geq j \geq i \geq k + 1 \geq 1, \end{aligned}$$

respectively.

Proof We denote the value of a theta foam by $\Theta(\pi_{p,q,r}, \pi_{j,k}, i)$. Since the q -degree of a nondecorated theta foam is equal to $-(N - 1) - 2(N - 2) - 3(N - 3) = -(6N - 14)$, we can have nonzero values of $\Theta(\pi_{p,q,r}, \pi_{j,k}, i)$ only if $p + q + r + j + k + i = 3N - 7$. Thus, if the 3-facet is not decorated, ie $p = q = r = 0$, we have only four possibilities for the triple (j, k, i) – namely $(N - 2, N - 2, N - 3)$, $(N - 2, N - 3, N - 2)$, $(N - 2, N - 4, N - 1)$ and $(N - 3, N - 3, N - 1)$. By Proposition 6.2 we have

$$\Theta(\pi_{0,0,0}, \pi_{N-2,N-2}, N - 3) = -1.$$

However by dot migration, Lemma 3.1 and the fact that $\pi_{p,q,r} = 0$ if $p \geq N - 2$, we have

$$\begin{aligned} 0 &= \Theta(\pi_{N-2,N-2,N-3}, \pi_{0,0}, 0) \\ &= \Theta(\pi_{0,0,0}, \pi_{N-2,N-2}, N - 3) + \Theta(\pi_{0,0,0}, \pi_{N-2,N-3}, N - 2), \\ 0 &= \Theta(\pi_{N-1,N-3,N-3}, \pi_{0,0}, 0) \\ &= \Theta(\pi_{0,0,0}, \pi_{N-2,N-3}, N - 2) + \Theta(\pi_{0,0,0}, \pi_{N-3,N-3}, N - 1), \\ 0 &= \Theta(\pi_{N-1,N-2,N-4}, \pi_{0,0}, 0) \\ &= \Theta(\pi_{0,0,0}, \pi_{N-2,N-2}, N - 3) + \Theta(\pi_{0,0,0}, \pi_{N-2,N-3}, N - 2) \\ &\quad + \Theta(\pi_{0,0,0}, \pi_{N-2,N-4}, N - 1). \end{aligned}$$

Thus, the only nonzero values of the theta foams, when the 3-facet is nondecorated are

$$\begin{aligned}\Theta(\pi_{0,0,0}, \pi_{N-2,N-2}, N-3) &= \Theta(\pi_{0,0,0}, \pi_{N-3,N-3}, N-1) = -1, \\ \Theta(\pi_{0,0,0}, \pi_{N-2,N-3}, N-2) &= +1.\end{aligned}$$

Now we calculate the values of the general theta foam. Suppose first that $i \leq k$. Then we have

$$(35) \quad \Theta(\pi_{p,q,r}, \pi_{j,k}, i) = \Theta(\pi_{p,q,r}, \pi_{i,i}\pi_{j-i,k-i}, i) = \Theta(\pi_{p+i,q+i,r+i}, \pi_{j-i,k-i}, 0),$$

by dot migration. In order to calculate $\Theta(\pi_{x,y,z}, \pi_{w,u}, 0)$ for $N-3 \geq x \geq y \geq z \geq 0$ and $N-2 \geq w \geq u \geq 0$, we use Lemma 3.1. By dot migration we have

$$(36) \quad \Theta(\pi_{x,y,z}, \pi_{w,u}, 0) = \sum_{(a,b,c) \sqsubset (x,y,z)} \Theta(\pi_{0,0,0}, \pi_{w,u}\pi_{a,b}, c).$$

Since $c \leq p \leq N-3$, a summand on the right-hand side of (36) can be nonzero only for $c = N-3$ and a and b such that $\pi_{N-2,N-2} \in \pi_{w,u}\pi_{a,b}$, ie $a = N-2-u$ and $b = N-2-w$. Hence the value of (36) is equal to -1 if

$$(37) \quad (N-2-u, N-2-w, N-3) \sqsubset (x, y, z),$$

and 0 otherwise. Finally, (37) is equivalent to $x + y + z + w + u = 3N - 7$, $x \geq N-2-u \geq y$, $y \geq N-2-w \geq z$ and $x \geq N-3 \geq z$, so we must have $u > 0$ and

$$\begin{aligned}x &= N-3, \\ y &= N-2-u, \\ z &= N-2-w.\end{aligned}$$

Going back to (36), we have that the value of theta is equal to 0 if $l = i$, and in the case $l > i$ it is nonzero (and equal to -1) if and only if

$$\begin{aligned}p &= N-3-i, \\ q &= N-2-k, \\ r &= N-2-j,\end{aligned}$$

which gives the first family.

Suppose now that $k < i$. As in (35) we have

$$(38) \quad \Theta(\pi_{p,q,r}, \pi_{j,k}, i) = \Theta(\pi_{p+k,q+k,r+k}, \pi_{j-k,0}, i-k).$$

Hence, we now concentrate on $\Theta(\pi_{x,y,z}, \pi_{w,0}, u)$ for $N - 3 \geq x \geq y \geq z \geq 0$, $N - 2 \geq w \geq 0$ and $N - 1 \geq u \geq 1$. Again, by using Lemma 3.1 we have

$$(39) \quad \Theta(\pi_{x,y,z}, \pi_{w,0}, 0) = \sum_{(a,b,c) \sqsubset (x,y,z)} \Theta(\pi_{0,0,0}, \pi_{w,0}\pi_{a,b}, u + c).$$

Since $a \leq N - 3$, we cannot have $\pi_{N-2,N-2} \in \pi_{w,0}\pi_{a,b}$ and we can have $\pi_{N-2,N-3} \in \pi_{w,0}\pi_{a,b}$ if and only if $a = N - 3$ and $b = N - 2 - w$. In this case we have a nonzero summand (equal to 1) if and only if $c = N - 2 - u$. Finally $\pi_{N-3,N-3} \in \pi_{w,0}\pi_{a,b}$ if and only if $a = N - 3$ and $b = N - 3 - w$. In this case we have a nonzero summand (equal to -1) if and only if $c = N - 1 - u$. Thus we have a summand on the right-hand side of (39) equal to $+1$ if and only if

$$(40) \quad (N - 3, N - 2 - w, N - 2 - u) \sqsubset (x, y, z),$$

and a summand equal to -1 if and only if

$$(41) \quad (N - 3, N - 3 - w, N - 1 - u) \sqsubset (x, y, z).$$

Note that in both above cases we must have $x + y + z + w + u = 3N - 7$, $x = N - 3$ and $u \geq 1$. Finally, the value of the r.h.s of (39) will be nonzero if and only if exactly one of (40) and (41) holds.

In order to find the value of the sum on right-hand side of (39), we split the rest of the proof in three cases according to the relation between w and u .

If $w \geq u$, (40) is equivalent to $y \geq N - 2 - w$, $z \leq N - 2 - w$, while (41) is equivalent to $y \geq N - 3 - w$, $z \leq N - 3 - w$ and $u \geq 2$. Now, we can see that the sum is nonzero and equal to 1 if and only if $z = N - 2 - w$ and so $y = N - 2 - u$. Returning to (38), we have that the value of $\Theta(\pi_{p,q,r}, \pi_{j,k}, i)$ is equal to 1 for

$$\begin{aligned} p &= N - 3 - k, \\ q &= N - 2 - i, \\ r &= N - 2 - j, \end{aligned}$$

for $N - 2 \geq j \geq i \geq k + 1 \geq 1$, which is our third family.

If $w \leq u - 2$, (40) is equivalent to $y \geq N - 2 - w$, $z \leq N - 2 - u$ and $u \leq N - 2$ while (41) is equivalent to $y \geq N - 3 - w$, $z \leq N - 1 - u$. Hence, in this case we have that the total sum is nonzero and equal to -1 if and only if $y = N - 3 - w$ and $z = N - 1 - u$, which by returning to (38) gives that the value of $\Theta(\pi_{p,q,r}, \pi_{j,k}, i)$ is

equal to -1 for

$$\begin{aligned} p &= N - 3 - k, \\ q &= N - 3 - j, \\ r &= N - 1 - i, \end{aligned}$$

for $N - 3 \geq i - 2 \geq j \geq k \geq 0$, which is our second family.

Finally, if $u = w + 1$ (40) becomes equivalent to $y \geq N - 2 - w$ and $z \leq N - 3 - w$, while (41) becomes $y \geq N - 3 - w$ and $z \leq N - 3 - w$. Thus, in order to have a nonzero sum, we must have $y = N - 3 - w$. But in that case, because of the fixed total sum of indices, we would have $z = N - 1 - u = N - 2 - w > N - 3 - w$, which contradicts (41). Hence, in this case, the total value of the theta foam is 0. \square

As a direct consequence of the previous theorem, we have:

Corollary 6.4 *For fixed values of j, k and i , if $j \neq i - 1$ and $k \neq i$, there is exactly one triple (p, q, r) such that the value of $\Theta(\pi_{p,q,r}, \pi_{j,k}, i)$ is nonzero. Also, if $j = i - 1$ or $k = i$, the value of $\Theta(\pi_{p,q,r}, \pi_{j,k}, i)$ is equal to 0 for every triple (p, q, r) . Hence, for fixed i , there are $\binom{n-1}{2}$ 5-tuples (p, q, r, j, k) such that $\Theta(\pi_{p,q,r}, \pi_{j,k}, i)$ is nonzero.*

Conversely, for fixed p, q and r , there always exist three different triples (j, k, i) (one from each family), such that $\Theta(\pi_{p,q,r}, \pi_{j,k}, i)$ is nonzero.

Finally, for all p, q, r, j, k and i , we have

$$\Theta(\pi_{p,q,r}, \pi_{j,k}, i) = \Theta(\hat{\pi}_{p,q,r}, \hat{\pi}_{j,k}, N - 1 - i).$$

The following relations are an immediate consequence of Lemma 6.3, Corollary 6.4, (CN_1) , (CN_2) and (CN_*) .

Corollary 6.5

$$(42) \quad \begin{array}{c} \text{Diagram of a theta foam } \pi_{p,q,r} \text{ with indices } p, q, r \text{ and } i \text{ marked.} \end{array} = \begin{cases} - \text{ (q,r) } & \text{if } p = N - 3 - i \\ - \text{ (p+1, q+1) } & \text{if } r = N - 1 - i \\ - \text{ (p+1, r) } & \text{if } q = N - 2 - i \\ 0 & \text{else.} \end{cases}$$

$$(43) \quad \begin{array}{c} \text{Diagram} \\ (k,m) \end{array} = \begin{cases} -\boxed{\begin{array}{c} * \\ (k-1, m-1, i) \end{array}} & \text{if } N-2 \geq k \geq m \geq i+1 \geq 1 \\ -\boxed{\begin{array}{c} * \\ (i-2, k, m) \end{array}} & \text{if } N-1 \geq i \geq k+2 \geq m+2 \geq 2 \\ \boxed{\begin{array}{c} * \\ (k-1, i-1, m) \end{array}} & \text{if } N-2 \geq k \geq i \geq m+1 \geq 1 \\ 0 & \text{else.} \end{cases}$$

$$(44) \quad \begin{array}{c} \text{Diagram} \\ (p,q,r) \\ (k,m) \end{array} = \begin{cases} -\boxed{p+2} & \text{if } q = N-2-m, r = N-2-k \\ -\boxed{r} & \text{if } p = N-3-m, q = N-3-k \\ \boxed{q+1} & \text{if } p = N-3-m, r = N-2-k \\ 0 & \text{else.} \end{cases}$$

Lemma 6.6

$$\begin{array}{c} \text{Diagram} \\ (i,j) \\ m \\ k \end{array} = \begin{cases} -1 & \text{if } m+j = N-1 = i+k+1 \\ +1 & \text{if } j+k = N-1 = i+m+1. \end{cases}$$

Proof By the dot conversion formulas, we get:

$$\begin{array}{c} \text{Diagram} \\ (i,j) \\ m \\ k \end{array} = \sum_{\alpha=0}^{i-j} \begin{array}{c} \text{Diagram} \\ m+i-\alpha \\ k+j+\alpha \end{array}$$

By (\ominus) we have

$$\begin{array}{c} \text{Diagram} \\ m+i-\alpha \\ k+j+\alpha \end{array} = \begin{cases} -1 & \text{if } m+i-(N-1) = \alpha = N-2-(k+j) \\ +1 & \text{if } m+i-(N-2) = \alpha = N-1-(k+j) \\ 0 & \text{else.} \end{cases}$$

We see that, in the sum above, the summands for two consecutive values of α will cancel unless one of them is zero and the other is not. We see that the total sum is equal to -1 if the first nonzero summand is at $\alpha = i - j$ and $+1$ if the last nonzero summand is at $\alpha = 0$. □

The following bubble identities are an immediate consequence of Lemma 6.6 and (CN_1) and (CN_2) .

Corollary 6.7

$$(45) \quad \begin{array}{c} i \\ \text{---} \\ \text{---} \\ \text{---} \\ j \end{array} = \begin{cases} -\overline{(i-1, j)} & \text{if } i > j \geq 0 \\ \overline{(j-1, i)} & \text{if } j > i \geq 0 \\ 0 & \text{if } i = j. \end{cases}$$

$$(46) \quad \begin{array}{c} i \\ \text{---} \\ \text{---} \\ \text{---} \\ (k, j) \end{array} = \begin{cases} -\overline{k+1} & \text{if } i + j = N - 1 \\ \overline{j} & \text{if } i + k = N - 2 \\ 0 & \text{else.} \end{cases}$$

The following identities follow easily from (CN_1) , (CN_2) , Lemma 6.3, Lemma 6.6 and their corollaries.

Corollary 6.8

$$(3C) \quad \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} + \sum_{a+b+c=N-2} \begin{array}{c} a \\ \text{---} \\ \text{---} \\ b \\ \text{---} \\ c \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}$$

$$(RD_1) \quad \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \bullet \\ \text{---} \end{array} \begin{array}{c} \text{---} \end{array} - \begin{array}{c} \text{---} \end{array} \begin{array}{c} \bullet \\ \text{---} \end{array}$$

$$(RD_2) \quad \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \bullet \\ \text{---} \end{array} \begin{array}{c} \text{---} \end{array} - \begin{array}{c} \bullet \\ \text{---} \end{array} \begin{array}{c} \text{---} \end{array} + \begin{array}{c} \text{---} \end{array} \begin{array}{c} \bullet \\ \text{---} \end{array}$$

$$(FC) \quad \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = - \begin{array}{c} \text{---} \end{array} \begin{array}{c} \text{---} \end{array}$$

Note that by the results above we are able to compute $\langle u \rangle_{KL}$ combinatorially, for any closed foam u whose singular graphs has no vertices, simply by using the cutting neck relations near all singular circles and evaluating the resulting spheres and theta foams. If the singular graph of u has vertices, then we do not know if our relations are sufficient to evaluate u . We conjecture that they are sufficient, and that therefore our theory is strictly combinatorial, but we do not have a complete proof.

Proposition 6.9 *The following identities hold in \mathbf{Foam}_N :*

The digon removal relations:

(DR₁)

(DR₂)

(DR_{3,1})

(DR_{3,2})

(DR_{3,3})

The first square removal relation:

(SqR₁)

Proof We first explain the idea of the proof. Naively one could try to consider all closures of the foams in a relation and compare their KL-evaluations. However, in practice we are not able to compute the KL-evaluations of arbitrary closed foams with singular vertices. Therefore we use a different strategy. We consider any foam in the proposition as a foam from \emptyset to its boundary, rather than as a foam between one part of its boundary to another part. If u is such a foam whose boundary is a closed

web Γ , then by the properties explained in Section 5 the KL–formula associates to u an element in $H^*(\Gamma)$, which is the homology of the complex associated to Γ in [11]. By Definition 6.1 and by the gluing properties of the KL–formula, as explained in Section 5, the induced linear map from $\langle \cdot |_{\text{KL}}: \mathbf{Foam}_N(\emptyset, \Gamma) \rightarrow H^*(\Gamma)$ is injective. The KL–formula also defines an inner product on $\mathbf{Foam}_N(\emptyset, \Gamma)$ by

$$(u, v) \mapsto \langle u\widehat{v} \rangle_{\text{KL}}.$$

By \widehat{v} we mean the foam in $\mathbf{Foam}_N(\Gamma, \emptyset)$ obtained by rotating v along the axis which corresponds to the y –axis (ie the horizontal axis parallel to the computer screen) in the original picture in this proposition. By the results in [11] we know the dimension of $H^*(\Gamma)$. Suppose it is equal to m and that we can find two sets of elements u_i and u_i^* in $\mathbf{Foam}_N(\emptyset, \Gamma)$, $i = 1, 2, \dots, m$, such that


$$\langle u_i\widehat{u_j^*} \rangle_{\text{KL}} = \delta_{i,j},$$

where $\delta_{i,j}$ is the Kronecker delta. Then $\{u_i\}$ and $\{u_i^*\}$ are mutually dual bases of $\mathbf{Foam}_N(\emptyset, \Gamma)$ and $\langle \cdot |_{\text{KL}}$ is an isomorphism. Therefore, two elements $f, g \in \mathbf{Foam}_N(\emptyset, \Gamma)$ are equal if and only if


$$\langle f\widehat{u_i} \rangle_{\text{KL}} = \langle g\widehat{u_i} \rangle_{\text{KL}},$$

for all $i = 1, 2, \dots, m$ (alternatively one can use the u_i^* of course). In practice this only helps if the left-hand side and the right-hand side of these m equations can be computed, eg if f, g and the u_i are all foams with singular graphs without vertices. Fortunately that is the case for all the relations in this proposition.

Let us now prove (DR₁) in detail. Note that the boundary of any of the foams in (DR₁), denoted Γ , is homeomorphic to the web:



Recall that the dimension of $H^*(\Gamma)$ is equal to $2N(N - 1)$ [11]. For $0 \leq i, j \leq 1$ and $0 \leq m \leq k \leq N - 2$, let $u_{i,j;(k,m)}$ denote the following foam:



Let
$$u_{i,j;(k,m)}^* = u_{1-j,1-i;(N-2-m,N-2-k)}.$$

From Equation (45) and the sphere relation (S_2) it easily follows that


$$\langle u_{i,j;(k,m)} \widehat{u_{r,s;(t,v)}^*} \rangle_{\text{KL}} = \delta_{i,r} \delta_{j,s} \delta_{k,t} \delta_{m,v},$$

where δ denotes the Kronecker delta. Note that there are exactly $2N(N-1)$ quadruples $(i, j; (k, l))$ which satisfy the conditions. Therefore the $u_{i,j;(k,m)}$ define a basis of $H^*(\Gamma)$ and the $u_{i,j;(k,m)}^*$ define its dual basis. In order to prove (DR_1) all we need to do next is check that


$$\langle (\text{left-hand side of } (\text{DR}_1)) \widehat{u_{i,j;(k,m)}} \rangle_{\text{KL}} = \langle (\text{right-hand side of } (\text{DR}_1)) \widehat{u_{i,j;(k,m)}} \rangle_{\text{KL}},$$

for all i, j and (k, m) . This again follows easily from equation (45) and the sphere relation (S_2).

The other digon removal relations are proved in the same way. We do not repeat the whole argument again for each digon removal relation, but will only give the relevant mutually dual bases. For (DR_2) , note that Γ is equal to:



Let $u_{i,k,m}$ denote the foam




for $0 \leq i, m \leq N-2$ and $0 \leq k \leq N-1$. The dual basis is defined by


$$u_{i,k,m}^* = -u_{N-2-m, N-1-k, N-2-i},$$

for the same range of indices. Note that there are $N(N-1)^2$ possible indices, which corresponds exactly to the dimension of $H^*(\Gamma)$.

For (DR_{3_1}) , the web Γ is equal to:



let $u_{i;(k,m);(p,q,r)}$ denote the foam




for $0 \leq i \leq 2$, $0 \leq m \leq k \leq 1$ and $0 \leq r \leq q \leq p \leq N - 3$. The dual basis is given by


$$u_{i;(k,m);(p,q,r)}^* = u_{2-(k+m);(1-\lfloor i/2 \rfloor, 1-\lceil i/2 \rceil);(N-3-r, N-3-q, N-3-p)},$$

for $0 \leq t \leq s \leq 1$, $0 \leq m \leq k \leq 1$ and $0 \leq r \leq q \leq p \leq N - 3$. Note that there are $3^2 \binom{N}{3}$ possible indices, which corresponds exactly to the dimension of $H^*(\Gamma)$.

For (DR_{3_2}) , take Γ to be:



Let $u_{i;(k,m);(s,t)}$ denote the foam




for $0 \leq i \leq N - 1$, $0 \leq m \leq k \leq N - 3$ and $0 \leq t \leq s \leq N - 3$. The dual basis is given by


$$u_{i;(k,m);(s,t)}^* = u_{N-1-i;(N-3-t, N-3-s);(N-3-m, N-3-k)},$$

for the same range of indices. Note that there are $N \binom{N-1}{2}^2$ indices, which corresponds exactly to the dimension of $H^*(\Gamma)$.

For (DR_{3_3}) , take Γ to be:



Let $u_{i,j;(k,m)}$ denote the foam




for $0 \leq i, j \leq N - 3$ and $0 \leq m \leq k \leq N - 2$. Define


$$u_{i,j;(k,m)}^* = u_{N-3-j, N-3-i;(N-2-m, N-2-k)},$$

for the same range of indices. Note that there are $(N - 2)^2 \binom{N}{2}$ indices, which corresponds exactly to the dimension of $H^*(\Gamma)$.

For (SqR_1) , the relevant web Γ is equal to:



By the results in [11], the dimension of $H^*(\Gamma)$ equals $N^2 + 2N(N - 2) + N^2(N - 2)^2$. The proof of this relation is very similar, except that it is slightly harder to find the mutually dual bases in $H^*(\Gamma)$. The problem is that the two terms on the right-hand side of (SqR_1) are topologically distinct. Therefore we get four different types of basis elements, which are gluings of the upper or lower half of the first term and the upper or lower half of the second term. For $0 \leq i, j \leq N - 1$, let $u_{i,j}$ denote the foam



with the top simple facet labelled by i and the bottom one by j . Take


$$u_{i,j}^* = u_{N-1-j, N-1-i}.$$

Note that


$$\langle u_{i,j} u_{k,m}^* \rangle_{\text{KL}} = \delta_{i,k} \delta_{j,m}$$

by the (FC) relation in Corollary 6.8 and the sphere relation (S_1) .

For $0 \leq i \leq N - 1$ and $0 \leq k \leq N - 3$, let $v'_{i,k}$ denote the foam



with the simple square on the right-hand side labelled by i and the other simple facet by k . Note that the latter is indeed only one facet, because it has a saddle-point in the middle where the dotted lines meet. For the same range of indices, we define $w'_{i,k}$ by



with the simple square on the left-hand side labelled by i and the other simple facet by k . The basis elements are now defined by

$$v_{i,k} = \sum_{a+b+c=N-3-k} v'_{c,a+b+i},$$

$$w_{i,j} = \sum_{a+b+c=N-3-j} w'_{c,a+b+i}.$$

The respective duals are defined by


$$v_{i,k}^* = w'_{k,N-1-i} \quad \text{and} \quad w_{i,k}^* = v'_{k,N-1-i}.$$

We show that

$$\langle v_{i,j} \widehat{v_{k,m}^*} \rangle_{\text{KL}} = \delta_{i,k} \delta_{j,m} = \langle v_{i,j} \widehat{v_{k,m}^*} \rangle_{\text{KL}}$$

holds. First apply the (FC) relation of Corollary 6.8. Then apply (RD₁) of the same corollary twice and finally use the sphere relation (S₁).

For $0 \leq i, j \leq N - 1$ and $0 \leq k, m \leq N - 3$, let $s'_{i,j,k,m}$ denote the foam



with the simple squares labelled by k and m , from left to right respectively, and the other two simple facets by i and j , from front to back respectively. The basis elements are defined by

$$s_{i,j,k,m} = \sum_{\substack{a+b+c=N-3-k \\ d+e+f=N-3-m}} s'_{c,f,i+a+d,j+b+e}.$$

For the same range of indices, the dual elements of this shape are given by

$$s_{i,j,k,m}^* = s'_{m,k,N-1-i,N-1-j}.$$

From (RD₁) of Corollary 6.8, applied twice, and the sphere relation (S₁), the following holds:

$$\langle s_{a,b,c,d} \widehat{s_{i,j,k,m}^*} \rangle_{\text{KL}} = \delta_{a,i} \delta_{b,j} \delta_{c,k} \delta_{d,m}.$$

It is also easy to see that the inner product $\langle \rangle_{\text{KL}}$ of a basis element and a dual basis element of distinct shapes, ie indicated by different letters above, gives zero. For example, consider

$$\langle u_{i,j} \widehat{v_{k,m}^*} \rangle_{\text{KL}}$$

for any valid choice of i, j, k, m . At the place where the two different shapes are glued,

$$u_{i,j} \widehat{v_{k,m}^*}$$


contains a simple saddle with a simple-double bubble. By Equation (46) that bubble kills

$$\langle u_{i,j} \widehat{v_{k,m}^*} \rangle_{\text{KL}},$$

because $m \leq N - 3$. The same argument holds for the other cases. This shows that $\{u, v, w, s\}$ and $\{u^*, v^*, w^*, s^*\}$ form dual bases of $H^*(\Gamma)$, because the number of possible indices equals $N^2 + 2N(N - 2) + N^2(N - 2)^2$.

In order to prove (SqR₁) one now has to compute the inner product of the left-hand side and the right-hand side with any basis element of $H^*(\Gamma)$ and show that they are equal. We leave this to the reader, since the arguments one has to use are the same as we used above. \square

Corollary 6.10 (The second square removal relation)



Proof Apply the relation (SqR₁) to the simple-double square tube perpendicular to the triple facet of the second term on the right-hand side of (SqR₂). The first term on the right-hand side of (SqR₁) yields minus the first term on the right-hand side of (SqR₂) after applying the relations (DR_{3₂}), (MP) and the bubble relation (42). The second term on the right-hand side of (SqR₁), ie the whole sum, yields the left-hand side of (SqR₂) after applying the relations (DR_{3₃}), (MP) and the bubble relation (42). Note that the signs come out right because in both cases we get two bubbles with opposite orientations. \square

7 Invariance under the Reidemeister moves


Let $\mathbf{Kom}(\mathbf{Foam}_N)$ and $\mathbf{Kom}_h(\mathbf{Foam}_N)$ denote the category of complexes in \mathbf{Foam}_N and the same category modulo homotopies respectively. As in [1] and [12] we can take all different flattenings of D to obtain an object in $\mathbf{Kom}(\mathbf{Foam}_N)$ which we call $\langle D \rangle$. The construction is well known by now and is indicated in Figure 7.

Theorem 7.1 The bracket $\langle \rangle$ is invariant in $\mathbf{Kom}_h(\mathbf{Foam}_N)$ under the Reidemeister moves.

$$\begin{aligned}
 \langle \times \rangle = 0 &\longrightarrow \underline{\langle \rangle} \{1 - N\} \xrightarrow{\text{foam}} \langle \times \rangle \{-N\} \longrightarrow 0 \\
 \langle \times \rangle = 0 &\longrightarrow \langle \times \rangle \{N\} \xrightarrow{\text{foam}} \underline{\langle \rangle} \{N - 1\} \longrightarrow 0
 \end{aligned}$$

Figure 7: Complex associated to a crossing. Underlined terms correspond to homological degree zero and $\{a\}$ is a positive shift by a in the q -degree.

Proof Reidemeister I Consider diagrams D and D' that differ in a circular region, as in the figure below.



We give the homotopy between complexes $\langle D \rangle$ and $\langle D' \rangle$ in Figure 8². By the sphere




Figure 8: Invariance under Reidemeister I


relation (S_1) , we get $g^0 f^0 = \text{Id}_{\langle D' \rangle^0}$. To see that $df^0 = 0$ holds, one can use dot mutation to get a new labelling of the same foam with the double facet labelled by $\pi_{N-1,0}$, which kills the foam by the dot conversion relations. The equality $dh = \text{Id}_{\langle D \rangle^1}$ follows from (DR_2) . To show that $f^0 g^0 + hd = \text{Id}_{\langle D \rangle^0}$, apply (RD_1) to hd and then cancel all terms which appear twice with opposite signs. What is left is the sum of

²We thank Christian Blanchet for spotting a mistake in a previous version of this diagram.

N terms which is equal to $\text{Id}_{\langle D \rangle^0}$ by (CN_1) . Therefore $\langle D' \rangle$ is homotopy-equivalent to $\langle D \rangle$.

For the Reidemeister I move with a positive crossing one can easily write down a similar homotopy equivalence.

Reidemeister IIa Consider diagrams D and D' that differ in a circular region, as in the figure below.



We only sketch the arguments that the diagram in Figure 9 defines a homotopy equivalence between the complexes $\langle D \rangle$ and $\langle D' \rangle$:





Figure 9: Invariance under Reidemeister IIa

- g and f are morphisms of complexes (use only isotopies);
- $g^1 f^1 = \text{Id}_{\langle D' \rangle^1}$ (uses Equation (45));
- $f^0 g^0 + hd = \text{Id}_{\langle D \rangle^0}$ and $f^2 g^2 + dh = \text{Id}_{\langle D \rangle^2}$ (use isotopies);
- $f^1 g^1 + dh + hd = \text{Id}_{\langle D \rangle^1}$ (use (DR_1)).

Reidemeister IIb Consider diagrams D and D' that differ only in a circular region, as in the figure below.



Again, we sketch the arguments that the diagram in Figure 10 defines a homotopy equivalence between the complexes $\langle D \rangle$ and $\langle D' \rangle$:





Figure 10: Invariance under Reidemeister IIb

- g and f are morphisms of complexes (use isotopies and (DR_2));
- $g^1 f^1 = Id_{\langle D' \rangle^1}$ (use (FC) and (S_1));
- $f^0 g^0 + hd = Id_{\langle D \rangle^0}$ and $f^2 g^2 + dh = Id_{\langle D \rangle^2}$ (use (RD_1) and (DR_2));
- $f^1 g^1 + dh + hd = Id_{\langle D \rangle^1}$ (use (DR_2) , (RD_1) , $(3C)$ and (SqR_1)).

Reidemeister III Consider diagrams D and D' that differ only in a circular region, as in the figure below.



In order to prove that $\langle D' \rangle$ is homotopy equivalent to $\langle D \rangle$ we show that the latter is homotopy equivalent to a third complex denoted $\langle Q \rangle$ in Figure 11. The differential in $\langle Q \rangle$ in homological degree 0 is defined by



for one summand and a similar foam for the other summand. By applying a symmetry relative to a horizontal axis crossing each diagram in $\langle D \rangle$ and $\langle Q \rangle$ we obtain a homotopy equivalence between $\langle D' \rangle$ and $\langle Q' \rangle$. It is easy to see that $\langle Q \rangle$ and $\langle Q' \rangle$ are isomorphic. In homological degree 0 the isomorphism is given by the obvious foam with two singular vertices. In the other degrees the isomorphism is given by the identity (in degrees 1 and 2 one has to swap the two terms of course). The fact that this defines an isomorphism follows immediately from the (MP) relation. We conclude that $\langle D \rangle$ and $\langle D' \rangle$ are homotopy equivalent. \square

The other versions of the Reidemeister III move can be obtained from the one above and Reidemeister II moves. The homotopy equivalences can then be defined as composites. For more details we refer the reader to Clark, Morrison and Walker [2].

From Theorem 7.1 we see that we can use any diagram D of L to obtain the invariant in $\mathbf{Kom}_h(\mathbf{Foam}_N)$ which justifies the notation $\langle L \rangle$ for $\langle D \rangle$.

8 Functoriality

The proof of the functoriality of our homology follows the same line of reasoning as in [1] and [12]. As in those papers, it is clear that the construction and the results of the previous sections can be extended to the category of tangles, following a similar approach using open webs and foams with corners. A foam with corners should be considered as living inside a cylinder, as in [1], such that the intersection with the cylinder is just a disjoint set of vertical edges.




Figure 11: Invariance under Reidemeister III. A circle attached to the tail of an arrow indicates that the corresponding morphism has a minus sign.

The degree formula can be extended to the category of open webs and foams with corners by:


Definition 8.1 Let u be a foam with d_{\bullet} dots of type \bullet , d_{\circ} dots of type \circ and d_{\odot} dots of type \odot . Let b_i be the number of vertical edges of type i of the boundary of u . The q -grading of u is given by

$$q(u) = - \sum_{i=1}^3 i(N-i)q_i(u) - 2(N-2)q_{s_v}(u) + \frac{1}{2} \sum_{i=1}^3 i(N-i)b_i + 2d_{\bullet} + 4d_{\circ} + 6d_{\odot}.$$

Note that the Kapustin–Li formula also induces a grading on foams with corners, because for any foam u between two (open) webs Γ_1 and Γ_2 , it gives an element in the graded vector space $\text{Ext}(M_1, M_2)$, where M_i is the matrix factorization associated to Γ_i in [11], for $i = 1, 2$. Recall that the Ext groups have a $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}$ -grading. For foams there is no $\mathbb{Z}/2\mathbb{Z}$ -grading, but the \mathbb{Z} -grading survives.

Lemma 8.2 For any foam u , the Kapustin–Li grading of u is equal to $q(u)$.

Proof Both gradings are additive under horizontal and vertical gluing and are preserved by the relations in \mathbf{Foam}_N . Also the degrees of the dots are the same in both gradings. Therefore it is enough to establish the equality between the gradings for the foams which generate \mathbf{Foam}_N . For any foam without a singular graph the gradings are obviously equal, so let us concentrate on the singular cups and caps, the singular saddle point cobordisms and the cobordisms with one singular vertex in Figure 4. To compute the degree of the singular cups and caps, for both gradings, one can use the digon removal relations. For example, let us consider the singular cup:



Any grading that preserves relation (DR_1) has to attribute the value of -1 to that foam, because the foam on the left-hand side of (DR_1) has degree 0, being an identity, and the dot on the right-hand side has degree 2. Similarly one can compute the degrees of the other singular cups and caps. To compute the degree of the singular saddle-point cobordisms, one can use the removing disc relations (RD_1) and (RD_2) . For example, the saddle-point cobordism in Figure 4 has to have degree 1. Finally, using the (MP) relation one shows that both foams on the right-hand side in Figure 4 have to have degree 0. □

Corollary 8.3 For any closed foam u we have that $\langle u \rangle_{\text{KL}}$ is zero if $q(u) \neq 0$.

As in [12] we have the following lemma, which is the analogue of [1, Lemma 8.6]:

Lemma 8.4 *For a crossingless tangle diagram T we have that $\text{Hom}_{\mathbf{Foam}_N}(T, T)$ is zero in negative degrees and \mathbb{Q} in degree zero.*

Proof Let T be a crossingless tangle diagram and $u \in \text{Hom}_{\mathbf{Foam}_N}(T, T)$. Recall that u can be considered to be in a cylinder with vertical edges intersecting the latter. The boundary of u consists of a disjoint union of circles (topologically speaking). By dragging these circles slightly into the interior of u one gets a disjoint union of circles in the interior of u . Apply relation (CN_1) to each of these circles. We get a linear combination of terms in $\text{Hom}_{\mathbf{Foam}_N}(T, T)$ each of which is the disjoint union of the identity on T , possibly decorated with dots, and a closed foam, which can be evaluated by $\langle \rangle_{\text{KL}}$. Note that the identity of T with any number of dots has always nonnegative degree. Therefore, if u has negative degree, the closed foams above have negative degree as well and evaluate to zero. This shows the first claim in the lemma. If u has degree 0, the only terms which survive after evaluating the closed foams have degree 0 as well and are therefore a multiple of the identity on T . This proves the second claim in the lemma. \square

The proofs of Lemmas 8.7–8.9 in [1] are “identical”. The proofs of the analogues of Theorem 4 and Theorem 5 in [1] follow the same reasoning but have to be adapted as in [12]. One has to use the homotopies of our Section 7 instead of the homotopies used in [1]. Without giving further details, we state the main result. Let $\mathbf{Kom}_{/\mathbb{Q}^*h}(\mathbf{Foam}_N)$ denote the category $\mathbf{Kom}_{/h}(\mathbf{Foam}_N)$ modded out by \mathbb{Q}^* , the invertible rational numbers.

Proposition 8.5 $\langle \rangle$ defines a functor $\mathbf{Link} \rightarrow \mathbf{Kom}_{/\mathbb{Q}^*h}(\mathbf{Foam}_N)$.

We do not know if the above projective functor can actually be considered as a functor to $\mathbf{Kom}_{/h}(\mathbf{Foam}_N)$. For certain applications, such as the $\mathfrak{sl}(N)$ -version of Rasmussen’s s -invariant or the coloured $\mathfrak{sl}(N)$ link homologies, one needs to have an affirmative answer to this question at least up to a sign.

9 The $\mathfrak{sl}(N)$ -link homology

Definition 9.1 Let Γ, Γ' be closed webs and $f \in \text{Hom}_{\mathbf{Foam}_N}(\Gamma, \Gamma')$. Define a functor \mathcal{F} between the categories \mathbf{Foam}_N and the category $\mathbf{Vect}_{\mathbb{Z}}$ of \mathbb{Z} -graded rational vector spaces and \mathbb{Z} -graded linear maps as:

- (1) $\mathcal{F}(\Gamma) = \text{Hom}_{\mathbf{Foam}_N}(\emptyset, \Gamma)$.
- (2) $\mathcal{F}(f)$ is the \mathbb{Q} -linear map $\mathcal{F}(f) : \text{Hom}_{\mathbf{Foam}_N}(\emptyset, \Gamma) \rightarrow \text{Hom}_{\mathbf{Foam}_N}(\emptyset, \Gamma')$ given by composition.

Note that \mathcal{F} is a tensor functor and that the degree of $\mathcal{F}(f)$ equals $q(f)$. Note also that $\mathcal{F}(\bigcirc) \cong H^*(\mathbb{C}P^{N-1})\{-N + 1\}$ and $\mathcal{F}(\bigodot) \cong H^*(\mathcal{G}_{2,N})\{-2N + 4\}$.

The following are a categorified version of the relations in Figure 3.

Lemma 9.2 (MOY decomposition) *We have the following decompositions under the functor \mathcal{F} :*

$$\begin{aligned}
 (1) \quad & \mathcal{F}\left(\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array}\right) \cong \mathcal{F}\left(\begin{array}{c} \parallel \\ \parallel \end{array}\right) \{-1\} \oplus \mathcal{F}\left(\begin{array}{c} \parallel \\ \parallel \end{array}\right) \{1\}. \\
 (2) \quad & \mathcal{F}\left(\begin{array}{c} \diagup \quad \diagdown \\ \diagup \quad \diagdown \end{array}\right) \cong \bigoplus_{i=0}^{N-2} \mathcal{F}\left(\begin{array}{c} \parallel \\ \parallel \end{array}\right) \{2 - N + 2i\}. \\
 (3) \quad & \mathcal{F}\left(\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array}\right) \cong \mathcal{F}\left(\begin{array}{c} \diagup \quad \diagdown \\ \diagup \quad \diagdown \end{array}\right) \oplus \left(\bigoplus_{i=0}^{N-3} \mathcal{F}\left(\begin{array}{c} \parallel \\ \parallel \end{array}\right) \left(\begin{array}{c} \parallel \\ \parallel \end{array}\right) \{3 - N + 2i\}\right). \\
 (4) \quad & \mathcal{F}\left(\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array}\right) \oplus \mathcal{F}\left(\begin{array}{c} \parallel \\ \parallel \end{array}\right) \cong \mathcal{F}\left(\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array}\right) \oplus \mathcal{F}\left(\begin{array}{c} \parallel \\ \parallel \end{array}\right).
 \end{aligned}$$

Proof (1) Define grading-preserving maps

$$\varphi_0: \mathcal{F}\left(\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array}\right) \{1\} \rightarrow \mathcal{F}\left(\begin{array}{c} \parallel \\ \parallel \end{array}\right) \qquad \varphi_1: \mathcal{F}\left(\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array}\right) \{-1\} \rightarrow \mathcal{F}\left(\begin{array}{c} \parallel \\ \parallel \end{array}\right)$$

$$\psi_0: \mathcal{F}\left(\begin{array}{c} \parallel \\ \parallel \end{array}\right) \rightarrow \mathcal{F}\left(\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array}\right) \{1\} \qquad \psi_1: \mathcal{F}\left(\begin{array}{c} \parallel \\ \parallel \end{array}\right) \rightarrow \mathcal{F}\left(\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array}\right) \{-1\}$$

as

$$\varphi_0 = \mathcal{F}\left(\begin{array}{c} \text{blue box} \\ \text{bubble} \end{array}\right), \qquad \varphi_1 = \mathcal{F}\left(\begin{array}{c} \text{blue box} \\ \text{bubble} \end{array}\right),$$

$$\psi_0 = \mathcal{F}\left(\begin{array}{c} \text{blue box} \\ \text{bubble} \end{array}\right), \qquad \psi_1 = -\mathcal{F}\left(\begin{array}{c} \text{blue box} \\ \text{bubble} \end{array}\right).$$

The bubble identities imply that $\varphi_i \psi_j = \delta_{i,j}$ (for $i, j = 0, 1$) and from the (DR₁) relation it follows that $\psi_0 \varphi_0 + \psi_1 \varphi_1$ is the identity map in

$$\mathcal{F}\left(\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array}\right).$$

(2) Define grading-preserving maps

$$\varphi_i: \mathcal{F}\left(\begin{array}{c} \diagup \quad \diagdown \\ \diagup \quad \diagdown \end{array}\right) \{N - 2 - 2i\} \rightarrow \mathcal{F}\left(\begin{array}{c} \parallel \\ \parallel \end{array}\right), \quad \psi_i: \mathcal{F}\left(\begin{array}{c} \parallel \\ \parallel \end{array}\right) \rightarrow \mathcal{F}\left(\begin{array}{c} \diagup \quad \diagdown \\ \diagup \quad \diagdown \end{array}\right) \{N - 2 - 2i\},$$

for $0 \leq i \leq N - 2$, as

$$\varphi_i = \mathcal{F} \left(\text{Diagram with blue cap and } N-2-j \right), \quad \psi_i = \sum_{j=0}^i \mathcal{F} \left(\text{Diagram with blue cap and } i-j \right).$$

We have $\varphi_i \psi_k = \delta_{i,k}$ and

$$\sum_{i=0}^{N-2} \psi_i \varphi_i = \text{Id} \left(\mathcal{F} \left(\text{Diagram with blue cap} \right) \right).$$

The first assertion is straightforward and can be checked using the (RD_1) and (S_1) relations. The second is immediate from the (DR_2) relation, which can be written as:

$$(DR_2) \quad \text{Diagram with blue vertical strip} = \sum_{i=0}^{N-2} \sum_{j=0}^i \text{Diagram with blue cap and } i-j \text{ and } N-2-j$$

(3) Define grading-preserving maps

$$\begin{aligned} \varphi_i &: \mathcal{F} \left(\text{Diagram with blue vertical strip} \right) \{N - 3 + 2i\} \rightarrow \mathcal{F} \left(\text{Diagram with blue cap} \right) \left(\right), \\ \psi_i &: \mathcal{F} \left(\text{Diagram with blue cap} \right) \left(\right) \rightarrow \mathcal{F} \left(\text{Diagram with blue vertical strip} \right) \{N - 3 + 2i\}, \end{aligned}$$

for $0 \leq i \leq N - 3$, and

$$\rho: \mathcal{F} \left(\text{Diagram with blue vertical strip} \right) \rightarrow \mathcal{F} \left(\text{Diagram with blue cap} \right), \quad \tau: \mathcal{F} \left(\text{Diagram with blue cap} \right) \rightarrow \mathcal{F} \left(\text{Diagram with blue vertical strip} \right),$$

$$\text{as } \varphi_i = \mathcal{F} \left(\text{Diagram with blue cap and } i \right), \quad \psi_i = \sum_{a+b+c=N-3-i} \mathcal{F} \left(\text{Diagram with blue cap and } a, b, c \right),$$

$$\rho = \mathcal{F} \left(\text{Diagram with blue vertical strip} \right), \quad \tau = -\mathcal{F} \left(\text{Diagram with blue vertical strip} \right).$$

Checking that $\varphi_i \psi_k = \delta_{i,k}$ for $0 \leq i, k \leq N-3$, $\varphi_i \tau = 0$ and $\rho \psi_i = 0$, for $0 \leq i \leq N-3$, and $\rho \tau = -1$ is left to the reader. From the (SqR_1) relation it follows that

$$\tau \rho + \sum_{i=0}^{N-3} \psi_i \varphi_i = \text{Id} \left(\mathcal{F} \left(\square \square \right) \right).$$

Direct sum decomposition (4) We prove direct decomposition (4) showing that

$$(a) \quad \mathcal{F} \left(\begin{array}{c} \uparrow \uparrow \\ \downarrow \downarrow \end{array} \right) \cong \mathcal{F} \left(\begin{array}{c} \uparrow \uparrow \\ \downarrow \downarrow \end{array} \right) \oplus \mathcal{F} \left(\begin{array}{c} \uparrow \downarrow \\ \downarrow \uparrow \end{array} \right), \quad (b) \quad \mathcal{F} \left(\begin{array}{c} \uparrow \uparrow \\ \downarrow \downarrow \end{array} \right) \cong \mathcal{F} \left(\begin{array}{c} \uparrow \downarrow \\ \downarrow \uparrow \end{array} \right) \oplus \mathcal{F} \left(\begin{array}{c} \uparrow \downarrow \\ \downarrow \uparrow \end{array} \right).$$

Note that this suffices because the last term on the right-hand side of (a) is isomorphic to the last term on the right-hand side of (b) by the (MP) relation.

To prove (a) we define grading-preserving maps

$$\begin{aligned} \varphi_0: \mathcal{F} \left(\begin{array}{c} \uparrow \uparrow \\ \downarrow \downarrow \end{array} \right) &\rightarrow \mathcal{F} \left(\begin{array}{c} \uparrow \uparrow \\ \downarrow \downarrow \end{array} \right), & \varphi_1: \mathcal{F} \left(\begin{array}{c} \uparrow \uparrow \\ \downarrow \downarrow \end{array} \right) &\rightarrow \mathcal{F} \left(\begin{array}{c} \uparrow \downarrow \\ \downarrow \uparrow \end{array} \right), \\ \psi_0: \mathcal{F} \left(\begin{array}{c} \uparrow \uparrow \\ \downarrow \downarrow \end{array} \right) &\rightarrow \mathcal{F} \left(\begin{array}{c} \uparrow \downarrow \\ \downarrow \uparrow \end{array} \right), & \psi_1: \mathcal{F} \left(\begin{array}{c} \uparrow \downarrow \\ \downarrow \uparrow \end{array} \right) &\rightarrow \mathcal{F} \left(\begin{array}{c} \uparrow \downarrow \\ \downarrow \uparrow \end{array} \right), \end{aligned}$$

$$\begin{aligned} \text{by } \varphi_0 &= -\mathcal{F} \left(\begin{array}{c} \text{foam diagram} \end{array} \right), & \varphi_1 &= -\mathcal{F} \left(\begin{array}{c} \text{foam diagram with *} \end{array} \right), \\ \psi_0 &= \mathcal{F} \left(\begin{array}{c} \text{foam diagram} \end{array} \right), & \psi_1 &= \mathcal{F} \left(\begin{array}{c} \text{foam diagram with *} \end{array} \right). \end{aligned}$$

We have that $\varphi_i \psi_j = \delta_{i,j}$ for $i, j = 0, 1$ (we leave the details to the reader). From the (SqR_2) relation it follows that

$$\psi_0 \varphi_0 + \psi_1 \varphi_1 = \text{Id} \left(\mathcal{F} \left(\begin{array}{c} \uparrow \uparrow \\ \downarrow \downarrow \end{array} \right) \right).$$

Applying a symmetry to all diagrams in decomposition (a) gives decomposition (b). \square

In order to relate our construction to the $\mathfrak{sl}(N)$ polynomial we need to introduce shifts. We denote by $\{n\}$ an upward shift in the q -grading by n and by $[m]$ an upward shift in the homological grading by m .

Definition 9.3 Let $\langle L \rangle_i$ denote the i -th homological degree of the complex $\langle L \rangle$. We define the i -th homological degree of the complex $\mathcal{F}(L)$ to be

$$\mathcal{F}_i(L) = \mathcal{F}\langle L \rangle_i[-n_-]\{(N-1)n_+ - Nn_- + i\},$$

where n_+ and n_- denote the number of positive and negative crossings in the diagram used to calculate $\langle L \rangle$.

We now have a homology functor $\mathbf{Link} \rightarrow \mathbf{Vect}_{\mathbb{Z}/\mathbb{Q}^*}$ which we still call \mathcal{F} . Definition 9.3, Theorem 7.1 and Lemma 9.2 imply:


Theorem 9.4 For a link L the graded Euler characteristic of $H^*(\mathcal{F}(L))$ equals $P_N(L)$, the $\mathfrak{sl}(N)$ polynomial of L .

The MOY-relations are also the last bit that we need in order to show the following theorem.


Theorem 9.5 For any link L , the bigraded complex $\mathcal{F}(L)$ is isomorphic to the Khovanov–Rozansky complex $\mathbf{KR}(L)$ in [11].

Proof The map $\langle |_{\mathbf{KL}}$ defines a grading preserving linear injection $\mathcal{F}(\Gamma) \rightarrow \mathbf{KR}(\Gamma)$, for any web Γ . Lemma 9.2 implies that the graded dimensions of $\mathcal{F}(\Gamma)$ and $\mathbf{KR}(\Gamma)$ are equal, so $\langle |_{\mathbf{KL}}$ is a grading preserving linear isomorphism, for any web Γ .

To prove the theorem we would have to show that $\langle |_{\mathbf{KL}}$ commutes with the differentials. We call



the *zip* and



the *unzip*. Note that both the zip and the unzip have q -degree 1. Let Γ_1 be the source web of the zip and Γ_2 its target web, and let Γ be the theta web, which is the total boundary of the zip where the vertical edges have been pinched. The q -dimension of $\text{Ext}(\Gamma_1, \Gamma_2)$ is equal to

$$q^{2N-2} \text{qdim}(\Gamma) = q + q^2(\dots),$$

where (\dots) is a polynomial in q . Therefore the differentials in the two complexes commute up to a scalar. By the removing disc relation (RD_1) we see that if the “zips” commute up to λ , then the “unzips” commute up to λ^{-1} . If $\lambda \neq 1$, we have to modify our map between the two complexes slightly, in order to get an honest morphism of complexes. We use Khovanov’s idea of “twist equivalence” in [8]. For a given link consider the hypercube of resolutions. If an arrow in the hypercube corresponds to a zip, multiply $\langle(\text{target})|_{\text{KL}}$ by λ , where target means the target of the arrow. If it corresponds to an unzip, multiply $\langle(\text{target})|_{\text{KL}}$ by λ^{-1} . This is well-defined, because all squares in the hypercube (anti-)commute. By definition this new map commutes with the differentials and therefore proves that the two complexes are isomorphic. \square

We conjecture that the above isomorphism actually extends to link cobordisms, giving a projective natural isomorphism between the two projective link homology functors. Proving this would require a careful comparison between the two functors for all the elementary link cobordisms.

Acknowledgements The authors thank Mikhail Khovanov for helpful comments and Lev Rozansky for patiently explaining the Kapustin–Li formula in detail to us.

The authors were supported by the Fundação para a Ciência e a Tecnologia (ISR/IST plurianual funding) through the programme “Programa Operacional Ciência, Tecnologia, Inovação” (POCTI) and the POS Conhecimento programme, cofinanced by the European Community fund FEDER. Marko Stošić was also partially supported by the Ministry of Science of Serbia, project 144032.

References

- [1] **D Bar-Natan**, *Khovanov’s homology for tangles and cobordisms*, *Geom. Topol.* 9 (2005) 1443–1499 MR2174270
- [2] **D Clark, S Morrison, K Walker**, *Fixing the functoriality of Khovanov homology* arXiv:math/0701339
- [3] **W Fulton**, *Young tableaux. With applications to representation theory and geometry*, London Math. Soc. Student Texts 35, Cambridge Univ. Press (1997) MR1464693
- [4] **W Fulton, J Harris**, *Representation theory. A first course*, Graduate Texts in Math. 129, Springer, New York (1991) MR1153249
- [5] **H Hiller**, *Geometry of Coxeter groups*, Research Notes in Math. 54, Pitman (Advanced Publishing Program), Boston (1982) MR649068
- [6] **A Kapustin, Y Li**, *Topological correlators in Landau–Ginzburg models with boundaries*, *Adv. Theor. Math. Phys.* 7 (2003) 727–749 MR2039036

- [7] **M Khovanov**, *sl(3) link homology*, *Algebr. Geom. Topol.* 4 (2004) 1045–1081 MR2100691
- [8] **M Khovanov**, *Link homology and Frobenius extensions*, *Fund. Math.* 190 (2006) 179–190 MR2232858
- [9] **M Khovanov, L Rozansky**, *Topological Landau–Ginzburg models on the world-sheet foam*, *Adv. Theor. Math. Phys.* 11 (2007) 233–259 MR2322554
- [10] **M Khovanov, L Rozansky**, *Virtual crossing, convolutions and a categorification of the SO(2N) Kauffman polynomial*, *J. Gökova Geom. Topol. GGT* 1 (2007) 116–214 MR2386537
- [11] **M Khovanov, L Rozansky**, *Matrix factorizations and link homology*, *Fund. Math.* 199 (2008) 1–91 MR2391017
- [12] **M Mackaay, P Vaz**, *The universal sl₃–link homology*, *Algebr. Geom. Topol.* 7 (2007) 1135–1169 MR2336253
- [13] **S Morrison**, *A diagrammatic category for the representation theory of U_q(sl(N))*, PhD thesis, UC Berkeley (2007) arXiv:0704.1503
- [14] **H Murakami, T Ohtsuki, S Yamada**, *Homfly polynomial via an invariant of colored plane graphs*, *Enseign. Math. (2)* 44 (1998) 325–360 MR1659228
- [15] **L Rozansky**, *Topological A–models on seamed Riemann surfaces*, *Adv. Theor. Math. Phys.* 11 (2007) 517–529 MR2354074

MM, PV: Departamento de Matemática, Universidade do Algarve
Campus de Gambelas, 8005-139 Faro, Portugal

MS: Instituto de Sistemas e Robótica and CAMGSD, Instituto Superior Técnico
Avenida Rovisco Pais, 1049-001 Lisboa, Portugal

mmackaay@ualg.pt, mstosic@math.ist.utl.pt, pfortevaz@ualg.pt

Proposed: Peter Ozsváth

Received: 4 December 2007

Seconded: Peter Teichner, Robion Kirby

Revised: 1 August 2008