

## CAN AN ANISOTROPIC REDUCTIVE GROUP ADMIT A TITS SYSTEM?

PIERRE-EMMANUEL CAPRACE\* AND TIMOTHÉE MARQUIS

ABSTRACT. Seeking for a converse to a well-known theorem by Borel–Tits, we address the question whether the group of rational points  $G(k)$  of an anisotropic reductive  $k$ -group may admit a split spherical BN-pair. We show that if  $k$  is a perfect field or a local field, then such a BN-pair must be virtually trivial. We also consider arbitrary compact groups and show that the only abstract BN-pairs they can admit are spherical, and even virtually trivial provided they are split.

## 1. INTRODUCTION

In a seminal paper [5], Armand Borel and Jacques Tits established — amongst other things — that the group  $G(k)$  of  $k$ -rational points of a (connected) reductive linear algebraic  $k$ -group  $G$  always possesses a canonical BN-pair, where  $k$  is an arbitrary ground field. More precisely, they showed that if  $P$  is a minimal parabolic  $k$ -subgroup of  $G$ , and if  $N$  is the normalizer in  $G$  of some maximal  $k$ -split torus contained in  $P$ , then  $(P(k), N(k))$  is a BN-pair for  $G(k)$ . This result constitutes a cornerstone in understanding the abstract group structure of the group of  $k$ -rational points  $G(k)$ . As an application, it yields for example the celebrated simplicity result of Tits [20]. Of course, the aforementioned BN-pair is trivial when  $G$  is **anisotropic** over  $k$ . (Abusing slightly the standard conventions, we shall say that  $G$  is anisotropic if it has no proper  $k$ -parabolic subgroup, *i.e.* if  $P = G$ . As is well-known, this definition coincides with the standard one in case  $G$  is semi-simple (see [4, 11.21])). In fact, the abstract group structure of  $G(k)$  remains intriguing and mysterious to a large extent in the anisotropic case. In this context, we propose the following.

**Conjecture** (Converse to Borel–Tits). *Let  $G$  be a reductive algebraic  $k$ -group which is anisotropic over  $k$ . Then every split spherical BN-pair for  $G(k)$  is trivial.*

Recall that a BN-pair  $(B, N)$  for a group  $G$  is called **spherical** if the associated Weyl group  $W := N/T$  is finite, where  $T := B \cap N$ . It is said to be **split** if it is saturated (*i.e.*  $T = \bigcap_{w \in W} wBw^{-1}$ ), and if there exists a nilpotent normal subgroup  $U \triangleleft B$  such that  $B \cong U \rtimes T$ . Note that if  $(B, N)$  is irreducible of rank at least 2, one can show that  $U$  is automatically nilpotent (see [19]). The BN-pair for  $G(k)$  described above is always split in the above sense ([4, 14.19]).

Besides the natural search for a converse to Borel–Tits, a motivation to consider the above conjecture is provided by the recent work of Peter Abramenko and Ken Brown [1], who constructed Weyl transitive actions on trees for certain anisotropic groups over global function fields. We refer to [2, Ch. 6] for more details on the relations and distinctions between BN-pairs, strong transitivity and Weyl transitivity.

Our first contribution concerns the special case when the ground field  $k$  is a local field. The  $k$ -anisotropy of  $G$  is then equivalent to the compactness of  $G(k)$  (see [13]). In fact, our first step will be to establish the following two results, which concern arbitrary compact topological groups (not necessarily associated with algebraic groups).

**Theorem 1.** *Let  $G$  be a compact group. Then every BN-pair for  $G$  is spherical.*

**Theorem 2.** *Let  $G$  be a compact group possessing a split spherical BN-pair  $(B, N)$ . Then, the associated building is finite. In other words,  $[G : B] < \infty$ .*

---

Date: August 2009.

2000 *Mathematics Subject Classification.* 20E42, 20G15 (Primary); 22C05 (Secondary).

\*Supported by the Fund for Scientific Research–F.N.R.S., Belgium.

We emphasize that the BN-pairs appearing in these statements are *abstract*: The corresponding subgroups  $B$  and  $N$  are *not* supposed to be closed in  $G$ . Specializing to anisotropic groups over local fields, we deduce the following immediate corollary.

**Theorem 3.** *Let  $k$  be a local field and  $G$  be a connected semi-simple algebraic  $k$ -group which is anisotropic over  $k$ . Then:*

- (1) *Every BN-pair for  $G(k)$  is spherical.*
- (2) *Every split spherical BN-pair  $(B, N)$  for  $G(k)$  is ‘virtually trivial’, in the sense that  $B$  has finite index in  $G(k)$ .*

Finally, we consider the case of perfect ground fields.

**Theorem 4.** *Let  $k$  be a perfect field and  $G$  be a reductive algebraic  $k$ -group which is anisotropic over  $k$ . Then every split spherical BN-pair for  $G(k)$  is virtually trivial.*

Notice that Theorems 3 and 4 are logically independent, since there exist local fields which are not perfect and vice-versa.

It would be very interesting to sharpen the conclusion of Theorems 3 and 4, that is, to show that the BN-pair must be trivial, and not only virtually trivial. However, we expect this to be quite difficult, since it is closely related to a conjecture due to Andrei Rapinchuk and Gopal Prasad (see [14]), which may be stated as follows: “*Let  $G$  be a reductive  $k$ -group which is anisotropic over  $k$ . Then, every finite quotient of  $G(k)$  is solvable.*” As of today, this conjecture was confirmed only when  $G$  is the multiplicative group of a finite dimensional division algebra (see [15]). We now sketch informally how these two problems are related.

On one side, if  $G(k)$  possesses a BN-pair with finite associated building  $\Delta$ , and if  $K := \ker(G(k) \curvearrowright \Delta)$  is the kernel of the corresponding action, then  $G(k)/K$  is a finite group whose action on  $\Delta$  is faithful, and thus  $G(k)/K$  possesses a faithful BN-pair. But these groups have been classified: they are simple Chevalley groups, and in particular are not solvable (up to two exceptions). Thus, if the BN-pair for  $G(k)$  were nontrivial, there would exist (modulo the two exceptions) a non-solvable finite quotient of  $G(k)$ .

Conversely, suppose that  $G(k)$  possesses a nontrivial and non-solvable finite quotient  $F' := G(k)/K$ . Let  $R \triangleleft F'$  be the solvable radical of  $F'$ , that is, its largest solvable normal subgroup. Going to the quotient  $F := F'/R$ , we thus know that  $G(k)$  surjects onto a nontrivial finite group with trivial solvable radical (namely,  $F$ ). Let now  $M$  be a minimal normal subgroup of  $F$ . Then  $M$  is a direct product of non-Abelian simple groups which are pairwise isomorphic, say  $M \cong S_1 \times \cdots \times S_k$  with  $S_i \cong S$  for all  $i \in \{1, \dots, k\}$ . By the classification of finite simple groups,  $S$  is very likely to be a Chevalley group. Such a group possesses a root datum, and thus also a nontrivial BN-pair whose associated (finite) building is in bijection with  $S/B$ . Repeating this construction for each  $S_i$ , we then get a finite building  $\Delta = \Delta_1 \times \cdots \times \Delta_k$  on which  $M = S_1 \times \cdots \times S_k$  acts strongly transitively. Finally, the action of  $\text{Aut}(M)$  on the set of  $p$ -Sylow subgroups of  $M$  (where  $p = \text{char } k$ ) induces an action of  $\text{Aut}(M)$  on  $\Delta$  making the diagram

$$\begin{array}{ccc} F & \xrightarrow{\alpha} & \text{Aut}(M) \\ \iota \uparrow & & \downarrow \\ M & \xrightarrow{\text{strongly tr.}} & \text{Aut}(\Delta) \end{array}$$

commutative, where  $\alpha(f)$  denotes the conjugation by  $f$  for all  $f \in F$ . In particular, we get a strongly transitive action of  $F$ , and thus also of  $G(k)$ , on the finite building  $\Delta$ . This yields a nontrivial and virtually trivial BN-pair for  $G(k)$ .

**General conventions.** All algebraic groups considered here are supposed to be affine, all topological groups are assumed Hausdorff and all BN-pairs have finite rank.

**Acknowledgement.** We are very grateful to the anonymous referee for his/her useful detailed comments.

## 2. PROOF OF THEOREM 1

**2.1. Heuristic sketch.** Let  $G$  be a compact group and let  $(B, N)$  be a BN-pair for  $G$ . Also, let  $\Delta$  be the associated building. We consider the Davis realization of  $\Delta$ , noted  $|\Delta|_{\text{CAT}(0)}$  in this paper, and which is a complete CAT(0) space, as well as a simplicial complex, on which  $G$  acts by simplicial isometries. The key step in the proof of Theorem 1 is then to establish that this action is elliptic (Theorem 2.5 below). To do so, we use a result of Martin Bridson stating that such an action is always semi-simple, and we then argue by contradiction, assuming that  $G$  possesses an element with no fixed point. Such an element would then generate a subgroup  $Q$  of  $G$  which acts by translations on  $|\Delta|_{\text{CAT}(0)}$ . Moreover, the structure of simplicial complex of  $|\Delta|_{\text{CAT}(0)}$  implies that the set of translation lengths of the elements of  $Q$  is discrete at 0. The contradiction now comes from divisibility properties of compact and procyclic groups, which we apply to  $Q$ .

**2.2. Procyclic groups.** Let  $G$  be a profinite group. Recall that  $G$  is said to be **procyclic** if there exists a  $g \in G$  such that the subgroup generated by  $g$  is dense in  $G$ , that is,  $G = \overline{\langle g \rangle}$ . Moreover  $G$  is said to be **pro- $p$**  for some prime  $p$  if every finite Hausdorff quotient of  $G$  is a  $p$ -group.

The following basic properties of procyclic groups can be found in [16, 2.7]. The symbol  $\mathbb{P}$  denotes the set of all primes.

**Proposition 2.1.** *Let  $G$  be a procyclic group. Then,*

- (i)  $G$  is the direct product  $G = \prod_{p \in \mathbb{P}} G_p$  of its  $p$ -Sylow subgroups, and each  $G_p$  is a pro- $p$  procyclic group.
- (ii)  $G$  is, in a unique way, a quotient of  $\hat{\mathbb{Z}} := \prod_{p \in \mathbb{P}} \mathbb{Z}_p$ . If  $G$  is pro- $p$  for some  $p \in \mathbb{P}$ , then it is a quotient of  $\mathbb{Z}_p$ .

**2.3. Divisible groups.** Recall that an element  $g \in G$  is said to be  **$n$ -divisible** for some  $n \in \mathbb{N}$  if there exists an  $h \in G$  such that  $h^n = g$ . We say that  $g$  is **divisible** if it is  $n$ -divisible for each  $n \geq 1$ . The group  $G$  is called  **$n$ -divisible** (respectively **divisible**) when all its elements are.

Now, every prime  $q$  different from  $p$  is invertible in  $\mathbb{Z}_p$  since its  $p$ -adic valuation is zero. Hence, the additive group  $\mathbb{Z}_p$  is  $q$ -divisible for each  $q \in \mathbb{P} \setminus \{p\}$ . In particular, Proposition 2.1 shows that if a procyclic group  $G$  has trivial  $q$ -Sylow subgroups, then  $G$  is  $q$ -divisible.

We conclude this paragraph by stating the following characterization of divisibility for compact groups (see [12, Corollaire 2]).

**Proposition 2.2.** *Let  $G$  be a compact topological group. Then,  $G$  is divisible if and only if it is connected.*

**2.4. Semi-simple actions on CAT(0) spaces.** Let  $G$  be a group acting on a metric space  $(X, d)$ . For every  $g \in G$ , we define the **translation length** of  $g$  by  $|g| := \inf\{d(x, g \cdot x) \mid x \in X\} \in [0, \infty)$  and the **minimal set** of  $g$  by  $\text{Min}(g) := \{x \in X \mid d(x, g \cdot x) = |g|\}$ . An element  $g \in G$  is said to be **semi-simple** when  $\text{Min}(g)$  is nonempty. In that case, we say that  $g$  is **elliptic** if it fixes some point, that is, if  $|g| = 0$ ; otherwise, if  $|g| > 0$ , we call  $g$  **hyperbolic**.

A **geodesic line** (respectively, **geodesic segment**) in  $X$  is an isometry  $f: \mathbb{R} \rightarrow X$  (respectively,  $f: [0; 1] \rightarrow X$ ); by abuse of language, we will identify  $f$  with its image in  $X$ .

The following lemma follows from Proposition 2.4 in [6].

**Lemma 2.3.** *Let  $(X, d)$  be a complete CAT(0) metric space, and let  $C$  be a closed convex nonempty subset of  $X$ . Then:*

- (i) For every  $x \in X$ , there is a unique  $y \in C$  such that  $d(x, y) = d(x, C)$ , where  $d(x, C) := \inf_{z \in C} d(x, z)$ . We call  $y$  the **projection** of  $x$  on  $C$  and we write  $y = \text{proj}_C x$ .
- (ii) For all  $x_1, x_2 \in X$ , we have  $d(\text{proj}_C x_1, \text{proj}_C x_2) \leq d(x_1, x_2)$ .

Suppose now that  $(X, d)$  is a cell complex. We then say that  $G$  acts by **cellular isometries** on  $X$  if it preserves the metric, as well as the cell decomposition of  $X$ .

The following result is due to Martin Bridson [7].

**Proposition 2.4.** *Let  $X$  be a locally Euclidean CAT(0) cell complex with finitely many isometry types of cells, and  $G$  be a group acting on  $X$  by cellular isometries. Then every element of  $G$  is semi-simple. Moreover,  $\inf\{|g| \neq 0 \mid g \in G\} > 0$ .*

We now establish the following result, which is the key ingredient for the proof of Theorem 1:

**Theorem 2.5.** *Let  $X$  be a locally Euclidean CAT(0) cell complex with finitely many isometry types of cells, and  $G$  be a compact group acting on  $X$  by cellular isometries (not necessarily continuously). Then every element of  $G$  is elliptic.*

*Proof.* Suppose for a contradiction there exists a  $g \in G$  without fixed point. Proposition 2.4 then implies that  $g$  is hyperbolic. Let  $Q = \overline{\langle g \rangle}$  be the closure of the subgroup generated by  $g$  in  $G$ . So,  $Q$  is compact.

Claim 1:  $Q$  is Abelian.

This is clear since it contains a dense Abelian (in fact cyclic) subgroup.

Claim 2: For every  $h \in Q$ , the minimal set  $\text{Min}(h)$  is a closed convex subset of  $X$  which is stabilized by  $Q$ .

This follows from [6, Proposition II.6.2].

Claim 3: For every  $h \in Q$  and every nonempty closed convex subset  $C$  of  $X$  stabilized by  $Q$ , the set  $C \cap \text{Min}(h)$  is nonempty.

Note first that  $\text{Min}(h)$  is nonempty by Proposition 2.4. Let  $x \in \text{Min}(h)$  and consider the projections  $y := \text{proj}_C x$  and  $z := \text{proj}_C hx$  provided by Lemma 2.3. Since  $hC = C$ , we then obtain

$$d(x, y) = \inf_{c \in C} d(x, c) = \inf_{c \in C} d(hx, hc) = \inf_{c \in C} d(hx, c) = d(hx, z).$$

Hence  $d(hx, hy) = d(x, y) = d(hx, z)$ , and so  $z = hy = \text{proj}_C hx$  by uniqueness of projections. Since in addition  $d(y, z) \leq d(x, hx) = |h|$  by Lemma 2.3, we finally get  $d(y, hy) = |h|$  and therefore  $y \in C \cap \text{Min}(h)$ .

Claim 4: For all  $h_1, h_2 \in Q$ , the set  $\text{Min}(h_1) \cap \text{Min}(h_2)$  is nonempty.

As  $\text{Min}(h_1)$  and  $\text{Min}(h_2)$  are nonempty by Proposition 2.4, the claim follows from Claims 2 and 3.

Claim 5: Let  $h \in Q$  and let  $C$  be a nonempty closed convex subset of  $X$  stabilized by  $Q$ . We may thus consider the action of  $h$  on  $C$ . Denote by  $|h|_C$  the translation length of  $h$  for this action. Then,  $h$  is semi-simple in  $C$  and  $|h| = |h|_C$ .

Claim 3 yields that if  $x \in \text{Min}(h)$ , then  $y := \text{proj}_C x \in \text{Min}(h)$ . Since  $\text{Min}(h)$  is nonempty by Proposition 2.4, the claim follows.

Claim 6: For every  $h \in Q$  and  $n \geq 1$ , we have  $|h^n| = n|h|$ .

By Claim 4, we may choose an  $x \in \text{Min}(h) \cap \text{Min}(h^n)$ . Note that  $h$  is elliptic (respectively hyperbolic) if and only if  $h^n$  is so (see [6, II.6.7 and II.6.8]). In particular, if  $h$  is hyperbolic, then  $x$  belongs to some  $h$ -axis, which is also an  $h^n$ -axis. In any case, we obtain  $d(x, h^n x) = nd(x, hx)$ , whence  $|h^n| = d(x, h^n x) = nd(x, hx) = n|h|$ .

Claim 7: Every divisible element of  $Q$  is elliptic.

Let  $h \in Q$  be divisible and suppose for a contradiction it is not elliptic. Then  $h$  is hyperbolic by Proposition 2.4. For each natural number  $n \geq 1$ , choose an  $h_n \in Q$  such that  $h_n^n = h$ . In particular, all  $h_n$  are hyperbolic. Moreover,  $|h_n^n| = n|h_n|$  by Claim 6. Therefore, we obtain a sequence  $(h_n)$  of elements of  $Q$  such that  $|h_n| = |h|/n > 0$ , contradicting the second part of Proposition 2.4.

We now establish the desired contradiction to the hyperbolicity of  $g$ . First note that the component group  $P := Q/Q^0$  of  $Q$  is a profinite group. In fact, it is even procyclic, since the subgroup generated by the projection of  $g$  in  $P$  is dense in  $P$ , the natural mapping  $\pi: Q \rightarrow Q/Q^0$  being continuous. In particular, it follows from Proposition 2.1 that  $P$  is the product of its  $p$ -Sylow subgroups  $P_p$ . Moreover, each  $P_p$  is a pro- $p$  group and is therefore  $q$ -divisible for every  $q \in \mathbb{P} \setminus \{p\}$ . For each  $p \in \mathbb{P}$ , let  $Q_p$  be the subgroup of  $Q$  which is the pre-image of  $P_p$  under  $\pi$ .

Claim 8: If  $h, a, d \in Q$  with  $ha = d^n$  for some  $n \geq 1$  and  $a$  is elliptic, then  $|h| = n|d|$ .

Write  $C := \text{Min}(h) \cap \text{Min}(a)$ . Then  $C$  is nonempty by Claim 4. Since  $d^n$  stabilizes  $C$ , Claim 5 implies that it is semi-simple in  $C$  with translation length  $|d^n|_C = |d^n|$ . Thus,  $|d^n|_C = |d^n| = n|d|$  by Claim 6. Note also that  $ha$  is semi-simple in  $C$  with translation length  $|ha|_C = |h|$ . Therefore,  $|h| = |ha|_C = |d^n|_C = n|d|$ , as desired.

**Claim 9:** *Let  $h \in Q$  be hyperbolic. Suppose that  $ha_i = d_i^{n_i}$  for all  $i \geq 1$ , where  $a_i, d_i \in Q$ , each  $a_i$  is elliptic and where  $n_i \geq 1$ . Then the set  $\{n_i \mid i \geq 1\}$  is bounded.*

Indeed, by Claim 8, the sequence  $(d_i)$  of elements of  $Q$  is such that  $|d_i| = |h|/n_i > 0$ . The claim now follows from the second part of Proposition 2.4.

**Claim 10:** *Let  $p \in \mathbb{P}$ . Then all elements of  $Q_p$  are elliptic.*

Suppose for a contradiction there exists an  $h \in Q_p$  which is not elliptic, and is thus hyperbolic by Proposition 2.4. Let  $q \in \mathbb{P} \setminus \{p\}$ . Since  $P_p = \pi(Q_p)$  is  $q$ -divisible, there exists an  $h_q \in Q$  such that  $h_q^q Q^0 = hQ^0$ . Let  $a \in Q^0$  such that  $ha = h_q^q$ . By Proposition 2.2, since  $Q^0$  is compact and connected, it is divisible, and so  $a$  is elliptic by Claim 7. Since the set of natural prime numbers distinct from  $p$  is unbounded, the desired contradiction now comes from Claim 9.

Let now  $gQ^0 = (g_p)_{p \in \mathbb{P}}$  be the decomposition of  $\pi(g)$  in  $P = \prod_{p \in \mathbb{P}} P_p$  (that is, each  $g_p \in P_p$ ). Let  $q \in \mathbb{P}$ , and choose an  $a_q \in Q_p$  such that  $\pi(a_q) = g_q^{-1}$ . Then  $\pi(ga_q)$  has no component in the  $q$ -Sylow of  $P$ , and is therefore  $q$ -divisible in  $P$ . Hence, there exist an  $h_q \in Q$  and an  $a \in Q^0$  such that  $ga_q a = h_q^q$ . By Claim 10, we know that  $a_q$  is elliptic. But so is  $a$ , and hence the product  $a' := a_q a$  is also elliptic by Claim 4. Since  $q$  is an arbitrary prime, Claim 9 again yields the desired contradiction.  $\square$

**2.5. The Davis realization of a building.** We recall from [10] that any building  $\Delta$  admits a metric realization, denoted by  $|\Delta|_{\text{CAT}(0)}$ , which is a locally Euclidean  $\text{CAT}(0)$  cell complex with finitely many types of cells. Moreover any group of type-preserving automorphisms of  $\Delta$  acts in a canonical way by cellular isometries on  $|\Delta|_{\text{CAT}(0)}$ . Finally, the cell supporting any point of  $|\Delta|_{\text{CAT}(0)}$  determines a unique spherical residue of  $\Delta$ . In particular, an automorphism of  $\Delta$  which fixes a point in  $|\Delta|_{\text{CAT}(0)}$  must stabilize the corresponding spherical residue in  $\Delta$ .

Here is a reformulation of Theorem 1.

**Theorem 2.6.** *Let  $G$  be a compact group acting strongly transitively by type-preserving automorphisms on a thick building  $\Delta$ . Then,  $\Delta$  is spherical.*

*Proof.* Let  $(W, S)$  be the Coxeter system associated to  $\Delta$ , and let  $\Sigma$  be the fundamental apartment of  $\Delta$ . Then, the action of the stabilizer in  $G$  of  $\Sigma$  can be identified with the action of  $W$  on this apartment ([21, 2.8]).

**Claim 1:**  $|\Sigma|_{\text{CAT}(0)}$  is a closed convex subset of  $|\Delta|_{\text{CAT}(0)}$ .

A basic fact about buildings is the existence, for each pair  $(\Sigma, C)$  consisting of an apartment  $\Sigma$  and of a chamber  $C \in \Sigma$ , of a *retraction of  $\Delta$  onto  $\Sigma$  centered at  $C$* , that is, of a simplicial map  $\rho = \rho_{\Sigma, C}: \Delta \rightarrow \Sigma$  preserving minimal galleries from  $C$  and such that  $\rho|_{\Sigma} = \text{id}_{\Sigma}$ . The induced mapping  $\bar{\rho}: |\Delta|_{\text{CAT}(0)} \rightarrow |\Sigma|_{\text{CAT}(0)}$  then maps every geodesic segment of  $|\Delta|_{\text{CAT}(0)}$  onto a piecewise geodesic segment of  $|\Sigma|_{\text{CAT}(0)}$  of same length. In particular, the mapping  $\bar{\rho}$  is distance decreasing (see [10, Lemme 11.2]). Hence, if  $x$  and  $y$  are two points in  $|\Sigma|_{\text{CAT}(0)}$ , then the geodesic segment from  $x$  to  $y$  is entirely contained in  $|\Sigma|_{\text{CAT}(0)}$  since its image by  $\bar{\rho}$  is also a geodesic from  $x$  to  $y$ . This proves that  $|\Sigma|_{\text{CAT}(0)}$  is convex. To see it is closed, it suffices to note that it is complete as a metric space since it is precisely the Davis realization of the building  $\Sigma$ .

**Claim 2:** *If  $g \in G$  is elliptic in  $X = |\Delta|_{\text{CAT}(0)}$  and stabilizes  $|\Sigma|_{\text{CAT}(0)}$ , then  $g$  is also elliptic in  $|\Sigma|_{\text{CAT}(0)}$ .*

This follows from Claim 5 in the proof of Theorem 2.5.

Theorem 2.5 now implies that the induced action of  $W$  on  $|\Sigma|_{\text{CAT}(0)}$  is elliptic, that is, every  $w \in W$  is elliptic. Notice that the  $W$ -action on  $|\Sigma|_{\text{CAT}(0)}$  is proper, since by construction, it is cellular and the stabilizer of every point is a spherical (in particular finite) parabolic subgroup of  $W$ . Recalling now that every infinite finitely generated Coxeter group contains elements of infinite

order (in fact, so do all finitely generated infinite linear groups by a classical result of Schur [17]; in the special case of Coxeter groups, a direct argument may be found in [2, Proposition 2.74]), we deduce that  $W$  is finite. In other words  $\Delta$  is spherical.  $\square$

### 3. PROOF OF THEOREM 2

**3.1. Heuristic sketch.** Let  $G$  be a compact group possessing a split spherical BN-pair, and let  $\Delta$  be the associated building. We first establish Theorem 2 when  $G$  acts continuously on  $\Delta$ . In that case, 2-transitive actions (which are closely related to strongly transitive actions) of  $G$  on subspaces  $X$  of  $\Delta$  are easily seen to be possible only for finite  $X$ . The second step is then to show that the action of  $G$  on  $\Delta$  has to be continuous. This uses the fact that buildings arising from split spherical BN-pairs are Moufang (see Proposition 3.3 below).

**3.2. Continuous actions on buildings.** Recall that a topological space  $X$  is said to satisfy the  $T_1$  separation axiom when all its singletons are closed. The following is probably well-known.

**Lemma 3.1.** *Let  $G$  be a compact group. If  $G$  admits a continuous 2-transitive action on a  $T_1$  topological space  $X$ , then  $X$  is finite.*

*Proof.* Define  $Y := \{(x, y) \in X \times X \mid x \neq y\} \subset X \times X$ , and fix  $x, y \in X$  with  $x \neq y$ . Since the mapping  $\alpha_x: G \rightarrow X: g \mapsto g \cdot x$  is continuous, so is  $\alpha_x \times \alpha_y: G \rightarrow X \times X: g \mapsto (g \cdot x, g \cdot y)$ . By 2-transitivity, we get  $Y = (\alpha_x \times \alpha_y)(G)$ , and so  $Y$  is compact.

Note also that the mapping  $f: X \times X \rightarrow X \times X: (a, b) \mapsto (x, b)$  is continuous. Setting  $Z := X \setminus \{x\}$ , we then get  $Z \times \{x\} = f^{-1}(\{(x, x)\}) \cap Y$ , so that  $Z \times \{x\}$  is closed in  $Y$ , and hence compact. It follows that  $Z$  is compact, being the image of  $Z \times \{x\}$  by the projection on the first factor  $X \times X \rightarrow X$ , which is of course continuous.

In particular,  $Z$  is closed, and hence  $\{x\}$  is open. It follows that  $X$  is discrete, and therefore finite since  $X = \alpha_x(G)$  is compact.  $\square$

Let  $\Delta$  be a building of type  $(W, S)$ , and denote by  $\text{Ch } \Delta$  the set of its chambers. Consider the chamber system  $\Gamma$  of  $\Delta$ , which is the labelled graph with vertex set  $\text{Ch } \Delta$  and with an edge labelled by  $s \in S$  for each pair of  $s$ -adjacent chambers of  $\Delta$  (see [8, Ch.I Appendix D]). Let  $J \subset S$ . A  **$J$ -gallery** in  $\Gamma$  between two chambers  $x$  and  $y$  of  $\Delta$  is a sequence  $(x = x_0, x_1, \dots, x_l = y)$  of chambers of  $\Delta$  such that for each  $i \in \{1, \dots, l\}$ , there exists an  $s \in J$  such that  $x_{i-1}$  is  $s$ -adjacent to  $x_i$ . The natural number  $l$  is called the **length** of the gallery. A **minimal** gallery is a gallery of minimal length. The **distance** in  $\Delta$  between two chambers  $x, y \in \text{Ch } \Delta$  is the length of a minimal gallery joining  $x$  to  $y$ . The **diameter** of  $\Gamma$  is the supremum (in  $\mathbb{N} \cup \{\infty\}$ ) of the distances between its vertices.

Let  $J \subset S$ . The  **$J$ -residue**  $R = R_J(x)$  of some chamber  $x \in \text{Ch } \Delta$  is the set of chambers of  $\Delta$  which are connected to  $x$  by a  $J$ -gallery. When  $J$  has cardinality 1, we call  $R$  a **panel**.

In this paper, we will say that a group  $G$  acts **continuously** on  $\Delta$  if the stabilizers of the residues of  $\Delta$  are closed in  $G$ . Note that we can of course restrict our attention to the maximal proper residues, the others being obtained as intersections of those.

**Lemma 3.2.** *Let  $G$  be a compact group acting continuously and strongly transitively by type-preserving automorphisms on a spherical thick building  $\Delta$ . Then  $\Delta$  is finite.*

*Proof.* The stabilizer  $H$  in  $G$  of a panel  $P$  of  $\Delta$  is a closed and thus compact subgroup of  $G$ .

Claim 1:  *$H$  acts 2-transitively on  $P$ .*

Indeed, let  $C$  be a chamber of  $P$  and let  $B := \text{Stab}_G(C) \subset H$ . We first show that  $B$ , and thus also  $H$ , is transitive on the set  $\mathcal{C} = P \setminus \{C\}$ . Let  $C_1, C_2 \in \mathcal{C}$  and let  $\Sigma_1$  (respectively,  $\Sigma_2$ ) be an apartment containing  $C$  and  $C_1$  (respectively,  $C$  and  $C_2$ ). By strong transitivity,  $B$  is transitive on the set of apartments containing  $C$ , and so there exists a  $b \in B$  such that  $b\Sigma_1 = \Sigma_2$ . Hence  $bC_1 = C_2$ . It now remains to show that  $H$  is transitive on  $P$ . But if  $C_1, C_2 \in P$ , then since  $\Delta$  is thick, we may choose a chamber  $C$  in  $P$  different from  $C_1, C_2$ . The stabilizer  $B'$  of  $C$  in  $G$  is then contained in  $H$  and is transitive on  $P \setminus \{C\}$  by the previous argument.

Now, identifying  $\Delta$  with  $\Delta(G, B)$ , so that  $H = B \cup BsB$  for some generator  $s$  of the corresponding Weyl group, we get a 2-transitive, continuous action by left translation of the compact group  $H$  on the topological space  $H/B$ . Moreover, this space is  $T_1$  since  $B$  is closed in  $G$  by hypothesis. Lemma 3.1 then implies that  $P$  is finite. In other words, as  $P$  was arbitrary, the building  $\Delta$  is *locally finite*, that is, every panel is finite. The following observation now allows us to conclude:

**Claim 2:** *Every locally finite spherical building is finite.*

Indeed, let  $\Gamma = \text{Ch } \Delta$  be the graph whose vertices are the chambers of  $\Delta$ , and such that two chambers of  $\Delta$  are adjacent if they share a common panel. Since  $\Delta$  is locally finite, so is  $\Gamma$ . Hence, fixing a vertex  $x \in \Gamma$ , each ball in  $\Gamma$  centered at  $x$  with radius  $n$  ( $n \in \mathbb{N}$ ) possesses a finite number of vertices. Moreover, as  $\Delta$  is spherical, the diameter of  $\Delta$  is finite ([8, Ch.IV, 3]), and hence the diameter of  $\Gamma$  is also finite. Thus  $\Gamma$  is contained in such a ball, and is therefore finite.  $\square$

**3.3. Moufang buildings.** Let  $\Delta = \Delta(G, B)$  be the building associated to a split spherical BN-pair  $(B = T \times U, N)$  of type  $(W, S)$ . It is well-known (see the main result of [11]) that the existence of a splitting for the above BN-pair is equivalent to the fact that the building  $\Delta$  enjoys the Moufang property, as defined in [21, Chapter 11].

Two chambers  $x, y \in \text{Ch } \Delta$  are called **opposite** if they are at maximal distance in the chamber system of  $\Delta$ . Similarly, one can define *opposite residues* (see for instance [2, 5.7]). The set of chambers (respectively, residues) of  $\Delta$  which are opposite to a given chamber  $C$  (respectively, residue  $R$ ) will be denoted by  $C^{\text{op}}$  (respectively,  $R^{\text{op}}$ ).

**Proposition 3.3.** *Let  $P = BW_J B$  be a proper standard parabolic subgroup of  $\Delta = \Delta(G, B)$  for some proper subset  $J$  of  $S$ , let  $C$  be the fundamental chamber (i.e. the unique chamber fixed by  $B$ ) and let  $R$  be the unique  $J$ -residue containing  $C$ . Define the subgroup  $V := \bigcap_{p \in P} pUp^{-1}$  of  $G$ . Then  $V$  acts simply transitively on  $R^{\text{op}}$ .*

*Proof.* Let  $\Sigma$  be an apartment containing  $C$ . By [21, 9.11], there exists a minimal gallery  $\gamma_{R'}$ , one for each residue  $R' \in R^{\text{op}}$ , beginning at  $C$  and ending at a chamber  $C'$  in  $R'$  such that the type of  $\gamma_{R'}$  is independent of the choice of  $R'$  and  $C = \text{proj}_R C'$ . Let  $R' \in R^{\text{op}}$  be the unique residue of  $\Sigma$  opposite  $R$  and let  $C'$  be the last chamber of  $\gamma_{R'}$ . Let also  $\alpha$  be a root of  $\Sigma$  containing  $C$  but not  $C'$ . By [21, 8.21],  $R \cap \Sigma \subset \alpha$ . By [21, 9.7], therefore,  $R$  is fixed pointwise by the root group  $U_\alpha$ . Since  $P$  maps  $R$  to itself, we have  $C \in R \subset \alpha^P$  and hence  $p^{-1}U_\alpha p \subset U$  for all  $p \in P$  by the definition of root subgroups (see [21, 11.1]) and the fact that the ‘radical’  $U$  does not depend on the choice of the apartment  $\Sigma$  (see [21, Proposition 11.11(iii)]). Thus  $U_\alpha \subset V$ . Now, as in [2, 7.67], one shows that the subgroup of  $V$  generated by all  $U_\alpha$ ’s of the latter form acts transitively on the set  $\{\gamma_{R'} \mid R' \in R^{\text{op}}\}$ , and hence also on  $R^{\text{op}}$ .

Suppose  $h \in V$  maps  $R' \in R^{\text{op}}$  to itself. Then  $h$  acts trivially on  $R$ . Since the restriction of  $\text{proj}_{R'}$  to  $R$  is a bijection from  $R$  to  $R'$  (by [21, 9.11] again), it follows that  $h$  acts trivially on  $R'$ . By [21, 9.8], therefore,  $h$  fixes two opposite chambers of  $\Sigma$  and hence  $h$  fixes  $\Sigma$ . By [21, 9.7] again, we conclude that  $h = 1$ .  $\square$

In particular, we have the following (compare [8, Ch.IV, 5]).

**Lemma 3.4.** *Let  $C$  be the fundamental chamber of  $\Delta$ . Then  $U$  acts simply transitively on  $C^{\text{op}}$ . Equivalently,  $U$  acts simply transitively on the set of apartments containing  $C$ .*

**Lemma 3.5.** *Let  $P = BW_J B$  be a proper standard parabolic subgroup of  $\Delta = \Delta(G, B)$  for some proper subset  $J$  of  $S$ , let  $C$  be the fundamental chamber and let  $R$  be the unique  $J$ -residue containing  $C$ . Then there exist two chambers in  $C^{\text{op}}$  which are opposite to one another. In particular,  $|R^{\text{op}}| \geq 2$ .*

*Proof.* The first assertion holds by [2, Proposition 4.104] and the second follows since no proper residue contains two opposite chambers.  $\square$

We are now ready to complete the proof of Theorem 2.

**Theorem 3.6.** *Let  $G$  be a compact topological group possessing a spherical split BN-pair  $(B = T \times U, N)$ . Then the associated building is finite.*

*Proof.* Let  $\Delta = \Delta(G, B)$  be the building associated to  $(B, N)$ , and let  $(W, S)$  be the corresponding Coxeter system.

We start with some basic observations in the case  $(W, S)$  is not irreducible. Suppose thus that  $S$  decomposes as  $S = S_1 \amalg S_2$  with  $s_1 s_2 = s_2 s_1$  for all  $s_1 \in S_1$  and  $s_2 \in S_2$ . Then  $W$  splits as a direct product  $W \cong W_1 \times W_2$ , where  $W_i = \langle S_i \rangle$ , and the building  $\Delta$  decomposes canonically as a product  $\Delta = \Delta_1 \times \Delta_2$  of buildings of type  $(W_1, S_1)$  and  $(W_2, S_2)$  respectively (see [21, Proposition 7.33]).

In particular, we obtain induced actions of  $G$  on both  $\Delta_1$  and  $\Delta_2$ , which are obviously strongly transitive. The corresponding BN-pairs for  $G$  may be described as follows. Since each  $s \in S$  can be written as a coset  $nT \in N/T = W$ , we may choose, for  $i = 1, 2$ , a set  $\overline{N}_i$  of representatives in  $N$  for the elements of  $S_i$ . For each  $i = 1, 2$ , consider now the subgroup  $N_i$  of  $N$  generated by  $\overline{N}_i$  and  $T$ , and set  $B_i := \langle B \cup N_{3-i} \rangle = BN_{3-i}B \leq G$ . Then  $(B_i, N_i)$  is a spherical BN-pair for  $G$ , and the associated building is nothing but  $\Delta_i = \Delta(G, B_i)$ .

We claim that the BN-pair  $(B_i, N_i)$  is split. This follows readily from the aforementioned equivalence between splittings of BN-pairs and the Moufang property for the associated buildings. More precisely, consider the group  $U_i = \bigcap_{g \in B_i} gUg^{-1}$  which is the kernel of the  $U$ -action on  $\Delta_{3-i}$ . Then  $U_i$  acts sharply transitively on the chambers of  $\Delta_i$  which are opposite the standard chamber  $C$ , which by definition is the unique chamber fixed by  $B_i$ . Therefore we have  $B_i \cong T_i \rtimes U_i$ , where  $T_i = \bigcap_{w \in W_i} wB_iw^{-1}$ , and  $U_i$  induces a splitting of the BN-pair  $(B_i, N_i)$  as claimed.

This shows that the given split BN-pair for  $G$  yields various split BN-pairs for  $G$  corresponding to the various irreducible components of  $\Delta$ . Since  $\text{Ch } \Delta$  is naturally in one-to-one correspondence with the Cartesian product  $\text{Ch } \Delta_1 \times \cdots \times \text{Ch } \Delta_n$  of the chamber sets of the various irreducible components of  $\Delta$ , the desired finiteness result readily follows provided we establish it for each irreducible BN-pair  $(B_i, N_i)$  as above. In other words, there is no loss of generality in assuming that the building  $\Delta$  is irreducible. We adopt henceforth this additional assumption.

Let now  $\mathcal{P}$  denote the set of maximal proper standard parabolic subgroups of  $G$ . Pick any  $P \in \mathcal{P}$ . Thus  $P$  is of the form  $P = BW_JB$  for some maximal subset  $J \subsetneq S$ , where  $W_J = \langle J \rangle$ . In particular,  $P$  is a maximal subgroup of  $G$  (see [2, Lemma 6.43(1)]). Define the normal subgroup

$$V := \bigcap_{p \in P} pUp^{-1} \trianglelefteq P$$

of  $P$ . As  $V$  is contained in  $U$ , it is also nilpotent. Moreover,  $V$  acts faithfully on  $\Delta$ . Indeed, the kernel  $\ker(G \curvearrowright \Delta)$  of the action of  $G$  on  $\Delta$  is obviously contained in the stabilizer of the chambers of the fundamental apartment  $\Sigma$ , that is, in  $\bigcap_{w \in W} wBw^{-1} = T$ , and so

$$V \cap \ker(G \curvearrowright \Delta) \subseteq U \cap T = \{1\}.$$

Now, since  $V$  is normal in  $P$ , we have  $P \subseteq \mathcal{N}_G(V)$ . Moreover, as the conjugation automorphism  $\kappa_g: G \rightarrow G: x \mapsto gxg^{-1}$  is continuous, we get  $\mathcal{N}_G(\overline{V}) \supseteq \mathcal{N}_G(V)$  and so  $\mathcal{N}_G(\overline{V}) \supseteq P$ . Hence, by maximality of  $P$ , we obtain that either  $\mathcal{N}_G(\overline{V}) = P$  or  $\mathcal{N}_G(\overline{V}) = G$ .

**Claim:**  $\mathcal{N}_G(\overline{V}) = P$  for all  $P \in \mathcal{P}$ .

Assume for a contradiction that  $\mathcal{N}_G(\overline{V}) = G$  for some  $P \in \mathcal{P}$ . In other words,  $\overline{V} \triangleleft G$ . In particular, the center  $\mathcal{Z}(\overline{V}) \subseteq \overline{V}$  of  $\overline{V}$  is also a normal subgroup of  $G$ . Moreover,  $V$  is nontrivial since, by Proposition 3.3, it acts transitively on  $R^{\text{op}}$  and since  $|R^{\text{op}}| \geq 2$  by Lemma 3.5. As  $V$  is nilpotent, this implies that  $\mathcal{Z}(V)$  is also nontrivial.

Now, using again the continuity of the conjugation automorphism  $\kappa_h$  (for  $h \in G$ ), we see that  $\mathcal{Z}(V) = \mathcal{Z}_G(V) \cap V$  is contained in  $\mathcal{Z}(\overline{V}) = \mathcal{Z}_G(\overline{V}) \cap \overline{V}$ . Moreover, as  $V$  acts faithfully on  $\Delta$ , so does  $\mathcal{Z}(V)$ . This implies in particular that  $\mathcal{Z}(V)$ , and thus also  $\mathcal{Z}(\overline{V})$ , act nontrivially on  $\Delta$ .

Tits' transitivity Lemma (see [8, Lemma 6.61]) then guarantees that the group  $\mathcal{Z}(\overline{V})$  is transitive on the chambers of  $\Delta$ . In fact, this action is even simply transitive. Indeed, the stabilizers in  $\mathcal{Z}(\overline{V})$  of the chambers of  $\Delta$  are all conjugate by transitivity. They are thus all equal since  $\mathcal{Z}(\overline{V})$  is Abelian, and are therefore contained in the kernel  $\ker(G \curvearrowright \Delta)$  of the action of  $G$  on  $\Delta$ . Since  $\mathcal{Z}(V) \subseteq \mathcal{Z}(\overline{V})$ , this implies that the action of  $\mathcal{Z}(V)$  on  $\text{Ch } \Delta$  is free. But since  $\mathcal{Z}(V) \subseteq V \subseteq U \subseteq B$ , and as  $B$  stabilizes the fundamental chamber, it follows that  $\mathcal{Z}(V)$  acts trivially on  $\Delta$ . This contradiction establishes the Claim.



Since the normalizer of a closed subgroup is closed, we deduce from the Claim that every  $P \in \mathcal{P}$  is closed. But this means that  $G$  acts continuously on  $\Delta$ , and so Lemma 3.2 ensures that  $\Delta$  is finite, as desired.  $\square$

#### 4. PROOF OF THEOREM 4

Let  $k$  be a perfect field and let  $K = \overline{k}$  be its algebraic closure. In what follows, we identify an algebraic  $k$ -group  $G$  with its group of  $K$ -rational points.

The main tool for the proof of Theorem 4 is the following characterization of anisotropy, due to Borel and Tits (see [3]).

**Proposition 4.1.** *Let  $G$  be a reductive algebraic  $k$ -group and let  $U$  be a unipotent  $k$ -subgroup of  $G$ . If  $k$  is perfect, then there exists a parabolic  $k$ -subgroup  $P$  of  $G$  whose unipotent radical  $R_u(P)$  contains  $U$ .*

In particular, if  $G$  is anisotropic over  $k$ , then  $U$  must be trivial.

*Proof of Theorem 4.* Suppose for a contradiction that the split spherical BN-pair  $(B, N)$  for the reductive  $k$ -group  $G$  is such that  $B$  has infinite index in  $G(k)$ . Let  $\Delta = \Delta(G(k), B)$  be the associated building, and let  $W$  be the corresponding (finite) Weyl group. Also, denote by  $\overline{B}$  the Zariski closure of  $B$  in  $G$ .

The Bruhat decomposition for  $G$  yields  $G = \coprod_{w \in W} BwB$ . Since  $G(k)$  is Zariski dense in  $G$  by [4, 18.3], we have

$$G = \overline{G(k)} = \overline{\coprod_{w \in W} BwB} \subseteq \coprod_{w \in W} \overline{BwB}.$$

As  $G$  is connected, it cannot be written as a finite union of closed subsets in a nontrivial way. Therefore, we deduce that  $BwB$  is dense in  $G$  for some  $w \in W$ . In particular, so is  $\overline{BwB}$ .

Let now  $A := (\overline{B})^0$  be the identity component of  $\overline{B}$ . Since  $A$  has finite index in  $\overline{B}$ , it follows that  $\overline{BwB}$  is a finite union of double cosets modulo  $A$ . As before, this implies that some double coset of the form  $AzA$  is dense in  $G$ .

Claim:  $\overline{B} \neq G$ .

Indeed, let  $U$  be the nilpotent normal subgroup of  $B$  arising from the splitting of the BN-pair, and suppose for a contradiction that  $B$  is dense in  $G$ . Then the Zariski closure  $\overline{U}$  of  $U$  in  $G$  is a nilpotent normal subgroup of  $\overline{B} = G$  ([4, 2.1]). Its identity component  $\overline{U}^0$  is thus contained in the radical of  $G$ , which coincides with the connected center  $\mathcal{Z}(G)^0$  ([4, 11.21]). Hence, since  $\overline{U}^0$  has finite index in  $\overline{U}$ , we get

$$[U : U \cap \mathcal{Z}(G)] \leq [U : U \cap \overline{U}^0] = [U\overline{U}^0 : \overline{U}^0] \leq [\overline{U} : \overline{U}^0] < \infty.$$

Now, if  $u \in U \cap \mathcal{Z}(G)$ , then  $u$  acts trivially on  $\Delta$  since for any chamber  $gB$ , we have  $ugB = guB = gB$ . As  $U$  acts simply transitively on  $C^{\text{op}}$  by Lemma 3.4, where  $C = 1_G B$  is the fundamental chamber of  $\Delta$ , this implies that  $u = 1$ : otherwise,  $\Delta$  would contain only one apartment, so that  $[G(k) : B] < \infty$ , a contradiction. So  $U \cap \mathcal{Z}(G) = \{1\}$  and therefore  $U$  is finite. Using again the sharp transitivity of  $U$  on  $C^{\text{op}}$ , we deduce that  $\Delta$  is the reunion of finitely many apartments, hence is finite, contradicting once more our initial hypothesis. The claim stands proven.

In particular  $A$  is a proper closed subgroup of  $G$  such that  $AzA$  is dense in  $G$  for some  $z \in G$ . The main result of [9] now implies that  $A$  is not reductive, *i.e.* the unipotent radical  $R_u(A)$  is nontrivial. Moreover, since  $B$  is contained in  $G(k)$  and is dense in  $\overline{B}$ , we know that  $\overline{B}$  is defined on  $k$  ([4, AG.14.4]). Hence,  $A$  is also  $k$ -defined ([4, 1.2]), and so is  $R_u(A)$  since  $k$  is perfect ([18, 12.1.7(d)]). Thus  $R_u(A)$  is a nontrivial unipotent  $k$ -subgroup of  $G$ . As we observed following Proposition 4.1, this contradicts the assumption that  $G$  is anisotropic over  $k$ .  $\square$

## REFERENCES

- [1] Peter Abramenko and Kenneth S. Brown. Transitivity properties for group actions on buildings. *J. Group Theory*, 10(3):267–277, 2007. 1
- [2] Peter Abramenko and Kenneth S. Brown. *Buildings*, volume 248 of *Graduate Texts in Mathematics*. Springer, New York, 2008. Theory and applications. 1, 6, 7, 8
- [3] Armand Borel and Jacques Tits. Éléments unipotents et sous-groupes paraboliques de groupes réductifs. I. *Invent. Math.*, 12:95–104, 1971. 9
- [4] Armand Borel. *Linear algebraic groups*, volume 126 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1991. 1, 9
- [5] Armand Borel and Jacques Tits. Groupes réductifs. *Inst. Hautes Études Sci. Publ. Math.*, (27):55–150, 1965. 1
- [6] Martin R. Bridson and André Haefliger. *Metric spaces of non-positive curvature*, volume 319 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin, 1999. 3, 4
- [7] Martin R. Bridson. On the semisimplicity of polyhedral isometries. *Proc. Amer. Math. Soc.*, 127(7):2143–2146, 1999. 3
- [8] Kenneth S. Brown. *Buildings*. Springer Monographs in Mathematics. Springer-Verlag, New York, 1998. Reprint of the 1989 original. 6, 7, 8
- [9] Jonathan Brundan. Double coset density in reductive algebraic groups. *J. Algebra*, 177(3):755–767, 1995. 9
- [10] Michael W. Davis. Buildings are CAT(0). In *Geometry and cohomology in group theory (Durham, 1994)*, volume 252 of *London Math. Soc. Lecture Note Ser.*, pages 108–123. Cambridge Univ. Press, Cambridge, 1998. 5
- [11] Tom De Medts, Fabienne Haot, Katrin Tent, and Hendrik Van Maldeghem. Split *BN*-pairs of rank at least 2 and the uniqueness of splittings. *J. Group Theory*, 8(1):1–10, 2005. 7
- [12] Jan Mycielski. Some properties of connected compact groups. *Colloq. Math.*, 5:162–166, 1958. 3
- [13] Gopal Prasad. Elementary proof of a theorem of Bruhat-Tits-Rousseau and of a theorem of Tits. *Bull. Soc. Math. France*, 110(2):197–202, 1982. 1
- [14] Andrei S. Rapinchuk. Algebraic and Abstract Simple Groups: Old and New. *Prepublication*, pages 55–150, 2003. 2
- [15] Andrei S. Rapinchuk, Yoav Segev, and Gary M. Seitz. Finite quotients of the multiplicative group of a finite dimensional division algebra are solvable. *J. Amer. Math. Soc.*, 15(4):929–978 (electronic), 2002. 2
- [16] Luis Ribes and Pavel Zalesskii. *Profinite groups*, volume 40 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*. Springer-Verlag, Berlin, 2000. 3
- [17] Issai Schur. Über Gruppen periodischer linearer Substitutionen. *Berl. Ber.*, 1911:619–627, 1911. 5
- [18] Tonny A. Springer. *Linear algebraic groups*, volume 9 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, second edition, 1998. 9
- [19] Katrin Tent. A short proof that root groups are nilpotent. *J. Algebra*, 277(2):765–768, 2004. 1
- [20] Jacques Tits. Algebraic and abstract simple groups. *Ann. of Math. (2)*, 80:313–329, 1964. 1
- [21] Richard M. Weiss. *The structure of spherical buildings*. Princeton University Press, Princeton, NJ, 2003. 5, 7, 8

UNIVERSITÉ CATHOLIQUE DE LOUVAIN, CHEMIN DU CYCLOTRON 2, 1348 LOUVAIN-LA-NEUVE, BELGIUM  
*E-mail address:* pierre-emmanuel.caprace@uclouvain.be

UNIVERSITÉ LIBRE DE BRUXELLES, BOULEVARD DU TRIOMPHE, 1050 BRUXELLES, BELGIUM  
*E-mail address:* tmarquis@ulb.ac.be