

## Multi-scale modelling of coastal, shelf and global ocean dynamics

**Eric Deleersnijder**  
**Vincent Legat**  
**Pierre Lermusiaux**

Editorial for a special issue in *Ocean Dynamics*  
November, 2010

Methods for widening the range of resolved scales (i.e. performing multi-scale simulations) in ocean sciences and engineering are developing rapidly, now allowing multi-scale ocean dynamics studies. Having recourse to grid nesting has been, and still is, a popular method for increasing marine models' resolution when and where needed, and for easily allowing the use of different dynamics at different resolution. However, this is not the only way to achieve this goal. Various techniques for modifying locally the grid resolution or dealing with complex-geometry domains are available. For instance, composite, structured grids and unstructured meshes offer an almost infinite geometrical flexibility.

This special issue focuses on such multi-scale modelling of coastal, shelf and global ocean dynamics, including the development of new methodologies and schemes and their applications to ocean process studies. Several articles focus on numerical aspects of unstructured-mesh space discretisation. Danilov shows that the noise developing on triangular meshes on which the location of the variables is inspired by Arakawa's C-grid is largest for regimes close to geostrophic balance. The noise can be reduced by specific operators, but cannot be entirely suppressed, *making the triangular C-grid a suboptimal choice for large-scale ocean modelling*. Then, the companion articles of Blaise et al. and Comblen et al. describe the space and time discretisation of a three-dimensional, baroclinic, finite-element model based on the Discontinuous Galerkin (DG) technique. This is a significant step forward in the field of finite-element ocean modelling, though this model cannot yet be regarded as suitable

---

E. Deleersnijder<sup>a,b,c,d</sup> . V. Legat<sup>a,b,c</sup>  
Université catholique de Louvain

<sup>a</sup> Institute of Mechanics, Materials and Civil Engineering (IMMC),

<sup>b</sup> Georges Lemaître Centre for Earth and Climate Research (TECLIM),

<sup>c</sup> Centre for systems engineering and applied mechanics (CESAME),

<sup>d</sup> Earth and Life Institute (ELI),

Euler, 4 Avenue G. Lemaître, B-1348 Louvain-la-Neuve, Belgium

P. Lermusiaux

Massachusetts Institute of Technology, Department of Mechanical Engineering, Room 5-207B,  
77 Massachusetts Avenue, Cambridge MA 02139-4307, USA

for tackling realistic applications. Ueckermann and Lermusiaux also consider DG finite element techniques, focusing on biological-physical dynamics in regions with complex bathymetric features. They compare low- to high-order discretisations, both in time and space, for regimes in which biology dominates, advection dominates, or terms are balanced. They find that higher order schemes on relatively coarse grids generally perform better than low-order schemes on fine grids. Kleptsova et al. assess various advection schemes for z-coordinate, three-dimensional models in which flooding and drying is taken into account. In this study, the ability to conserve momentum is regarded as the main criterion for selecting a suitable method. On the other hand, Maßmann assesses automatic differentiation for obtaining the adjoint of an unstructured-mesh, tidal model of the European Continental Shelf.

Two articles deal with grid nesting. Nash and Hartnett introduce a flooding and drying method that can be used in structured, nested-grid systems. This can be regarded as an alternative to flooding and drying techniques that are being developed for unstructured-mesh models (e.g. Kärnä et al. 2010). Then, Haley and Lermusiaux derive conservative time-dependent structured finite-volume discretisations and implicit two-way embedded schemes for primitive-equations with the intent to resolve tidal-to-mesoscale processes over large multiresolution telescoping domains with complex geometries including shallow seas with strong tides, steep shelfbreaks, and deep ocean interactions. The authors present realistic simulations with data assimilation in three regions with diverse dynamics and show that their developments enhance the predictive capability, leading to better match with ocean data.

Various multi-scale, realistic simulations are presented. Using a finite-element ice model and a slab ocean as in Lietaer et al (2008), Terwisscha van Scheltinga et al. model the Canadian Arctic Archipelago, focusing on the pathways for freshwater and sea-ice transport from the Arctic Ocean to the Labrador Sea and the Atlantic Ocean. The unstructured mesh can represent the complex geometry and narrow straits at high resolution, and allows improving transports of water masses and sea ice. Walters et al. have recourse to an unstructured mesh model to study tides and current in Greater Cook Strait (New Zealand). They identify the mechanisms causing residual currents. By means of the unstructured-mesh Finite Volume Coastal Ocean Model (FVCOM), Wang et al. study the hydrodynamics of the Bohai Sea. Xu et al. simulate coastal and urban inundation due to storm surges along U.S. East and Gulf Coasts. A sensitivity analysis reveals the importance of precise topographic data and the need for a bottom drag coefficient accounting for the presence of mangroves. Finally, Yang and Khangaonkar resort to FVCOM to simulate the three-dimensional circulation of Puget Sound, a large complex estuary system in the Pacific Northwest coastal ocean, including variable forcing from tides, the atmosphere, and river inflows. Comparisons of model estimates with measurements for tidal elevation, velocity, temperature, and salinity are deemed to be promising, from larger-scale circulation features to nearshore tide flats.

This special issue suggests that numerical techniques for multi-scale space discretisation are progressively becoming mature. One direction for future progress lies in the improvement of time discretisation methods for the new generation models, so that they can successfully compete with finite-difference, structured-mesh models based on (almost) constant resolution grids that have been developed and used over the past 40 years (e.g. Griffies et al. 2009).

**Acknowledgements** The guest-editors are indebted to the authors and anonymous reviewers for their contributions to this special issue. Eric Deleersnijder is a research associate with the Belgian Fund for Scientific Research (F.R.S.-FNRS). The contributions of Eric Deleersnijder and Vincent Legat to this special issue were achieved in the framework of the project *Taking up the challenges of multi-scale marine modelling*, which is funded by the Communauté Française de Belgique under contract ARC 10/15-028, and the Interuniversity Attraction Pole Timothy ([www.climate.be/timothy](http://www.climate.be/timothy)). The latter is funded by BELSPO ([www.belspo.be](http://www.belspo.be)) under contract IAP6.13. Pierre Lermusiaux is very grateful to the Office of Naval Research for support under grants N00014-07-1-1061 and N00014-08-1-1097 to the Massachusetts Institute of Technology. Some of the articles of this special issue derive from presentations made at the 8th International Workshop on Unstructured Mesh Numerical Modelling of Coastal, Shelf and Ocean Flows (16-18 September 2009, Louvain-la-Neuve, Belgium, [www.uclouvain.be/umm2009](http://www.uclouvain.be/umm2009)), for which the F.R.S.-FNRS and the contract ARC 04/09-316 (Communauté Française de Belgique) provided financial support.

## References

- . Blaise S., R. Comblen, V. Legat, J.-F. Remacle, E. Deleersnijder and J. Lambrechts: A discontinuous finite element baroclinic marine model on unstructured prismatic meshes. Part I: space discretization, *Ocean Dynamics* (this issue)
- . Comblen R., S. Blaise, V. Legat, J.-F. Remacle, E. Deleersnijder and J. Lambrechts: A discontinuous finite element baroclinic marine model on unstructured prismatic meshes. Part II: implicit/explicit time discretization, *Ocean Dynamics* (this issue)
- . Danilov S.. On utility of triangular C-grid type discretization for numerical modeling of large-scale ocean, *Ocean Dynamics* (this issue)
- . Griffies S.M., A.J. Adcroft, H. Banks, C.W. Böning, E.P. Chassignet, G. Danabasoglu, S. Danilov, E. Deleersnijder, H. Drange, M.H. England, B. Fox-Kemper, R. Gerdes, A. Gnanadesikan, R.J. Greatbatch, R.W. Hallberg, E. Hanert, M.J. Harrison, S.A. Legg, C.M. Little, G. Madec, S. Marsland, M. Nikurashin, A. Pirani, H.L. Simmons, J. Schröter, B.L. Samuels, A.-M. Treguier, J.R. Toggweiler, H. Tsujino, G.K. Vallis and L. White, 2009, Problems and prospects in large-scale ocean circulation models, in: J. Hall, D.E. Harrison and D. Stammer (Eds.), *Proceedings of the OceanObs'09 Conference: Sustained Ocean Observations and Information for Society* (Venice, Italy, 21-25 September 2009), ESA Publication WPP-306
- . Haley P.J., Jr and P.F.J. Lermusiaux, Multiscale two-way embedding schemes for free-surface primitive-equations in the *Multidisciplinary Simulation, Estimation and Assimilation System*, *Ocean Dynamics* (this issue)
- . Kärnä T., B. de Brye, O. Gourgue, J. Lambrechts, R. Comblen, V. Legat and E. Deleersnijder, 2010, A fully implicit wetting-drying method for DG-FEM shallow water models, with an application to the Scheldt Estuary, *Computer Methods in Applied Mechanics and Engineering* (in press), doi: 10.1016/j.cma.2010.07.001
- . Kleptsova O., G.S. Stelling and J.D. Pietrzak, An accurate momentum advection scheme for a z-level coordinate model, *Ocean Dynamics* (this issue)
- . Lietaer O., T. Fichefet and V. Legat, 2008, The effects of resolving the Canadian Arctic Archipelago in a finite element sea ice model, *Ocean Modelling*, 24, 140-152
- . Maßmann S., Sensitivities of an adjoint, unstructured mesh, tidal model on the European Continental Shelf, *Ocean Dynamics* (this issue)
- . Nash S. and M. Hartnett, Nested circulation modelling of inter-tidal zones: details of a nesting approach incorporating moving boundaries, *Ocean Dynamics* (this issue)
- . Terwisscha van Scheltinga A.D., P.G. Myers and J.D. Pietrzak. A finite element sea ice model of the Canadian Arctic Archipelago, *Ocean Dynamics* (this issue)
- . Ueckermann M.P. and P.F.J. Lermusiaux, High order schemes for 2D unsteady biogeochemical ocean models, *Ocean Dynamics* (this issue)
- . Walters R.A., P.A. Gillibrand, R.G. Bell and E.M. Lane, A study of tides and currents in Cook Strait, New Zealand, *Ocean Dynamic* (this issue)
- . Wang J., Y. Shen and Y. Guo, Seasonal circulation and influence factors of the Bohai Sea: a numerical study based on Lagrangian particle tracking method, *Ocean Dynamics* (this issue)
- . Xu H., K. Zhang, J. Shen and Y. Li, Storm surge simulation along the U.S. East and Gulf Coasts using a multi-scale numerical model approach, *Ocean Dynamics* (this issue)

- . Yang Z. and T. Khangaonkar, Multi-scale modeling of Puget Sound using an unstructured-grid coastal ocean model: from tide flats to estuaries and coastal waters, *Ocean Dynamics* (this issue)